



Entomofauna

ZEITSCHRIFT FÜR ENTOMOLOGIE

Band 37, Heft 23: 389-400

ISSN 0250-4413

Ansfelden, 4. Januar 2016

Study of the fruit flies (Diptera: Tephritidae) as biocontrol agents of asteraceus plants in Saqez region (Kurdestan province)

Mozhgan ZAMANI & Samad KHAGHANINIA

Abstract

Tephritidae is one of the most important families of the order Diptera. Larvae of this family are often associated with thistle of various plants specially family Asteraceae. The larvae of some species feeding of inedible asteraceus plants to grazing animals, reduce seed amount and cause stable control of them in grasslands. In this study, 13 species belonging to 9 genera of the family Tephritidae from 10 species belonging to 6 genera of the host plants were collected and identified during 2011-2013. Four species, *Onopordum acanthium* (LINNÉ, 1753), *Onopordum illyricum*, *Acroptilon repens* (LINNÉ, 1837) and *Echinops spinosissimus* TURRA, 1765, are recorded as the host plants of these flies for the first time from Iran. *Acanthiophilus helianthi* Rossi, 1794 from *Onopordum acanthium* and *Acroptilon repens*, *Tephritis postica* LOEW, 1844 and *Terellia gynae-cochroma* HERING, 1937 from *Onopordum illyricum* as well as *Tephritis myia lauta* LOEW, 1869 from *Echinops spinosissimus* were obtained by rearing in the laboratory.

Key words: Iran, Saqez, Asteraceae, Host plant association, Fruit flies.

Zusammenfassung

Die Tephritiden bilden eine der wichtigsten Familien der Ordnung Diptera (Zweiflügler). Die Larven dieser Familie sind oft vergemeinschaftet mit den Samenköpfen verschiedener Pflanzen, hauptsächlich der Familie Asteraceae (Korbblütler). Sofern die Larven sich von Korbblütlern ernähren, welche von grasesenden Nutzieren gemieden werden, sorgen sie für eine Reduzierung der Samenmenge dieser Pfanzen und damit für eine andauernde Erhaltung der Grasweide.

Für diese Studie wurden in den Jahren 2011–2013 13 Arten von Tephritiden aus 9 Gattungen dieser Familie gesammelt und identifiziert; diese 13 Arten lebten auf 10 Arten aus 6 Gattungen von Wirtspflanzen. Von diesen Wirtspflanzen wurden 4 Arten erstmals aus dem Iran gemeldet, und zwar *Onopordum acanthium* (LINNÉ, 1753), *Onopordum illyricum*, *Acroptilon repens* (LINNÉ, 1837) sowie *Echinops spinosissimus* TURRA, 1765. Folgende Larvenarten wurden aus Aufzucht im Laboratorium gewonnen: *Acanthiophilus helianthi* ROSSI, 1794 vom *Onopordum acanthium* und vom *Acroptilon repens*; *Tephritis postica* LOEW, 1844 und *Terellia gynaecochroma* HERING, 1937 beide vom *Onopordum illyricum*; *Tephritisomyia lauta* LOEW, 1869, vom *Echinops spinosissimus*.

Introduction

The Tephritidae, true fruit flies, with more than 4500 described species is a family of attractive picture-winged flies. The taxonomic diversity of this single family of two-winged flies is far greater than that of mammals (NORRBOM 2004). Fruit flies are distributed throughout the temperate and tropical areas of the world, being absent only from the high arctic and antarctic (THOMPSON 1998). This family consist of small to medium flies in 2–12 mm long, fronto-orbital plate usually bears one or more frontal bristles, antennae with bare or plumose arista, wings usually with a pattern consisting of brown strips and spots, costal vein with two interruptions (WHITE & ELSON-HARRIS 1992). They are almost all phytophagous and include numerous pests of fruit and vegetable crops as well as useful species to the biocontrol of weeds. Most Tephritidae larvae associated with thistle of plants particularly of the family Asteraceae. Family Asteraceae, usually known as Compositae or Sunflower, is one of the biggest families of the Angiosperms and the main features of this family is the type of their thistle flower. A few inflorescences of this family are compound and have environmental-economic importance and found in the most habitats. These plants are distributed in dry or semidry area (DLUSSKII et al. 2004).

Sometimes tephritids act as important biological control agents of asteraceus plants by reducing seed production and operation but some other attack leaves plant and cause leaf miner (KAPOOR et al. 1980; WHITE 1988). Some species like *Urophora stylata* (FABRICIUS) and *Chaetostomella cylindrical* (ROBINEAU-DESOUDY) are important agents in biological control programs against weeds such as *Cirsium vulgare* (SAVI) (WHITE & ELSON-HARRIS, 1992). Also *Urophora sirunaseva* (HERING) was used to control of some roughage species of Asteraceae (WHITE & ELSON-HARRIS 1992; WOODS et al. 2008).

Recently some studies have been done about identification of the fruit flies and their host plants in Iran (MOHAMMADZADE NAMIN et al. 2010; KARIMPOUR 2011). MOHAMMADZADE NAMIN & NOZARI (2011), introduced *Centaurea behen* LINNÉ as new host plant of *Urophora merzi* from Iran. KARIMPOUR (2011), introduced 13 species of Aseraceae as new host plants of Tephritids in Urmia province. Also the larvae of the species *Aciura afghana* (HERING, 1961) were collected from various species of Lamiaceae plants in Iran (KHAGHANINIA & GHARAJEDAGHI 2013). Before this study, the fruit flies fauna of Saqez region has not been studied. Since most of the tephritid flies are important as biological agents of some of the Asteraceus plants, this research has been done to increasing the knowledge of the tephritid flies fauna and their host plants in the region.

Material and methods

Flower heads of Asteraceus thistles were collected from Saqez region during 2012-2013. The collected host plants were deposited to the greenhouse of Plant Protection Department of Tabriz University. Materials including capitula of asteraceus plants and tephritids larvae preserved in net containers (20-30 °C and 40% humidity). Adult specimens exited of flower heads of the weeds in the greenhouse after 1-3 weeks. In order to identification, the containers filled by CO₂ and the adults were dropping to the alcohol and some of them were pinned for more studies. Tephritidae and asteraceus species were identified by WHITE et al. 1999; WHITE 1988 and Flouer of GHAHRAMAN & ATTAR 2000.

Results

In this study, 13 species belonging to 9 genera of the family Tephritidae from 10 species belonging to 6 genera of the weed asteraceus host plants were identified in Saqez region which totally are listed in table 1-1 and the new record ones are explained as follow.

Family: T e p h r i t i d a e

Tribe *Tephritisini* NEWMAN, 1834

Acanthiophilus helianthi (Rossi, 1794) (Fig. 1a)

M a t e r i a l e x a m i n e d : (1 ♂): from flower heads of *Onopordum acanthium* (new host plant record) and (52 ♂♂, 43 ♀♀): from flower heads of *Cirsium heterophyllum*, 36°04.134' N, 46°15.221' E, 1639 m. (1 ♂): from flower heads of *Acropitilon repens* (new host plant record) and (1 ♀): from *Senecio vulgaris* and (2 ♀): from *Centaurea solstitialis*, 36°10.471' N, 46°20.291' E, 1603 m.

H o s t p l a n t s i n S a q e z : From flower heads of some species such as *Acropitilon repens* (LINNÉ, 1837), *Onopordum acanthium* (LINNÉ, 1753), *Cirsium heterophyllum*

(HILL, 1768), *Centaurea solstitialis* and *Senecio vulgaris* were collected. *Acroptilon repens* and *Onopordum acanthium* are first records of the host plants in Iran.

Distribution: Poland, North Africa, Turkey, Spain, Sweden and Afghanistan (RICHTER 1988; NORRBOM et al. 1999); Iran: Tehran, Tabriz and Ilam (MOHAMMADZADE NAMIN et al. 2010).

***Tephritis postica* (LOEW, 1844) (Fig. 1b)**

Material examined: (66 ♂♂, 94 ♀♀): from flower heads of *Onopordum acanthium* and (249 ♂♂, 236 ♀♀): from *Onopordum illyricum* (new host plant record), 36°044.134' N, 46°15.221' E, 1639 m.

Host plants in S a q e z : From flower heads of *Onopordum acanthium* and *Onopordum illyricum* were collected. *Onopordum illyricum* is a new host plant record from Iran.

Distribution: France, Okarina, Uzbekistan, South to North Africa and Cent. Europe (RICHTER, 1988; NORRBOM et al. 1999); Iran: Tehran and Tabriz (MOHAMMADZADE NAMIN et al. 2010; GHARAJEDAGHI et al. 2011).

***Tephritisomyia lauta* (LOEW, 1869) (Fig. 1c)**

Material examined: (22 ♂♂, 20 ♀♀): from *Echinops spinosissimus* (new host plant record) and (28 ♂♂, 15 ♀♀): from *Echinops viscosus*, 36°10.365' N, 46°04.061' E, 1570 m.

Host plants in S a q e z : from flower heads of *Echinops viscosus* and *Echinops spinosissimus* were collected. *Echinops spinosissimus* is first record of the host plant from Iran.

Distribution: Greece, Tunisia, and Egypt (NORRBOM et al. 1999); Iran: Tehran and Ilam (MOHAMMADZADE NAMIN et al. 2010; GHARAJEDAGHI et al. 2011).

Tribe *Terellini* HENDEL, 1927

***Terellia gynaecochroma* (HERING, 1937) (Figs.1j)**

Material examined: (46 ♂♂, 40 ♀♀): from *Onopordum acanthium* and (46 ♂♂, 64 ♀♀): from *Onopordum illyricum*, 36°10.471' N, 46°20.290' E, 1603 m.

Host plants in S a q e z : From flower heads of *Onopordum acanthium* and *Onopordum illyricum* were collected which *Onopordum illyricum* reported as a new host plant from Iran.

Distribution: North Europe, Turkey and Lebanon (NORRBOM et al., 1999; KNIO et al. 2002; KUTUK & VAROL 2006); Iran: Tehran and Tabriz (MOHAMMADZADE NAMIN et al. 2010).

Table 1-1: The list of the studied Tephritidae species and their host plants in Saqez regio

Tribe	Species	Host plant	
Myopitini	<i>Urophora stylata</i> (FABRICIUS 1775)	<i>Cirsium arvens</i> (Fig2.f)	(2♂, 4♀) (Fig1.d)
	<i>Urophora terebrans</i> (LOEW 1850)	<i>Onopordum acanthium</i> (Fig2.g)	(2♂) (Fig1.f)
	<i>Urophora xanthippe</i> (MUNRO 1934)	<i>Acroptilon repens</i> (Fig2.a)	(2♂, 4♀) (Fig1.e)
Xyphosiini	<i>Xyphosia miliaria</i> (SCHRANK 1781)	<i>Cirsium arvens</i>	(2♂, 2♀) (Fig1.g)
Tephritisini	<i>Sphenella marginata</i> (FALLEN 1814)	<i>Senecio vulgaris</i> (Fig2.c)	(7♂, 3♀) (Fig1.k)
Terellini	<i>Chaetorellia carthami</i> (STACKELBERG 1929)	<i>Centaurea solstitialis</i> (Fig2.e)	(16♂, 11♀) (Fig1.h)
	<i>Chaetostomella cylindrical</i> ROBINEAU-DESOVIDY 1830	<i>Onopordum</i>	(8♂, 3♀)
		<i>acanthium</i>	(17♂, 13♀)
		<i>O. illyricum</i>	(2♂, 1♀)
		<i>Cirsium arvense</i>	(1♀)
		<i>Cirsium palustre</i> (Fig2.g, b, f, j)	(Fig1. i)
	<i>Terellia quadratula</i> (LOEW 1869)	<i>Echinops viscosus</i> (Fig2.i)	(2♂, 1♀) (Fig1.l)
	<i>Terellia ruficauda</i> (FABRICIUS 1794)	<i>Cirsium arvens</i>	(6♂, 3♀) (Fig1.o)

Discussion

This study showed that the large number of these flies feeding from roughage species of Asteraceae which predicted to be as their biocontrol agents. WHITE (1988), THOMPSON (1998) & NORRBOM et al (1999) introduced roughage species of Asteraceae as host plants of the most tribes of Tephritidae. Also HARRIS (1980) showed that the two species of *Urophora affinis* and *Urophora quadrifasciata* can be used for the biological control of *Centaurea diffusa* and *Centaurea maculosa* in Canada. In this study most of the fruit flies were collected from roughage species such as *Cirsium* and *Onopordum*. However some asteraceus species such as the members of the genus *Acroptilon* are not roughage but because of their toxic secondary products, their control in pastures could be important too. Roughage plants make pastures useless for grazing and as they have high resistance to herbicides, their chemical control is so difficult (HARRIS 1980). Studies on application of these flies for biological control of weeds have been done already in the world but no study has been done in this aspect in Iran so far. Thus it is suggested that use of these flies as biological agents as integrated control programs increase the efficiency and utilization of pasture.

Acknowledgements

The authors would like to thank to Dr. Valery A. Korneyev (I.I. Schmalhausen Institute of Zoology, NAS of Ukraine) and Dr. S. Mohamadzade Namin (Islamic Azad university of Varamin) for confirm the identification of the species in this study.

References

- DLUSSKII G., GLASUNOVA K. & N. LAVROVA (2004): The flower and blossom morphology of Asteraceae correlates with composition pf their polnators. – Journal of General Biology **65** (6): 490-499.
- GHARAJEDAGHI Y., KHAGHANINIA S. & S. MOHAMMADZADE NAMIN (2011): Study of the genus *Urophora* ROBINEAU-DESOVIDY, 1830 (Diptera: Tephritidae) in Ecebsir region with two species as new records for Iran. – Munis Entomology and Zoology 6 (1): 358-362.
- GHAHRAMAN A. & F. ATTAR (2000): Biodiversity of plant species in Iran. – Tehran University Publication: 120p. Volume 3.
- HARRIS P. (1980): Establishment of *Urophora affinis* and *Urophora quadrifasciata* (Diptera: Tephritidae) in Canada for the biological control of diffuse and spotted knapweed. – Zeitschrift für angewandte Entomologie **89**: 504-514.
- KAPOOR V.C., HARDY D.E., AGARWAL M.L. & J.S. GREWAL (1980): Fruit fly (Diptera: Tephritidae) systematics of the Indian subcontinent. – Export India Publications, Jallundur, India: 111 pp.
- KARIMPOUR Y. (2011): Fruit fly (Diptera: Tephritidae) reared from capitula of Asteraceae in the Urmia region, Iran. – Journal of Entomological Society of Iran **30** (2): 53-66.
- KNIO K.M., KALASH S.H. & I.M. WHITE (2002): Flower head-infesting fruit flies (Diptera: Tephritidae) on thistles (Asteraceae) in Lebanon. – Journal of Natural History **36**: 617-629.
- KHAGHANINIA S. & Y. GHARAJEDAGHI (2013): Study of fruit flies (Diptera: Tephritidae) of weeds in East Azerbaijan province and introduction of a new species for Iran fauna. – Plant Pest Research No. 3, Volume 3.
- KUTUK M. & M.I. VAROL (2006): Records of two new species of *Terellia* ROBINEAU-DESOVIDY, 1830 (Diptera: Tephritidae) new to the Turkish fauna. – Turkish Journal of Zoology **30**: 99-102.
- MOHAMMADZADE NAMIN S., NOZARI J. & G.H. RASOLIAN (2010): The fruit flies (Diptera, Tephritidae) in Tehran province, with new records for Iranian fauna. – Vestnik Zoologii **44** (1): 20-31.
- MOHAMMADZADE NAMIN S. & J. NOZARI (2011): A new species of *Urophora* ROBINEAU-DESOVIDY, 1830 (Diptera: Tephritidae) from Iran. – Zookeys **152**: 63-70.
- NORRBOM A.L., CARROLL L.E., THOMPSON F.C., WHITE I.M. & A. FREIDBERG (1999): Systematic Database of Names. – In: Thompson, F. C., ed. Fruit Fly Expert Identification System and Systematic Information Database, Myia: 65-299.


- NORRBOM A.L. (2004): Updates to a biosystematic database of world Diptera for Tephritidae throughout 1999. – Diptera Data Dissemination Disk (CD-ROM) 2.
- RICHTER V.A. (1988): Family Tephritidae (Tryptidae); Keys to the insects of the European part of the USSR, In: Volume 5: Diptera and Siphonaptera. Part 2. Bei-Bienko, G. Y. (Ed.). – Institute of zoology, Academy of Sciences, USSR: pp. 212-276.
- THOMPSON F.C. (1998): Fruit fly expert identification system and systematic information database. – North American Dipteristso Society, Backhuys Publishers, Leiden, The Netherlands: 524 pp.
- WHITE I.M. (1988): Tephritisid flies. In: Handbooks for the identification of british insects. – Royal Entomological Society of London 10: 134 pp.
- WHITE I.M. & M.M. ELSON-HARRIS (1992): Fruit flies of economic significance: their identification and bionomics. – CAB International, London: 601 p.
- WHITE I.M., HEADRICH D.H., NORRBOM A.L. & L.E. CARROLL (1999): Glossary. – In: M. Aluja & A. L. Norrbom (Eds.), Fruit flies (Tephritidae): Phylogeny and evolution of behavior. – CRC Press. Boca Raton, USA: pp. 881-920.
- WOODS D.M., PITCARIN M.J., JOLEY D.B. & C.E. TURNER (2008): Seasonal phenology and impact of *Urophora sirunaseva* on yellow star thistle seed production in California. – Biological Control 47: 172-179.

Author's address:


Mozhgan ZAMANI
University of Tabriz
Department of Plant Protection
Faculty of Agriculture
51664, Tabriz, I.R. Iran

Correspongding author:

e-mail: skhaghaninia@gmail.com



Figs. 1: Lateral view of (a) *Acanthiophilus helianthi*; (b) *Tephritis postica*, (c) *Tephritomyia lauta*; (d) *Urophora stylsta*; (e) *Urophora xanthippe*; (f) *Urophora terebrans*; (g) *Xyphosia mialiaria*, (h) *Chaetorellia carthami*; (i) *Chaetostomella cylindrica*; (j) *Terellia gynacochroma*; (k) *Sphenella marginata*; (l) *Terellia quadratulla*; (o) *Terellia ruficauda* (original).



Figs 2: Flower head of (a) *Acroptilon repens*; (b) *Onopordum Illyricum*; (c) *Senecio vulgaris*; (d) *Cirsium heterophyllum*; (e) *Centaurea solstitialis*; (f) *Cirsium arvens*; (g) *Onopordum acanthium*; (h) *Echinops spinosissimus*; (i) *Echinops viscosus*; (j) *Cirsium palustre* (original).

Druck, Eigentümer, Herausgeber, Verleger und für den Inhalt verantwortlich:
Maximilian SCHWARZ, Konsulent f. Wissenschaft der Oberösterreichischen Landesregierung, Eibenweg 6,
A-4052 Ansfelden, Austria; maximilian.schwarz@liwest.at.

Redaktion: Fritz GUSENLEITNER, Biologiezentrum Linz, f.gusenleitner@landesmuseum.at;
Roland GERSTMAYER, Lehrstuhl f. Zoologie, TU München, gerstmei@wzw.tum.de;
Thomas WITT, Tengstraße 33, D-80796 München, thomas@witt-thomas.com;
Berthold CLEWING, Akademischer Verlag München, avm@druckmedien.de;
Harald SULAK, Museum Witt München, h.sulak@atelier-sulak.de;

Mitarbeiter: Karin TRAXLER, Biologiezentrum Linz, bio.redaktion@landesmuseum.at;
Heike REICHERT, Museum Witt München, heike_reichert66@web.de;
Erich DILLER, Zool. Staatssammlung München, Erich.Diller@zsm.mwn.de.

Adresse: Entomofauna, Redaktion und Schrifttausch Thomas WITT, c/o Museum Witt München,
Tengstr. 33, 80796 München, Deutschland, thomas@witt-thomas.com;
Entomofauna, Redaktion c/o Fritz GUSENLEITNER, Lungitzerstr. 51, 4222 St. Georgen/Gusen,
Austria, f.gusenleitner@landesmuseum.at.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Entomofauna](#)

Jahr/Year: 2016

Band/Volume: [0037](#)

Autor(en)/Author(s): Zamani Mozghan, Khaghaninia Samad

Artikel/Article: [Study of the fruit flies \(Diptera: Tephritidae\) as biocontrol agents of asteraceus plants in Saeqez region \(Kurdestan province\) 389-400](#)