

Entomofauna	40/2	Heft 17: 391-413	Ansfelden, 10. Okt. 2019
-------------	------	------------------	--------------------------

The Geometridae (Lepidoptera) fauna of Bartın province in Black Sea Region, with a new record *(Pachycnemia tibiaria* (RAMBUR, 1829)) for Turkey

Mustafa ÖZDEMİR

Abstract

In order to determine the Geometridae fauna of Bartın province where costal region of Black Sea, 1216 specimens collected between the years 2008 and 2012 by light trap in collection of Nazife Tuatay Plant Protection Museum and previous records from study area were evaluated. Totally 102 Geometridae species belong 4 subfamily were determined. There were 5 recorded species up to date but 4 recorded species could not be founded. 93 species were recorded for the first time from Bartın province. *Pachycnemia tibiaria* (RAMBUR, 1829) were recorded newly for Turkish Geometridae fauna also.

Key words: *Pachycnemia tibiaria*, new record, Geometridae, Black Sea, Bartın, Turkey

Zusammenfassung

Um die Geometridae-Fauna der Bartın-Provinz, in der Küstenregion des Schwarzen Meeres, zu bestimmen, wurden 1216 Exemplare, gesammelt durch Lichtfalle in den Jahren von 2008 bis 2012, aus der Sammlung des Nazife Tuatay Pflanzenschutzmuseums und frühere Aufzeichnungen aus dem Untersuchungsgebiet ausgewertet. Insgesamt 102 Geometridae-Arten wurden als zu 4 Unterfamilien gehörig bestimmt. Bisher wird von 5 Arten berichtet, 4 Arten aber konnten nicht nachgewiesen werden. 93 Arten wurden zum ersten Mal in der Provinz Bartın registriert. *Pachycnemia tibiaria* (RAMBUR, 1829) wurde in die türkische Geometridae-Fauna neu aufgenommen.

Introduction

The geometrids are the most species-rich family of Lepidoptera apart from Noctuidae and Pyralidae. There are over 23000 decrypted species known in worldwide (SCOBLE & HAUSMANN, 2007), 900 in Europe (HAUSMANN, 2001) and 608 in Turkey (KOÇAK & KEMAL, 2009). Geometrid species are ordinarily nocturnal, they often fly preferably at dusk, others are said to be active mainly after midnight but some species fly by day. Most geometrids

are sensitive to changing of habitat quality. Compared with most other macrolepidopterous families, species of Geometridae tend to be more specific to certain habitats and also highly suitable for use as an environmental indicator group. (SCOBLE, 1999; HAUSMANN, 2001).

Bartin is one of the provinces of Turkey where state on Southwestern coast of Black Sea. It is located in the Euxin province within the borders of the Euro-Siberian region as biogeographically. (DEMİRÖRS & KURT, 2005).

The study area is oceanic climate feature in dominantly and characterized humid forests, pseudo-maquis and primary structure degraded farmland as plant communities. The main tree species of humid forests on the more north-facing surface of the mountains are *Fagus orientalis*, *Quercus petraea*, *Castanea sativa* and in higher levels *Abies bornmulleriana*. Among these species are common as scattered *Carpinus betulus*, *Populus tremula*, *Alnus glutinosa*, *Acer platanoides*, ve *Tilia tomentosa*, *Rhododendron ponticum*, *Daphne pontica* and *Ilex aquifolium* as the undergrowth plant. Dry forests cover much narrower areas than humid forests. Ravaged forest areas at the costal regions are covered by pseudomaquis. Pseudomaquis starts from the costal line, it goes up to 350-400 m. It is also seen throughout the valleys of the sea exposure. However, in some regions primary vegetation is disturbed by factors such as agricultural activities and overgrazing (DÖNMEZ, 2014).

The studies on determine the geometrid fauna of the Black Sea region of Turkey were firstly started with MANN (1859, 1861) in Amasya and SCHWINGENSCHUSS (1939a, 1939b) in Bolu province. SEVEN (1991, 1995), KOÇAK & SEVEN (1991), ÖZDEMİR (2007), CAN (2008), OKYAR (2012) and ÖZDEMİR (2016, 2017) can be given as important studies on Geometridae fauna from different part of Black Sea. AKBULUT et al. (2003), OKYAR et. al. (2009) and AKKUZU et al. (2015) are studied on geometrids where close locations of Bartın. The knowledge on Geometridae fauna of Bartın province is quite limited. There were 9 species recorded from Bartın until now (TOPER KAYGIN et al., 2009; OKYAR, 2012).

The importance of faunistic records is great because the spread of species can be accurately assessed. In this article, the findings of the study carried out in the province of Bartın on the southwestern coastal line of the Black Sea and the previous records from this area were evaluated jointly. It is believed that the study will contribute to the evaluation of species diversity in the black sea coastal line and neighboring areas.

Material and Methods

In this study, the specimens collected by light trap (8W black light fluorescent lamp) from Bartın Province during the years 2008-2012 and previous records were investigated. The specimens were determined according to external characters and male or female genitalia morphology. For genitalia preparations, the abdomens of materials were boiled in 10% KOH solution, dissected under a stereo microscope, and slides were mounted with Entellan. VARDIKJAN (1985), HAUSMANN (2001, 2004) MIRONOV (2003) HAUSMANN & VIIDALEPP (2012) and SKOU & SIHVONEN (2015) were used for the identification.




Fig. 1: The map to the localities sampled and previous records in the study area.

Results

Subfamily: Geometrinae STEPHENS, 1829

Chlorissa cloraria (HÜBNER, 1813)

E x e m i n e d m a t e r i a l : Amasra, Aliobasi 150m 22.V.2012 1♂, 41°46'39"N 32°30'59"E; Çakrazova 65m 21.V.2012 1♂, 41°46'38"N 32°30'17"E; Center, Bahçecik 244m 19.VII.2012 1♀ 2♂♂, 41°28'49"N 32°36'41"E.

D i s t r i b u t i o n i n T u r k e y : Balikesir Bolu, Düzce Edirne Zonguldak (KOÇAK & KEMAL 2009, ALTIPARMAK 2010; OKYAR 2012)

Eucrostes indigenata (DE VILLERS, 1789)

E x e m i n e d m a t e r i a l : Kuruçâile, Uğurlu 210m 2.VIII.2011 1♀ 1♂, 41°50'8"N 32°41'40"E.

D i s t r i b u t i o n i n T u r k e y : Antalya, Bursa, Hatay (KOÇAK & KEMAL 2009; CAN 2010).

Hemithea aestivaria (HÜBNER, 1789)

E x e m i n e d m a t e r i a l : Amasra, Akkonak 50m 19.VII.2012 1♂, 41°47'25"N 32°30'12"E; Hatipler 100m 2.VIII.2011 1♀, 41°46'39"N 32°30'18"E.

D i s t r i b u t i o n i n T u r k e y : Amasya, Bolu, Bursa, Düzce, Edirne, İstanbul, Ordu, Trabzon, (OKYAR & MIRONOV 2008; KOÇAK & KEMAL 2009; ÖZDEMİR 2016).

***Jodis lactearia* (LINNAEUS, 1758)**

Ex em i n e d m a t e r i a l : Amasra, Akkonak 50m 19.VII.2012 1♂, 41°47'25"N 32°30'12"E.
D i s t r i b u t i o n i n T u r k e y : Bolu, Bursa, Düzce, Kastamonu, Kayseri, Ordu, Rize,
Tekirdağ, Trabzon (KOÇAK & KEMAL 2009; OKYAR 2010b; ÖZDEMİR 2016).

***Phaiogramma etruscaria* (ZELLER, 1849)**

Ex em i n e d m a t e r i a l : Amasra, Akkonak 200m 2.VIII.2011 1♂, 41°47'19"N 32°30'51"E;
Çakrazova 65m 21.V.2012 1♂, 41°46'38"N 32°30'17"E; Hatipler 100m 2.VIII.2011 2♂♂,
41°46'39"N 32°30'18"E; Kuruçâşile, Uğurlu 210m 2.VIII.2011 1♂, 41°50'08"N 32°41'40"E.
D i s t r i b u t i o n i n T u r k e y : Artvin, Adana, Amasya, Bolu, Bursa, Çanakkale, Edirne,
Gaziantep, Giresun, Hatay, İsparta, İçel, Kırklareli, Konya, Kahramanmaraş, Ordu, Osmaniye,
Siirt, Sivas, Trabzon, Van (DOĞANLAR 2003; OKYAR & MIRONOV 2008; KOÇAK & KEMAL 2009;
CAN 2010; GÖKTÜRK 2011; KOYUNCU 2011; KEMAL & SEVEN 2013; ÖZDEMİR 2016; SEVEN &
AKIN 2016).

***Proteuchloris neraria* (HERRICH-SCHÄFFER, 1852)**

P r e v i o u s r e c o r d f r o m B a r t i n : Bartın Credit and Housing Agency Park
01.VI.2007 (TOPER KAYGIN et al., 2009)

D i s t r i b u t i o n i n T u r k e y : Adana, Amasya, Bolu, Bursa, Bartın, Çanakkale, Edirne,
Gaziantep, Hatay, İçel, Kahramanmaraş, Osmaniye, Siirt (KOÇAK & KEMAL 2009; CAN 2010;
OKYAR 2010; KOYUNCU 2011; SEVEN 2014).

***Pseudoterpn a pruinata* (HUFNAGEL, 1767)**

E x em i n e d m a t e r i a l : Amasra, Hatipler 100m 2.VIII.2011 1♀ 1♂, 41°46'39"N
32°30'18"E; Arit, Kocaköy 180m 6.IX.2010 3♂♂, 41°41'50"N 32°29'34"E.

D i s t r i b u t i o n i n T u r k e y : Ankara, Bolu, Bursa, Düzce, Karabük, Muğla, Sinop
(KOÇAK & KEMAL 2009; OKYAR 2012).

Subfamily: S t e r r h i n a e MEYRICK, 1892

***Cyclophora albiocellaria* (HÜBNER, 1789)**

E x em i n e d m a t e r i a l : Amasra, Çakrazova 65m 21.V.2012 1♂, 41°46'38"N 32°30'17"E.
D i s t r i b u t i o n i n T u r k e y : Amasya, Bursa, Düzce (AKBULUT et al. 2003; KOÇAK & KE-
MAL 2009).

***Cyclophora annularia* (FABRICIUS, 1775)**

E x em i n e d m a t e r i a l : Amasra, Akkonak 200m 2.VIII.2011 1♂, 41°47'19"N 32°30'51"E;
50m 19.VII.2012 1♀, 41°47'25"N 32°30'12"E; Çakrazova 65m 21.V.2012 1♀, 41°46'38"N
32°30'17"E; Kuruçâşile, Paşalar 400m 22.V.2012 2♀♀, 41°46'47"N 32°36'28"E.

D i s t r i b u t i o n i n T u r k e y : Bolu, Bursa, Düzce, İstanbul, Rize, Kahramanmaraş,
Trabzon, (KOÇAK & KEMAL 2009).

***Cyclophora linearia* (HÜBNER, 1799)**

E x e m i n e d m a t e r i a l : Amasra, Akkonak 200m 2.VIII.2011 1♂, 41°47'19"N 32°30'51"E; Çakrazova 65m 21.V.2012 1♀, 41°46'38"N 32°30'17"E; Kurucasile, Paşalar 400m 22.V.2012 21♀♀ 17♂♂, 41°46'47"N 32°36'28"E; Ulus, Aşağıemirce 240m 29.IX.2011 2♀♀ 6♂♂, 41°32'01"N 32°35'52"E.

D i s t r i b u t i o n i n T u r k e y : Adana, Bolu, Düzce, Edirne, Hatay, İstanbul, Kastamonu, Kırklareli Ordu, Rize, Trabzon, (KOÇAK & KEMAL 2009; OKYAR 2010b; ÖZDEMİR 2016)

***Cyclophora pupillaria* (HÜBNER, 1799)**

E x e m i n e d m a t e r i a l : Amasra, Akkonak 200m 2.VIII.2011 27♀♀ 24♂♂, 41°47'19"N 32°30'51"E; Çakrazova 65m 21.V.2012 1♂, 41°46'38"N 32°30'17"E; Kozpinari 327m 29.IX.2011 1♂, 41°42'41"N 32°21'50"E; Kurucasile, Curunlu 220m 2.VIII.2011 11♀♀ 19♂♂, 41°49'44"N 32°38'34"E; Dizlermezecili 220m 22.V.2012 1♀ 1♂, 41°47'52"N 32°35'01"E; Karaman Village 130m 19.IX.2012 1♀, 41°49'43"N 32°38'22"E; Bartın center, Arıönü 300m 28.IX.2011 1♂, 41°7'52"N 32°12'04"E; Bahçecik 244m 19.VII.2012 1♂, 41°28'49"N 32°36'41"E; Port area 130m 28.IX.2011 1♀ 8♂♂, 41°40'50"N 32°13'51"E; Ulus, 300m 29.IX.2011 1♂, 41°34'41"N 32°38'12"E.

D i s t r i b u t i o n i n T u r k e y : Adana, Ankara, Antalya, Bolu, Bursa, Çanakkale, Düzce, Hatay, İçel, İstanbul, Kahramanmaraş, Konya, Manisa, Muğla, Ordu, Osmaniye, Sivas (DOĞANLAR 2003; KOÇAK & KEMAL 2009; ÖZDEMİR 2016).

***Cyclophora quercimontaria* (BASTELBERGER, 1897)**

E x e m i n e d m a t e r i a l : Amasra, Çakrazova 65m 21.V.2012 3♀♀ 2♂♂, 41°46'38"N 32°30'17"E; Top Deresi 234m 21.V.2012 1♂, 41°42'39"N 32°23'49"E; Ulus, Aşağıemirce 240m 29.IX.2011 1♂, 41°32'01"N 32°35'52"E.

D i s t r i b u t i o n i n T u r k e y : Artvin, Bolu, Düzce, Ordu, Van (KOÇAK & KEMAL 2009; ÖZDEMİR 2016).

***Idaea aversata* (LINNAEUS, 1758)**

E x e m i n e d m a t e r i a l : Amasra, Hatipler 100m 2.VIII.2011 1♀, 41°46'39"N 32°30'18"E; Bartın center, Bahçecik 244m 19.VII.2012 6♀♀ 2♂♂, 41°28'49"N 32°36'41"E.

D i s t r i b u t i o n i n T u r k e y : Amasya, Ardahan, Bolu, Bursa, Çanakkale, Düzce, Edirne, Hatay, İstanbul, Kırklareli, Konya, Ordu, Sakarya, Siirt, Tekirdağ, Trabzon, Van (DOĞANLAR 2003; OKYAR & MIRONOV 2008; KOÇAK & KEMAL 2009; OKYAR 2012; SEVEN 2014; ÖZDEMİR 2016).

***Idaea consolidata* (LEDERER, 1853)**

E x e m i n e d m a t e r i a l : Amasra, Akkonak 200m 2.VIII.2011 1♀, 41°47'19"N 32°30'51"E; Bartın center, Bahçecik 244m 19.VII.2012 1♀ 4♂♂, 41°28'49"N 32°36'41"E.

D i s t r i b u t i o n i n T u r k e y : Amasya, Ankara, Antalya, Bursa, İstanbul, Kahramanmaraş (KOÇAK & KEMAL 2009).

***Idaea degeneraria* (HÜBNER, 1799)**

Previous record : Bartın Credit And Housing Agency Park 10.VI.2007 (TOPER KAYGIN et al. 2009).

E x e m i n e d m a t e r i a l : Amasra, Akkonak 200m 2.VIII.2011 1♀, 41°47'19"N 32°30'51"E; Aliobası 150m 22.V.2012 1♂, 41°46'39"N 32°30'59"E; Çakrazova 65m 21.V.2012 12♀♂ 5♂♂, 41°46'38"N 32°30'17"E; Kuruçâşile, Curunlu 220m 2.VIII.2011 1♀, 41°49'44"N 32°38'34"E; Paşalar 400m 22.V.2012 5♀♀ 3♂♂, 41°46'47"N 32°36'28"E; Port area 130m 28.IX.2011 1♂, 41°40'50"N 32°13'51"E; Ulus, Aşağıemirce 240m 29.IX.2011 1♀, 41°32'01"N 32°35'52"E; Hasanören 640m 29.IX.2011 1♀, 41°37'34"N 32°45'19"E.

D i s t r i b u t i o n i n T u r k e y : Adana, Amasya, Ankara, Bartın, Bolu, Bursa, Çanakkale, Edirne, Gaziantep, Hatay, Hakkari, İçel, İstanbul, Kahramanmaraş, Kırıkkale, Kırklareli, Konya, Ordu, Osmaniye, Siirt, Sivas, Tekirdağ, Tokat, Trabzon, Van (DOĞANLAR 2003; OKYAR & MIRONOV 2008; KOÇAK & KEMAL 2009; TOPER KAYGIN et al. 2009; ALTIPARMAK 2010; KOYUNCU & KÜTÜK 2011; KEMAL & SEVEN 2013; ÖZDEMİR 2016; KEMAL et al. 2017).

***Idaea dimidiata* (HUFNAGEL, 1767)**

E x e m i n e d m a t e r i a l : Amasra, Akkonak 200m 2.VIII.2011 3♂♂, 41°47'19"N 32°30'51"E; Çakrazova 65m 21.V.2012 1♂, 41°46'38"N 32°30'17"E; Arit, Kocaköy 180m 6.IX.2010 1♂, 41°41'50"N 32°29'34"E.

D i s t r i b u t i o n i n T u r k e y : Adana, Amasya, Ardahan, Bolu, Bursa, Düzce, Gaziantep, Hatay, İstanbul, Kırklareli, Manisa, Muğla, Ordu, Kahramanmaraş, Zonguldak (KOÇAK & KEMAL 2009; KOYUNCU & KÜTÜK 2011; OKYAR 2012; ÖZDEMİR 2016).

***Idaea filicata* (HÜBNER, 1799)**

E x e m i n e d m a t e r i a l : Amasra, Çakrazova 65m 21.V.2012 1♀ 1♂, 41°46'38"N 32°30'17"E; Top Deresi 234m 21.V.2012 1♂, 41°42'39"N 32°23'49"E; Kuruçâşile, Paşalar 400m 22.V.2012 4♀♀, 41°46'47"N 32°36'28"E.

D i s t r i b u t i o n i n T u r k e y : Adana, Amasya, Bolu, Bursa, Çanakkale, Edirne, Hakkari, İçel, İstanbul, Kırıkkale, Konya, Manisa, Kahramanmaraş, Muğla, Ordu, Osmaniye, Siirt, Sivas, Tekirdağ, Tokat, Trabzon, Van (DOĞANLAR 2003; KOÇAK & KEMAL 2009; SEVEN 2014; ÖZDEMİR 2016; KEMAL et al. 2017).

***Idaea infirmaria* (RAMBUR, 1833)**

E x e m i n e d m a t e r i a l : Kuruçâşile, Curunlu 220m 2.VIII.2011 1♀, 41°49'44"N 32°38'34"E.

D i s t r i b u t i o n i n T u r k e y : Çanakkale, Muğla (KOÇAK & KEMAL 2009).

***Idaea deversaria* (HERRICH & SCHAFER, 1847)**

E x e m i n e d m a t e r i a l : Amasra, Aliobası 150m 22.V.2012 5♀♀ 3♂♂, 41°46'39"N 32°30'59"E; Kazpinarı 60m 15.VI.2010 3♀♀, 41°42'05"N 32°21'54"E; Kuruçâşile, Karaman Village 130m 19.IX.2012 3♀♀ 8♂♂, 41°49'43"N 32°38'22"E; Paşalar 400m 22.V.2012 6♀♀ 5♂♂, 41°46'47"N 32°36'28"E.

D i s t r i b u t i o n i n T u r k e y : Amasya, Ankara, Antalya, Bolu, Bursa, Düzce, Gümüşhane, İçel, İstanbul, Kırklareli, Konya, Rize, Siirt, Trabzon (KOÇAK & KEMAL 2009; OKYAR 2010b; SEVEN 2014).

***Idaea obsoletaria* (RAMBUR, 1833)**

E x e m i n e d m a t e r i a l : Kuruçâşile, Uğurlu 210m 2.VIII.2011 2♂♂, 41°50'08"N 32°41'40"E.

D i s t r i b u t i o n i n T u r k e y : Amasya, Antalya, Bolu, Bursa, İzmir, Kırklareli Siirt Sivas (KOÇAK & KEMAL 2009; SEVEN 2014; KEMAL & KOÇAK 2015b).

***Idaea ochrata* (SCOPOLI, 1763)**

E x e m i n e d m a t e r i a l : Bartın center, Bahçecik 244m 19.VII.2012 1♀ 1♂, 41°28'49"N 32°36'41"E.

D i s t r i b u t i o n i n T u r k e y : Amasya, Bolu, Bursa, Çanakkale, Düzce, Edirne, Gaziantep, İğdır, İstanbul, Kırklareli, Kahramanmaraş, Karabük, Kastamonu, Ordu, Siirt, Tekirdağ, Van, Zonguldak (OKYAR & MIRONOV 2008; KOÇAK & KEMAL 2009; ALТИПАРМАК 2010; KOYUNCU & KÜTÜK 2011; OKYAR 2012; SEVEN 2014; ÖZDEMİR 2016)

***Idaea politaria* (HÜBNER, 1799)**

E x e m i n e d m a t e r i a l : Amasra, Akkonak 50m 19.VII.2012 2♂♂, 41°47'25"N 32°30'12"E; Bartın center, Şabankadı 250m 23.VI.2009 1♂, 41°29'38"N 32°17'42"E.

D i s t r i b u t i o n i n T u r k e y : Adana, Amasya, Antalya, Bolu, Bursa, Düzce, Edirne, Hatay, İçel, İstanbul, Kastamonu, Kırklareli, Manisa Ordu Siirt (KOÇAK & KEMAL 2009; OKYAR 2012; SEVEN 2014; ÖZDEMİR 2016).

***Idaea rubraria* (STAUDINGER, 1901)**

E x e m i n e d m a t e r i a l : Amasra, Hatipler 100m 2.VIII.2011 1♂, 41°46'39"N 32°30'18"E; Kuruçâşile, Curunlu 220m 2.VIII.2011 1♀, 41°49'44"N 32°38'34"E; Uğurlu 210m 2.VIII.2011 2♂♂, 41°50'08"N 32°41'40"E.

D i s t r i b u t i o n i n T u r k e y : Scattered populations in Turkey (HAUSMANN 2004).

***Idaea rufaria* (HÜBNER, 1799)**

E x e m i n e d m a t e r i a l : Amasra, Kalayci 142m 4.VI.2008 1♂, 41°47'30"N 32°33'34"E.

D i s t r i b u t i o n i n T u r k e y : Amasya, Bolu, Bursa, Çanakkale, Çorum, Edirne, İğdır, İçel, İstanbul, Karabük, Kastamonu, Kırklareli, Konya, Osmaniye, Sivas, Van (KOÇAK & KEMAL 2009; OKYAR 2012)

***Idaea straminata* (BORKHAUSEN, 1794)**

E x e m i n e d m a t e r i a l : Arit, Kocaköy 180m 6.IX.2010 1♀ 1♂, 41°41'50"N 32°29'34"E.

D i s t r i b u t i o n i n T u r k e y : Amasya, Bursa, Düzce, Kars, Konya, Ordu, Rize (KOÇAK & KEMAL 2009; OKYAR 2010b; ÖZDEMİR 2016)

***Idaea subsericeata* (HAWORTH, 1809)**

E x e m i n e d m a t e r i a l : Amasra, Akkonak 200m 2.VIII.2011 4♀♀ 7♂♂, 41°47'19"N 32°30'51"E; Çakrazova 65m 21.V.2012 2♀♀ 1♀, 41°46'38"N 32°30'17"E; Hatipler 100m 2.VIII.2011 5♂♂, 41°46'39"N 32°30'18"E; Arit, Kocaköy 180m 6.IX.2010 3♂♂, 41°41'50"N 32°29'34"E; Kuruçâşile, Curunlu 220m 2.VIII.2011 2♂♂, 41°49'44"N 32°38'34"E; Paşalar 400m 22.V.2012 1♀, 41°46'47"N 32°36'28"E; Uğurlu 210m 2.VIII.2011 2♂♂, 41°50'8"N 32°41'40"E.

D i s t r i b u t i o n i n T u r k e y : Adana, Amasya, Antalya, Bolu, Bursa, Düzce, Çanakkale, Edirne, Hatay, İçel, İstanbul, Kahramanmaraş, Ordu, Siirt, Tekirdağ (KOÇAK & KEMAL 2009; OKYAR 2012; SEVEN 2014; ÖZDEMİR 2016).

***Rhodostrophia discopunctata* AMSEL, 1935**

E x e m i n e d m a t e r i a l : Amasra, Çakrazova 65m 21.V.2012 3♂♂, 41°46'38"N 32°30'17"E; Kazpinarı 60m 15.VI.2010 1♂, 41°42'05"N 32°21'54"E; Bartın center, Şabankadı 250m 23.VI.2009 4♀♀ 3♂♂, 41°29'38"N 32°17'42"E; Şarköy 200m 23.VI.2009 1♀, 41°29'14"N 32°21'37"E.

D i s t r i b u t i o n i n T u r k e y : Amasya, Ankara, Batman, Bitlis, Bolu, Bursa, Çanakkale, Çankırı, Düzce, Gaziantep, Girasun, Gümüşhane, Hakkari, İçel, Karabük, Kastamonu, Kayseri, Konya, Kahramanmaraş, Nevşehir, Osmaniye, Rize, Siirt, Sivas, Tokat, Van (KOÇAK & KEMAL 2009; OKYAR 2010b; KOYUNCU & KÜTÜK 2011; OKYAR 2012; KEMAL & SEVEN 2013; KEMAL et al. 2017).

***Rhodostrophia vibicaria* (CLERCK, 1759)**

E x e m i n e d m a t e r i a l : Amasra, Çakrazova 65m 21.V.2012 1♀ 1♂, 41°46'38"N 32°30'17"E.

D i s t r i b u t i o n i n T u r k e y : Amasya, Ardahan, Bayburt, Bitlis, Bolu, Çanakkale, Düzce, Edirne, Gümüşhane, İğdır, İstanbul, Kastamonu, Kayseri, Kırklareli, Konya, Ordu, Siirt, Sinop, Sivas, Tekirdağ, Tokat, Trabzon, Van, Bayburt, İğdir (OKYAR & AKTAÇ 1999; AKBULUT et al. 2003; KOÇAK & KEMAL 2009; KEMAL & SEVEN 2013; OKYAR 2012; KEMAL & KOÇAK 2015b; ÖZDEMİR 2016).

***Scopula flaccidaria* (ZELLER, 1852)**

E x e m i n e d m a t e r i a l : Amasra, Akkonak 200m 2.VIII.2011 1♀ 1♂, 41°47'19"N 32°30'51"E; 32°30'19"E; 19.VII.2012 1♂, 41°47'25"N 32°30'12"E; Kalayçı 142m 4.VI.2008 1♂, 41°47'30"N 32°33'34"E; Arıt, Kocaköy 180m 6.IX.2010 1♂, 41°41'50"N 32°29'34"E; Kuruçışile, Uğurlu 210m 2.VIII.2011 1♂, 41°50'08"N 32°41'40"E.

D i s t r i b u t i o n i n T u r k e y : Adana, Amasya, Bursa, Edirne, Hatay, İçel, Kırklareli, Kahramanmaraş, Siirt, Tekirdağ (KOÇAK & KEMAL 2009; SEVEN 2014).

***Scopula imitaria* (Hübner, 1799)**

P r e v i o u s r e c o r d : 12.VI.2007, Catmaca (adult) (TOPER KAYGIN et al. 2009).

E x e m i n e d m a t e r i a l : Amasra, Akkonak 200m 2.VIII.2011 1♀, 41°47'19"N 32°30'51"E; Çakrazova 65m 21.V.2012 1♂, 41°46'38"N 32°30'17"E; Kazpinarı 60m 15.VI.2010 2♀♀ 4♂♂, 41°42'05"N 32°21'54"E; Kuruçışile, Karaman Village 130m 19.IX.2012 1♂, 41°49'43"N 32°38'22"E; Bartın center, Port area 130m 28.IX.2011 1♀, 41°40'50"N 32°13'51"E.

D i s t r i b u t i o n i n T u r k e y : Amasya, Antalya, Aydın, Balıkesir, Bartın, Bolu, Bursa, Düzce, Edirne, Hatay, İğdır, İstanbul, İzmir, Manisa, Muğla, Sinop, Tekirdağ (AKBULUT et al. 2003; KOÇAK & KEMAL 2009; ALTİPARMAK 2010).

***Scopula marginepunctata* (GOEZE, 1781)**

E x e m i n e d m a t e r i a l : Kuruçışile, Curunlu 220m 2.VIII.2011 1♂, 41°49'44"N 32°38'34"E; Karaman village 130m 19.IX.2012 2♀♀, 41°49'43"N 32°38'22"E; Uğurlu 210m 2.VIII.2011 1♀ 2♂♂, 41°50'08"N 32°41'40"E; Bartın center, Arıönü 300m 28.IX.2011 2♂♂, 41°37'52"N 32°12'04"E; Güzelcehisar 220m 28.IX.2011 1♀ 1♂, 41°39'13"N 32°13'21"E; Port area 130m 28.IX.2011 1♀, 41°40'50"N 32°13'51"E; Ulus, Gökpınar 730m 29.IX.2011 1♂ 41°38'35"N 32°50'34"E; Hasanören 640m 29.IX.2011 2♂♂, 41°37'34"N 32°45'19"E.

D i s t r i b u t i o n i n T u r k e y : Adana, Amasya, Bolu, Bursa, Çanakkale, Diyarbakır, Düzce, Edirne, Erzincan, Hakkari, İğdır, İstanbul, İzmir, Kayseri, Kırıkkale, Kırklareli, Konya, Kahramanmaraş, Ordu, Osmaniye, Siirt, Sivas, Tekirdağ, Tokat, Trabzon, Van (KOÇAK & KEMAL 2009; KEMAL et al. 2011; KEMAL & SEVEN 2013; ÖZDEMİR 2016; KEMAL et al. 2017).

***Scopula nigropunctata* (HUFNAGEL, 1767)**

E x e m i n e d m a t e r i a l : Amasra, Akkonak 200m 2.VIII.2011 1♂, 41°47'19"N 32°30'51"E; Bartın center, Port area 130m 28.IX.2011 1♀, 41°40'50"N 32°13'51"E.

D i s t r i b u t i o n i n T u r k e y : Ankara, Bolu, Düzce, Edirne, İstanbul, Kastamonu, Kırklareli, Ordu, Sakarya (KOÇAK & KEMAL 2009; OKYAR 2012; ÖZDEMİR 2016).

***Scopula ornata* (SCOPOLI, 1763)**

E x e m i n e d m a t e r i a l : Amasra, Akkonak 200m 2.VIII.2011 1♂, 41°47'19"N 32°30'51"E; Bartın center, Ariönü 300m 28.IX.2011 1♂, 41°37'52"N 32°12'04"E.

D i s t r i b u t i o n i n T u r k e y : Adana, Amasya, Ankara, Antalya, Aydın, Bilecik, Bitlis, Bolu, Bursa, Çanakkale, Çankırı, Düzce, Edirne, Gümüşhane, Hakkari, İğdır İstanbul, Kırklareli, Ordu, Tekirdağ, Trabzon, Van, Yalova (DOĞANLAR 2003; KOÇAK & KEMAL 2009; ÖZDEMİR 2016; KEMAL et al. 2017; KEMAL & KOÇAK 2017).

***Scopula rubiginata* (HUFNAGEL, 1767)**

P r e v i o u s r e c o r d : Ağdacı village 17.VI.2007, (TOPER KAYGIN et al. 2009).

D i s t r i b u t i o n i n T u r k e y : Amasya, Ankara, Antalya, Ardahan, Artvin, Bartın, Bitlis, Bolu, Bursa, Düzce, Erzincan, İstanbul, Kastamonu, Kırklareli, Konya, Malatya, Ordu, Sivas, Tekirdağ (KOÇAK & KEMAL 2009; GÖKTÜRK 2011; OKYAR 2012; ÖZDEMİR 2016).

***Scopula submutata* (TREITSCHKE, 1828)**

E x e m i n e d m a t e r i a l : Ulus, Gökpınar 730m 29.IX.2011 1♂, 41°38'35"N 32°50'34"E.

D i s t r i b u t i o n i n T u r k e y : Adana, Amasya, Ankara, Antalya, Bitlis, Bursa, Çanakkale, Erzincan, Hatay, İçel, Konya, Manisa, Kahramanmaraş, Kırıkkale, Mardin, Siirt, Sivas (KOÇAK & KEMAL 2009; KEMAL & SEVEN 2013).

***Timandra comae* SCHMIDT, 1931**

P r e v i o u s r e c o r d : Çamköy, 1♀, 1♂ (OKYAR 2012).

D i s t r i b u t i o n i n T u r k e y : Adana, Amasya, Artvin, Bartın, Bursa, Edirne, İçel, İstanbul, Kırklareli, Konya, Ordu, Rize, Samsun, Sivas, Tekirdağ (KOÇAK & KEMAL 2009; OKYAR 2012; ÖZDEMİR 2016).

Subfamily: L a r e n t i i n a e MEYRICK, 1892

***Aplocera efformata* (GUENÉE, 1858)**

E x e m i n e d m a t e r i a l : Kuruçışile, Curunlu 220m 2.VIII.2011 3♂♂, 41°49'44"N 32°38'34"E; Karaman village 130m 19.IX.2012 2♀♀ 3♂♂, 41°49'43"N 32°38'22"E; Bartın center, Güzelcehisar 220m 28.IX.2011 1♂, 41°39'13"N 32°13'21"E.

D i s t r i b u t i o n i n T u r k e y : Bolu, Çanakkale, Düzce, Edirne, İstanbul, Kırklareli, Manisa (KOÇAK & KEMAL 2009).

***Asthena albula* (HUFNAGEL, 1767)**

E x e m i n e d m a t e r i a l : Amasra, Akkonak 50m 19.VII.2012 1♂, 41°47'25"N 32°30'12"E; Çakrazova 65m 21.V.2012 2♀, 1♂, 41°46'38"N 32°30'17"E; Kurucasile, Paşalar 400m 22.V.2012 3♀, 7♂, 41°46'47"N 32°36'28"E; Bartın center, Bahçecik 244m 19.VII.2012 4♂, 41°28'49"N 32°36'41"E; Kaman 50m 21.V.2012 1♂, 41°40'25"N 32°21'56"E; Ulus, Aşağıemirce 240m 29.IX.2011 1♀, 41°32'01"N 32°35'52"E.

D i s t r i b u t i o n i n T u r k e y : Bolu, Düzce, Edirne, İstanbul, Kastamonu, Kırklareli, Trabzon Zonguldak (KOÇAK & KEMAL 2009; OKYAR 2012).

***Camptogramma bilineata* (LINNAEUS, 1758)**

E x e m i n e d m a t e r i a l : Amasra, Akkonak 50m 19.VII.2012 1♀, 41°47'25"N 32°30'12"E; Amasra, Çakrazova 65m 21.V.2012 1♀, 41°46'38"N 32°30'17"E; Kurucasile, Karaman village 130m 19.IX.2012 1♂, 41°49'43"N 32°38'22"E; Ulus, Hasanören 640m 29.IX.2011 1♀, 41°37'34"N 32°45'19"E.

D i s t r i b u t i o n i n T u r k e y : Adana, Amasya, Ankara, Balıkesir, Bitlis, Bolu, Bursa, Çanakkale, Çankırı, Edirne, Elazığ, Gaziantep, Giresun, Hakkari, Hatay, İçel, İstanbul, İzmir, Kahramanmaraş, Kars, Kayseri, Kırklareli, Konya, Malatya, Manisa, Muğla, Ordu, Osmaniye, Rize, Siirt, Sinop, Şırnak, Tekirdağ, Tokat, Van (DOĞANLAR 2003; KOÇAK & KEMAL 2009; OKYAR 2010b; KOYUNCU 2011; KOÇAK & KEMAL 2015; ÖZDEMİR 2016).

***Catarhoe rubidata* (DENIS & SCHIFFERMÜLLER, 1775)**

E x e m i n e d m a t e r i a l : Amasra, Çakrazova 65m 21.V.2012 1♀, 41°46'38"N 32°30'17"E; Kazpinarı 60m 15.VI.2010 1♀, 41°42'05"N 32°21'54"E; Kurucasile, Paşalar 400m 22.V.2012 1♀, 41°46'47"N 32°36'28"E.

D i s t r i b u t i o n i n T u r k e y : Amasya, Bingöl, Bitlis, Bursa, Edirne, İstanbul, Kırklareli, Van (KOÇAK & KEMAL 2009).

***Chloroclystis v-ata* (HAWORTH, 1809)**

E x e m i n e d m a t e r i a l : Amasra, Aliobası 150m 22.V.2012 3♀, 2♂, 41°46'39"N 32°30'59"E; Çakrazova 65m 21.V.2012 2♀, 2♂, 41°46'38"N 32°30'17"E; Kurucasile, Paşalar 400m 22.V.2012 11♀, 5♂, 41°46'47"N 32°36'28"E.

D i s t r i b u t i o n i n T u r k e y : Ordu, Rize (OKYAR 2010b; ÖZDEMİR 2016).

***Cidaria fulvata* (FORSTER, 1771)**

P r e v i o u s r e c o r d s : Sipahiler-Kayadibi 12.VII.2004 1250m, 1♂ (OKYAR 2012)

D i s t r i b u t i o n i n T u r k e y : Ankara, Ardahan, Bartın, Bayburt, Bitlis, Bolu, Erzincan, Gümüşhane, İğdir, Kastamonu, Konya, Sivas, Trabzon (KOÇAK & KEMAL 2009; OKYAR 2012).

***Coenotephria ablutaria hangayi* (VOJNITS, 1986)**

E x e m i n e d m a t e r i a l : Amasra, Çakrazova 65m 21.V.2012 1♀, 41°46'38"N 32°30'17"E; Kozpinarı 327m 29.IX.2011 7♂, 41°42'05"N 32°21'54"E; Bartın center, Ariönü 300m 28.IX.2011 1♂, 41°37'52"N 32°12'04"E.

D i s t r i b u t i o n i n T u r k e y : Hatay, İçel, Siirt, Van, Central Anatolia (HAUSMANN & VIIDALEPP 2012; SEVEN 2014; KEMAL & KOÇAK 2015a 2016c, 2017).

***Cosmorrhoe ocellata* (LINNAEUS, 1758)**

E x e m i n e d m a t e r i a l : Amasra, Çakrazova 65m 21.V.2012 2♂♂, 41°46'38"N 32°30'17"E; Hatıpler 100m 2.VIII.2011 1♂, 41°46'39"N 32°30'18"E; Kazpinarı 60m 15.VI.2010 1♂, 41°42'05"N 32°21'54"E; Arit, Kocaköy 180m 6.IX.2010 1♂, 41°41'50"N 32°29'34"E; Kurucasıle, Çurunlu 220m 2.VIII.2011 1♂, 41°49'44"N 32°38'34"E; Bartın center, Port area 130m 28.IX.2011 1♂, 41°40'50"N 32°13'51"E; Ulus, Aşağıemirce 240m 29.IX.2011 2♂♂, 41°32'01"N 32°35'52"E; Ulus, 300m 29.IX.2011 4♂♂, 41°34'41"N 32°38'12"E.

D i s t r i b u t i o n i n T u r k e y : Adana, Amasya, Ankara, Bolu, Bursa, Düzce, Edirne, Gaziantep, İcel, İstanbul, İzmir, Kastamonu, Kocaeli, Tekirdağ, Trabzon (KOÇAK & KEMAL 2009; KOYUNCU 2011; BEŞKARDEŞ 2012; OKYAR 2012).

***Dysstroma truncata* (HUFNAGEL, 1767)**

E x e m i n e d m a t e r i a l : Ulus, 300m 29.IX.2011 1♀, 41°34'41"N 32°38'12"E.

D i s t r i b u t i o n i n T u r k e y : Kars, Kastamonu, Rize, Trabzon (KOÇAK & KEMAL 2009; OKYAR 2010b, 2012).

***Epirrhoë galiata* (DENIS & SCHIFFERMÜLLER, 1775)**

E x e m i n e d m a t e r i a l : Amasra, Akkonak 200m 2.VIII.2011 1♂, 41°47'19"N 32°30'51"E; 50m 19.VII.2012 1♀; Çakrazova 65m 21.V.2012 1♀, 41°46'38"N 32°30'17"E; Kurucasıle, Karaman Village 130m 19.IX.2012 1♀ 3♂♂, 41°49'43"N 32°38'22"E; Üğurlu 210m 2.VIII.2011 2♀♀ 6♂♂, 41°50'08"N 32°41'40"E.

D i s t r i b u t i o n i n T u r k e y : Amasya, Ankara, Bolu, Bursa, Gümüşhane, Hatay, İstanbul, Konya, Kahramanmaraş, Muğla, Niğde (KOÇAK & KEMAL 2009).

***Euphyia biangulata* (HAWORTH, 1809)**

E x e m i n e d m a t e r i a l : Arit, Kocaköy 180m 6.IX.2010 1♂, 41°41'50"N 32°29'34"E; Kurucasıle, Paşalar 400m 22.V.2012 1♂, 41°46'47"N 32°36'28"E.

D i s t r i b u t i o n i n T u r k e y : Ardahan, Bolu, İstanbul, Ordu, Tokat (KOÇAK & KEMAL 2009; ÖZDEMİR 2016).

***Eupithecia breviculata* (DONZEL, 1837)**

E x e m i n e d m a t e r i a l : Bartın center, Şabankadı 250m 23.VI.2009 4♀♀, 41°29'38"N 32°17'42"E.

D i s t r i b u t i o n i n T u r k e y : Amasya, Ankara, Balıkesir, Bolu, Bursa, Çanakkale, Diyarbakır, Hakkari, İzmir, Kahramanmaraş, Kırıkkale, Kırklareli, Konya, Osmaniye, Siirt, Van (KOÇAK & KEMAL 2009; KEMAL & SEVEN 2013; KEMAL et al., 2017).

***Eupithecia ericeata* (RAMBUR, 1833)**

E x e m i n e d m a t e r i a l : Ulus, Hasanören 640m 29.IX.2011 1♀, 41°37'34"N 32°45'19"E.

D i s t r i b u t i o n i n T u r k e y : Bolu, Siirt (KOÇAK & KEMAL 2009; SEVEN 2014).

***Eupithecia haworthiata* DOUBLEDAY 1856**

E x e m i n e d m a t e r i a l : Amasra, Çakrazova 65m 21.V.2012 6♀♀ 1♂, 41°46'38"N 32°30'17"E; Aliobası 150m 22.V.2012 2♀♀, 41°46'39"N 32°30'59"E; Kurucasıle, Paşalar 400m 22.V.2012 1♀ 3♂♂, 41°46'47"N 32°36'28"E.

D i s t r i b u t i o n i n T u r k e y : Ordu, Northern Anatolia (MIRONOV 2003; ÖZDEMİR 2016).

***Eupithecia selinata* Herrich-Schäffer, 1861**

E x e m i n e d m a t e r i a l : Kuruçâile, Paşalar 400m 22.V.2012 1♂, 41°46'47"N 32°36'28"E.
D i s t r i b u t i o n i n T u r k e y : Ordu (ÖZDEMİR 2016)

***Eupithecia semigraphata* BRUAND, 1850**

E x e m i n e d m a t e r i a l : Amasra, Akkonak 200m 2.VIII.2011 1♂, 41°47'19"N 32°30'51"E; Kuruçâile, Karaman Village 130m 19.IX.2012 1♀ 1♂, 41°49'43"N 32°38'22"E; Bartın center, Arıönü 300m 28.IX.2011 1♀ 2♂♂, 41°37'52"N 32°12'04"E; Güzelcehisar 220m 28.IX.2011 2♀ 4♂♂, 41°39'13"N 32°13'21"E; Bartın center Port area 130m 28.IX.2011 3♂♂, 41°40'50"N 32°13'51"E; Ulus, Hasanören 640m 29.IX.2011 1♀, 41°37'34"N 32°45'19"E.

D i s t r i b u t i o n i n T u r k e y : Amasya, Ankara, Hakkari, İstanbul, Konya, Kahramanmaraş, Ordu (OKYAR & MIRONOV 2008; KOÇAK & KEMAL 2009; ÖZDEMİR 2016).

***Gymnoscelis rufifasciata* (HAWORTH, 1809)**

E x e m i n e d m a t e r i a l : Amasra, Akkonak 200m 2.VIII.2011 1♂, 41°47'19"N 32°30'51"E; 50m 19.VII.2012 1♂, 41°47'25"N 32°30'12"E; Çakrazova 65m 21.V.2012 1♀, 41°46'38"N 32°30'17"E; Hatipler 100m 2.VIII.2011 3♂♂, 41°46'39"N 32°30'18"E; Kuruçâile, Karaman Village 130m 19.IX.2012 1♀, 41°49'43"N 32°38'22"E; Uğurlu 210m 2.VIII.2011 1♀, 41°50'08"N 32°41'40"E; Bartın center, Bahçecik 244m 19.VII.2012 1♀, 41°28'49"N 32°36'41"E; Güzelcehisar 220m 28.IX.2011 1♀, 41°39'13"N 32°13'21"E; Ulus, Aşağıemirce 240m 29.IX.2011 1♀, 41°32'01"N 32°35'52"E; Hasanören 640m 29.IX.2011 1♀, 41°37'34"N 32°45'19"E.

D i s t r i b u t i o n i n T u r k e y : Adana, Amasya, Bolu, Bursa, Çanakkale, Edirne, Düzce, Hatay, İğdır, İçel, İstanbul, İzmir, Kirikkale, Konya, Mardin, Muğla, Osmaniye, Siirt (DOĞANLAR 2003; KOÇAK & KEMAL 2009; KEMAL & SEVEN 2013; KEMAL & KOÇAK 2016c).

***Horisme corticata* (TREITSCHKE, 1835)**

E x e m i n e d m a t e r i a l : Amasra, Çakrazova 65m 21.V.2012 1♂, 41°46'38"N 32°30'17"E; Kuruçâile, Paşalar 400m 22.V.2012 1♀ 3♂♂, 41°46'47"N 32°36'28"E; Bartın center, Bahçecik 244m 19.VII.2012 2♂♂, 41°28'49"N 32°36'41"E.

D i s t r i b u t i o n i n T u r k e y : Amasya, Ankara, Bolu, Bursa, Düzce, İstanbul, Kahramanmaraş (KOÇAK & KEMAL 2009).

***Horisme tersata* (DENIS & SCHIFFERMÜLLER, 1775)**

E x e m i n e d m a t e r i a l : Amasra, Aliobası 150m 22.V.2012 1♀, 41°46'39"N 32°30'59"E; Çakrazova 65m 21.V.2012 1♂, 41°46'38"N 32°30'17"E; Hatipler 100m 2.VIII.2011 1♂, 41°46'39"N 32°30'18"E; Kazpinarı 60m 15.VI.2010 1♂, 41°42'05"N 32°21'54"E.

D i s t r i b u t i o n i n T u r k e y : Amasya, Bolu, Bursa, İstanbul (KOÇAK & KEMAL 2009).

***Melanthis procellata* (DENIS & SCHIFFERMÜLLER, 1775)**

E x e m i n e d m a t e r i a l : Amasra, Akkonak 200m 2.VIII.2011 1♂, 41°47'19"N 32°30'51"E; 50m 19.VII.2012 4♂♂, 41°47'25"N 32°30'12"E; Aliobası 150m 22.V.2012 1♂, 41°46'39"N 32°30'59"E; Çakrazova 65m 21.V.2012 1♂, 41°46'38"N 32°30'17"E; Kuruçâile, Paşalar 400m 22.V.2012 1♂, 41°46'47"N 32°36'28"E; Ulus, Gökpınar 730m 29.IX.2011 1♂, 41°38'35"N 32°50'34"E.

D i s t r i b u t i o n i n T u r k e y : Bolu, Düzce, Edirne, İstanbul, Kırklareli, Ordu, Sinop, Trabzon (KOÇAK & KEMAL 2009; ÖZDEMİR 2016).

***Pasiphila rectangulata* (Linnaeus, 1758)**

Examined material: Amasra, Aliobası 150m 22.V.2012 1♂, 41°46'39"N 32°30'59"E; Kuruçışile, Paşalar 400m 22.V.2012 2♂♂, 41°46'47"N 32°36'28"E.

Distribution in Turkey: Bolu, Edirne, İğdır, Kırıkkale, Ordu, Trabzon, Van (KOÇAK & KEMAL 2009; ÖZDEMİR 2016).

***Pennithera firmata* (HÜBNER, 1822)**

Examined material: Amasra, Çakrazova 65m 21.V.2012 1♂, 41°46'38"N 32°30'17"E.

Distribution in Turkey: Bolu, Çanakkale, İstanbul (KOÇAK & KEMAL 2009).

***Thera britannica* (TURNER, 1925)**

Examined material: Ulus, Gökpınar 730m 29.IX.2011 8♀♀ 1♂, 41°38'35"N 32°50'34"E.

Distribution in Turkey: Bolu, Bursa, Düzce (KOÇAK & KEMAL 2009).

***Thera variata* (DENIS & SCHIFFERMÜLLER, 1775)**

Examined material: Ulus, Gökpınar 730m 29.IX.2011 7♀♀ 1♂, 41°38'35"N 32°50'34"E.

Distribution in Turkey: Adana, Antalya, Bolu, Düzce, İstanbul, Kastamonu, Kahramanmaraş, Sivas, Trabzon (KOÇAK & KEMAL 2009).

***Xanthorhoe ferrugata* (CLERCK, 1759)**

Examined material: Amasra, Hatipler 100m 2.VIII.2011 1♀, 41°46'39"N 32°30'18"E; Artı, Kocaköy 180m 6.IX.2010 1♀, 41°41'50"N 32°29'34"E.

Distribution in Turkey: Bursa, Bolu, Düzce, Kırklareli, Ordu, Trabzon, Zonguldak (ÖZDEMİR 2007; KOÇAK & KEMAL 2009; OKYAR 2012; ÖZDEMİR 2016).

Subfamily: Ennominae DUPONCHEL, 1845

***Apeira syringaria* (LINNAEUS, 1758)**

Examined material: Amasra, Akkonak 50m 19.VII.2012 1♂, 41°47'25"N 32°30'12"E; Bartın center, Bahçecik 244m 19.VII.2012 2♂, 41°28'49"N 32°36'41"E.

Distribution in Turkey: Bolu, İstanbul, Kırklareli, Ordu, Sinop (KOÇAK & KEMAL 2009; ÖZDEMİR 2016).

***Ascotis selenaria* (DENIS & SCHIFFERMÜLLER, 1775)**

Examined material: Amasra, Akkonak 200m 2.VIII.2011 1♀ 10♂♂, 41°47'19"N 32°30'51"E; Aliobası 150m 22.V.2012 1♂, 41°46'39"N 32°30'59"E; Hatipler 100m 2.VIII.2011 2♀♀ 12♂♂, 41°46'39"N 32°30'18"E; Kuruçışile, Curunlu 220m 2.VIII.2011 2♀♀ 1♂, 41°49'44"N 32°38'34"E; Kuruçışile, Uğurlu 210m 2.VIII.2011 3♂, 41°50'08"N 32°41'40"E.

Distribution in Turkey: Adana, Amasya, Balıkesir, Bolu, Bursa, Çanakkale, Düzce, Edirne, Hatay, İçel, İstanbul, Manisa, Kahramanmaraş, Ordu, Siirt, Sinop (DOĞANLAR 2003; KOÇAK & KEMAL 2009; SEVEN 2014; ÖZDEMİR 2016).

***Asovia maeoticaria* (ALPHÉRAKY, 1876)**

E x e m i n e d m a t e r i a l : Bartın center, Bahçecik 244m 19.VII.2012 9♂, 41°28'49"N 32°36'41"E.

D i s t r i b u t i o n i n T u r k e y : Amasya, Bolu, Kastamonu, Kırklareli, Konya, Manisa, Tokat (KOÇAK & KEMAL 2009; OKYAR 2012).

***Aspitates ochrearia* (ROSSI, 1794)**

P r e v i o u s r e c o r d : BFF Campus 10.IV.2007; Apdipaşa road 07.V.2007 (TOPER KAYGIN et al. 2009).

E x e m i n e d m a t e r i a l : Kurucasile, Karaman village 130m 19.IX.2012 1♂, 41°49'43"N 32°38'22"E; Bartın center, Ariönü 300m 28.IX.2011 1♂, 41°37'52"N 32°12'04"E.

D i s t r i b u t i o n i n T u r k e y : Amasya, Ankara, Bartın, Bursa, Çanakkale, Düzce, Diyarbakır, Edirne, Gaziantep, İçel, İstanbul, İzmir, Kırklareli, Kocaeli, Manisa, Kahramanmaraş, Kırıkkale, Kilis, Osmaniye, Siirt, Tekirdağ, Uşak, (AKBULUT et al. 2003; KOÇAK & KEMAL 2009; KOYUNCU 2011; SEVEN 2014).

***Cabera exanthemata* (SCOPOLI, 1763)**

E x e m i n e d m a t e r i a l : Amasra, Akkonak 50m 19.VII.2012 1♂, 41°47'25"N 32°30'12"E; Arit, Kocaköy 180m 6.IX.2010 1♀, 41°41'50"N 32°29'34"E; Bartın center, Kaman 50m 21.V.2012 1♂, 41°40'25"N 32°21'56"E.

D i s t r i b u t i o n i n T u r k e y : Aydın, Bursa, Edirne, Kastamonu (KOÇAK & KEMAL 2009; OKYAR 2012).

***Cabera pusaria* (LINNAEUS, 1758)**

E x e m i n e d m a t e r i a l : Amasra, Akkonak 50m 19.VII.2012 2♀♀ 8♂♂, 41°47'25"N 32°30'12"E; Çakrazova 65m 21.V.2012 1♂, 41°46'38"N 32°30'17"E; Top Deresi 234m 21.V.2012 1♂, 41°42'39"N 32°23'49"E; Kurucasile, Paşalar 400m 22.V.2012 1♀, 41°46'47"N 32°36'28"E.

D i s t r i b u t i o n i n T u r k e y : Bolu, Düzce İstanbul, Kastamonu, Kırklareli, Ordu, Tokat, Trabzon, Zonguldak (KOÇAK & KEMAL 2009; OKYAR 2012; ÖZDEMİR 2016).

***Campaea margaritaria* (LINNAEUS, 1761)**

E x e m i n e d m a t e r i a l : Amasra, Çakrazova 65m 21.V.2012 1♂, 41°46'38"N 32°30'17"E; Kazpinarı 60m 15.VI.2010 1♀, 41°42'05"N 32°21'54"E; Top Deresi 234m 21.V.2012 2♂♂, 41°42'39"N 32°23'49"E; Kurucasile, Paşalar 400m 22.V.2012 8♀♀ 22♂♂, 41°46'47"N 32°36'28"E; Ulus, Aşağıemirce 240m 29.IX.2011 1♀, 41°32'01"N 32°03'52"E; Gökpınar 730m 29.IX.2011 1♀ 2♂♂, 41°38'35"N 32°50'34"E; Hasanören 640m 29.IX.2011 1♀, 41°37'34"N 32°45'19"E.

D i s t r i b u t i o n i n T u r k e y : Amasya, Ankara, Ardahan, Artvin, Bolu, Bursa, Düzce, İstanbul, Kastamonu, Kırklareli, Ordu, Osmaniye, Sinop, Trabzon, Van, (KOÇAK & KEMAL 2009; GÖKTÜRK 2011; OKYAR 2012; ÖZDEMİR 2016).

***Charissa obscurata* (DENIS & SCHIFFERMÜLLER, 1775)**

E x e m i n e d m a t e r i a l : Arit, Kocaköy 180m 6.IX.2010 2♂♂, 41°41'50"N 32°29'34"E.

D i s t r i b u t i o n i n T u r k e y : Amasya, Bolu, Düzce, Edirne, Konya (KOÇAK & KEMAL 2009).

***Crocallis elinguaria* (LINNAEUS, 1758)**

Ex e m i n e d m a t e r i a l : Arit, Kocaköy 180m 6.IX.2010 1♂, 41°41'50"N 32°29'34"E; Kuruçışile, Curunlu 220m 2.VIII.2011 1♂, 41°49'44"N 32°38'34"E; Bartın center, Port area 130m 28.IX.2011 1♂, 41°40'50"N 32°13'51"E.

D i s t r i b u t i o n i n T u r k e y : Amasya, Bolu, Konya, Kahramanmaraş, Van (KOÇAK & KEMAL 2009).

***Ectropis crepuscularia* (DENIS & SCHIFFERMÜLLER, 1775)**

Ex e m i n e d m a t e r i a l : Amasra, Akkonak 50m 19.VII.2012 4♂♂, 41°47'25"N 32°30'12"E; Kuruçışile, Paşalar 400m 22.V.2012 1♀, 41°46'47"N 32°36'28"E.

D i s t r i b u t i o n i n T u r k e y : Bursa, Düzce (KOÇAK & KEMAL 2009).

***Ematurga atomaria* (LINNAEUS, 1758)**

P r e v i o u s r e c o r d : Catmaca 06.VIII. 2007; Kutlubey village 23.VIII.2007 (TOPER KAYGIN et al., 2009).

Ex e m i n e d m a t e r i a l : Amasra, Akkonak 200m 2.VIII.2011 2♀♀, 41°47'19"N 32°30'51"E; Aliobası 150m 22.V.2012 2♀♀, 41°46'39"N 32°30'59"E; Hatipler 100m 2.VIII.2011 4♀♀ 1♂, 41°46'39"N 32°30'18"E; Kuruçışile, Uğurlu 210m 2.VIII.2011 1♀ 1♂, 41°50'08"N 32°41'40"E.

D i s t r i b u t i o n i n T u r k e y : Amasya, Ankara, Bartın, Bilecik, Bitlis, Bolu, Bursa, Düzce, Edirne, İçel, İstanbul, Kastamonu, Kırklareli, Kocaeli, Manisa, Ordu, Sakarya, Samcun, Sinop, Sivas, Tokat, Zonguldak (KOÇAK & KEMAL 2009; TOPER KAYGIN et al. 2009).

***Ennomos fuscantaria* (HAWORTH ,1809)**

Ex e m i n e d m a t e r i a l : Arit, Kocaköy 180m 6.IX.2010 1♂, 41°41'50"N 32°29'34"E.

D i s t r i b u t i o n i n T u r k e y : Edirne, Hakkari, İstanbul (KOÇAK & KEMAL 2009; KOÇAK & KEMAL 2015; KEMAL & KOÇAK 2016).

***Ennomos quercaria* (HÜBNER, 1813)**

Ex e m i n e d m a t e r i a l : Arit, Kocaköy 180m 6.IX.2010 1♂, 41°41'50"N 32°29'34"E.

D i s t r i b u t i o n i n T u r k e y : Amasya, Bolu, İçel, İstanbul, Konya, Siirt, Tokat (KOÇAK & KEMAL 2009; SEVEN 2014).

***Ennomos quercinaria* (HUFNAGEL, 1767)**

Ex e m i n e d m a t e r i a l : Amasra, Akkonak 50m 19.VII.2012 1♂, 41°47'25"N 32°30'12"E; Bartın center, Bahçecik 244m 19.VII.2012 1♂, 41°28'49"N 32°36'41"E.

D i s t r i b u t i o n i n T u r k e y : Bolu, Düzce, İstanbul, Kırklareli, Kocaeli, (KOÇAK & KEMAL 2009; BEŞKARDEŞ 2012).

***Epione repandaria* (HUFNAGEL, 1767)**

Ex e m i n e d m a t e r i a l : Ulus, Gökpınar 730m 29.IX.2011 2♂♂, 41°38'35"N 32°50'34"E.

D i s t r i b u t i o n i n T u r k e y : İstanbul (KOÇAK & KEMAL 2009).

***Epione vespertaria* (LINNAEUS, 1767)**

Ex e m i n e d m a t e r i a l : Ulus, Gökpınar 730m 29.IX.2011 1♂, 41°38'35"N 32°50'34"E.

D i s t r i b u t i o n i n T u r k e y : Bursa, Gümüşhane, Kars, Trabzon (KOÇAK & KEMAL 2009).

***Eumannia oppositaria* (MANN, 1864)**

E x e m i n e d m a t e r i a l : Amasra, Akkonak 200m 2.VIII.2011 1♂, 41°47'19"N 32°30'51"E; Kuruçâşile, Curunlu 220m 2.VIII.2011 1♀, 41°49'44"N 32°38'34"E.

D i s t r i b u t i o n i n T u r k e y : Bursa, Konya (KOÇAK & KEMAL 2009).

***Gnophos sartata* TREITSCHKE, 1827**

E x e m i n e d m a t e r i a l : Amasra, Çakrazova 65m 21.V.2012 9♂♂, 41°46'38"N 32°30'17"E.

D i s t r i b u t i o n i n T u r k e y : Amasya, Bolu, Bursa, Çanakkale, Denizli, Hakkari, İstanbul, İzmir, Osmaniye, Kocaeli, Konya, Manisa, Kahramanmaraş, Siirt (KOÇAK & KEMAL 2009; BEŞKARDEŞ 2012; SEVEN 2014; KOÇAK & KEMAL 2015).

***Hylaea fasciaria* (LINNAEUS, 1758)**

E x e m i n e d m a t e r i a l : Bartın center, Port area 130m 28.IX.2011 1♂, 41°40'50"N 32°13'51"E; Ulus, 300m 29.IX.2011 1♂, 41°34'41"N 32°38'12"E.

D i s t r i b u t i o n i n T u r k e y : Ankara, Balıkesir, Bolu, Çanakkale, Edirne, İçel, Tekirdağ (KOÇAK & KEMAL 2009).

***Hypomecis punctinalis* (SCOPOLI, 1763)**

E x e m i n e d m a t e r i a l : Amasra, Aliobası 150m 22.V.2012 2♂♂, 41°46'39"N 32°30'59"E; Top Deresi 234m 21.V.2012 3♂♂, 41°42'39"N 32°23'49"E; Kuruçâşile, Paşalar 400m 22.V.2012 1♀ 2♂♂, 41°46'47"N 32°36'28"E.

D i s t r i b u t i o n i n T u r k e y : Amasya, Bolu, Bursa, Düzce, Hatay, Kırklareli, Ordu, Osmaniye, Sinop, Trabzon (DOĞANLAR 2003; KOÇAK & KEMAL 2009; ÖZDEMİR 2016).

***Hypomecis roboraria* ([DENIS & SCHIFFERMÜLLER], 1775)**

E x e m i n e d m a t e r i a l : Amasra, Akkonak 200m 2.VIII.2011 1♂, 41°47'19"N 32°30'51"E; 50m 19.VII.2012 4♂♂, 41°47'25"N 32°30'12"E; Aliobası 150m 22.V.2012 1♀ 1♂, 41°46'39"N 32°30'59"E; Hatipler 100m 2.VIII.2011 1♀ 1♂, 41°46'39"N 32°30'18"E; Amasra, Top Deresi 234m 21.V.2012 1♀ 8♂♂, 41°42'39"N 32°23'49"E; Kuruçâşile, Paşalar 400m 22.V.2012 1♀ 8♂♂, 41°46'47"N 32°36'28"E; Bartın center, Bahçecik 244m 19.VII.2012 1♂, 41°28'49"N 32°36'41"E.

D i s t r i b u t i o n i n T u r k e y : Artvin, Bolu, Düzce, Ordu (KOÇAK & KEMAL 2009; ÖZDEMİR 2016).

***Ligdia adustata* (DENIS & SCHIFFERMÜLLER, 1775)**

E x e m i n e d m a t e r i a l : Amasra, Çakrazova 65m 21.V.2012 4♂♂, 41°46'38"N 32°30'17"E; Kazpinarı 60m 15.VI.2010 2♂♂, 41°42'05"N 32°21'54"E; Kuruçâşile, Paşalar 400m 22.V.2012 1♀, 41°46'47"N 32°36'28"E.

D i s t r i b u t i o n i n T u r k e y : Amasya, Bolu, Bursa, Çankırı, Düzce, Edirne, İçel, Kırklareli, Rize, Trabzon (KOÇAK & KEMAL 2009; OKYAR 2010b).

***Lomaspilis bithynica* WEHRLI, 1939**

E x e m i n e d m a t e r i a l : Amasra, Akkonak 200m 2.VIII.2011 2♂♂, 41°47'19"N 32°30'51"E; Aliobası 150m 22.V.2012 1♂, 41°46'39"N 32°30'59"E; Hatipler 100m 2.VIII.2011 1♀ 1♂, 41°46'39"N 32°30'18"E; Kazpinarı 60m 15.VI.2010 1♀, 41°42'05"N 32°21'54"E; Kuruçâşile, Paşalar 400m 22.V.2012 1♀, 41°46'47"N 32°36'28"E; Bartın center, Kaman 50m 21.V.2012 1♂, 41°40'25"N 32°21'56"E.

D i s t r i b u t i o n i n T u r k e y : Ankara, Ardahan, Bolu, Düzce, İstanbul, Kastamonu, Kırklareli, Zonguldak (KOÇAK & KEMAL 2009; OKYAR 2012).

***Lomographa temerata* (DENIS & SCHIFFERMÜLLER, 1775)**

Ex e m i n e d m a t e r i a l : Amasra, Hatipler 100m 2.VIII.2011 3♂♂, 41°46'39"N 32°30'18"E; Kurucasile, Ugurlu 210m 2.VIII.2011 1♀, 41°50'08"N 32°41'40"E.

D i s t r i b u t i o n i n T u r k e y : Düzce, Ordu, Tokat (KOÇAK & KEMAL 2009; ÖZDEMİR 2016).

***Macaria alternata* (DENIS & SCHIFFERMÜLLER, 1775)**

Ex e m i n e d m a t e r i a l : Amasra, Akkonak 50m 19.VII.2012 3♂♂, 41°47'25"N 32°30'12"E; Hatipler 100m 2.VIII.2011 1♀ 5♂♂, 41°46'39"N 32°30'18"E; Bartın center, Bahçecik 244m 19.VII.2012 1♂, 41°28'49"N 32°36'41"E.

D i s t r i b u t i o n i n T u r k e y : Bursa (KOÇAK & KEMAL 2009).

***Macaria liturata* (CLERCK, 1759)**

Ex e m i n e d m a t e r i a l : Kurucasile, Paşalar 400m 22.V.2012 7♂♂, 41°46'47"N 32°36'28"E.

D i s t r i b u t i o n i n T u r k e y : Ardahan, Bolu, Düzce (KOÇAK & KEMAL 2009).

***Macaria notata* (LINNAEUS, 1758)**

Ex e m i n e d m a t e r i a l : Amasra, Akkonak 50m 19.VII.2012 3♂♂, 41°47'25"N 32°30'12"E; Aliobası 150m 22.V.2012 1♀ 3♂♂, 41°46'39"N 32°30'59"E; Kurucasile, Paşalar 400m 22.V.2012 4♀♀ 21♂, 41°46'47"N 32°36'28"E; Bartın center, Bahçecik 244m 19.VII.2012 1♂, 41°28'49"N 32°36'41"E.

D i s t r i b u t i o n i n T u r k e y : Düzce, Kırklareli, Rize, Sinop (AKBULUT et al. 2003; ÖZDEMİR 2007; CAN 2008; OKYAR 2010b)

***Nychiodes waltheri* WAGNER, 1919**

P r e v i o u s r e c o r d : Ağdacı village 10.VI.2008 (TOPER KAYGIN et al. 2009); Sipahiler-Kayadibi 12.VII.2004 1250m 1♀ (OKYAR 2012).

Ex e m i n e d m a t e r i a l : Amasra, Aliobası 150m 22.V.2012 1♀ 1♂, 41°46'39"N 32°30'59"E; Çakrazova 65m 21.V.2012 2♀♀, 41°46'38"N 32°30'17"E; Bartın center, Şabankadı 142m 23.VI.2009 1♀, 41°28'29"N 32°15'35"E.

D i s t r i b u t i o n i n T u r k e y : Bartın, Edirne, Hatay, İstanbul, Konya, Kahramanmaraş, Kırıkkale, Sivas (KOÇAK & KEMAL 2009; TOPER KAYGIN et al. 2009; ALТИPARMAK 2010; OKYAR 2012).

***Opisthograptis luteolata* (LINNAEUS, 1758)**

Ex e m i n e d m a t e r i a l : Amasra, Akkonak 50m 19.VII.2012 3♂♂, 41°47'25"N 32°30'12"E; Kurucasile, Karaman Village 130m 19.IX.2012 1♀, 41°40'25"N 32°21'56"E; Bartın center, Ariönü 300m 28.IX.2011 1♀, 41°37'52"N 32°12'04"E; Ulus, Aşağıemirce 240m 29.IX.2011 1♀ 1♂, 41°32'01"N 32°35'52"E; Gökpınar 730m 29.IX.2011 2♀♀ 2♂♂, 41°38'35"N 32°50'34"E.

D i s t r i b u t i o n i n T u r k e y : Amasya, Ankara, Artvin, Bolu, Bursa, Düzce, Çanakkale, Çankırı, İstanbul, Kastamonu, Kırklareli, Kocaeli, Manisa, Ordu, Rize, Sinop, Trabzon, Van (KOÇAK & KEMAL 2009; OKYAR 2010a, 2010b, 2012; GÖKTÜRK 2011; BEŞKARDEŞ 2012; ÖZDEMİR 2016).

***Pachycnemia hippocastanaria* (HÜBNER, 1799)**

E x e m i n e d m a t e r i a l : Amasra, Aliobasi 150m 22.V.2012 1♀ 1♂, 41°46'39"N 32°30'59"E; Çakrazova 65m 21.V.2012 9♂♂, 41°46'38"N 32°30'17"E; Hatipler 100m 2.VIII.2011 4♂♂, 41°46'39"N 32°30'18"E; Arit, Kocaköy 180m 6.IX.2010 1♀, 41°41'50"N 32°29'34"E; Kuruçâile, Karaman Village 130m 19.IX.2012 1♀ 1♂, 41°49'43"N 32°38'22"E.

D i s t r i b u t i o n i n T u r k e y : Amasya, Bolu, Bursa, Çanakkale, Düzce (KOÇAK & KEMAL 2009).

***Pachycnemia tibiaria* (RAMBUR, 1829)**

E x e m i n e d m a t e r i a l : Amasra, Akkonak 50m 19.VII.2012 1♂, 41°47'25"N 32°30'12"E; Kuruçâile, Karaman Village 130m 19.IX.2012 3♂♂, 41°49'43"N 32°38'22"E.

D i s t r i b u t i o n i n T u r k e y : This species is new record for Turkish Gaeometridae fauna.

D e s c r i p t i o n :

I m a g o : Wingspan 25-28 mm. Wings brownish grey and uniform, antemedial and postmedial lines shaped in small blackish dots, but indistinct in some. Medial area slightly darker. Outer margin of postmedial line lighter. Terminal line narrow, blackish, interrupted or fuzzy. Fringes concolorous with wings. Hindwings slightly lighter. Discal spots small, indistinct in upper side. Head, thorax and abdomen concolorous with wings (Fig. 2).

M a l e g e n i t a l i a : Uncus narrow subapically, long, broadened to apex, with two lobes and setose. Gnathos fused ventrally, upturned, apex slightly dentate. Valva broadened to apex, setose, apex round. Costa sclerotized and posterior end with short extension. Transtilla distinct. Saccus narrow. Aedeagus curved. Vesica with one long comutus (Fig. 2 a,b).



Fig. 2: Habitus of *Pachycnemia tibiaria* (RAMBUR, 1829) imago, a) male genitalia, b) aedeagus.

***Parectropis similaria* (HUFNAGEL, 1767)**

E x e m i n e d m a t e r i a l : Amasra, Aliobasi 150m 22.V.2012 1♀, 41°46'39"N 32°30'59"E; Top Deresi 234m 21.V.2012 1♂, 41°42'39"N 32°23'49"E; Kuruçâile, Paşalar 400m 22.V.2012 2♀♀ 3♂♂, 41°46'47"N 32°36'28"E.

D i s t r i b u t i o n i n T u r k e y : Bolu, Düzce, Ordu, Zonguldak (KOÇAK & KEMAL 2009; OKYAR 2012; ÖZDEMİR 2016)

***Peribatodes rhomboidaria* (DENIS & SCHIFFERMÜLLER, 1775)**

E x e m i n e d m a t e r i a l : Amasra, Akkonak 200m 2.VIII.2011 6♀♀ 10♂♂, 41°47'19"N 32°30'51"E; 50m 19.VII.2012 1♀ 1♂, 41°47'25"N 32°30'12"E; Aliobası 150m 22.V.2012 9♀♀ 8♂♂, 41°46'39"N 32°30'59"E; Çakrazova 65m 21.V.2012 1♀ 10♂♂, 41°46'38"N 32°30'17"E; Hatıpler 100m 2.VIII.2011 15♂♂, 41°46'39"N 32°30'18"E; Kazpinarı 60m 15.VI.2010 5♀♀ 3♂♂, 41°42'05"N 32°21'54"E; Kozpinarı 327m 29.IX.2011 2♀♀ 6♂♂, 41°42'41"N 32°21'50"E; Top Deresi 234m 21.V.2012 2♀♀, 41°42'39"N 32°23'49"E; Arit, Kocaköy 180m 6.IX.2010 17♂♂, 41°41'50"N 32°29'34"E; Kuruçışile, Karaman Village 130m 19.IX.2012 24♀♀ 8, 41°49'43"N 32°38'22"E; Paşalar 400m 22.V.2012 4♀♀ 5♂♂, 41°46'47"N 32°36'28"E.

D i s t r i b u t i o n i n T u r k e y : Adana, Amasya, Artvin, Bolu, Burdur, Bursa, Çanakkale, Düzce, Edirne, Hakkari, Hatay, İçel, İstanbul, Kars, Konya, Kahramanmaraş, Kastamonu, Kocaeli, Manisa, Muğla, Ordu, Osmaniye, Rize, Siirt, Sinop, Tekirdağ, Tokat, Trabzon, Van (DOĞANLAR 2003; KOÇAK & KEMAL 2009; GÖKTÜRK 2011; OKYAR 2010a; 2010b, 2012; BEŞKARDEŞ 2012; SEVEN 2014; ÖZDEMİR 2016; KEMAL et al. 2017).

***Petrophora chlorosata* (SCOPOLI, 1763)**

E x e m i n e d m a t e r i a l : Amasra, Çakrazova 65m 21.V.2012 1♂, 41°46'38"N 32°30'17"E.

D i s t r i b u t i o n i n T u r k e y : Amasya, Düzce, İstanbul Kırklareli, Ordu, Rize, Sinop, Trabzon (OKYAR & MIRONOV 2008; KOÇAK & KEMAL 2009; OKYAR 2010b; ÖZDEMİR 2016).

***Plagodis dolabraria* (LINNAEUS, 1767)**

E x e m i n e d m a t e r i a l : Amasra, Akkonak 200m 2.VIII.2011 1♂, 41°47'19"N 32°30'51"E; Çakrazova 65m 21.V.2012 1♂, 41°46'38"N 32°30'17"E; Hatıpler 100m 2.VIII.2011 1♂, 41°46'39"N 32°30'18"E; Kuruçışile, Paşalar 400m 22.V.2012 2♀♀ 16♂♂, 41°46'47"N 32°36'28"E; Bartın center, Bahçecik 244m 19.VII.2012 1♀ 1♂, 41°28'49"N 32°36'41"E.

D i s t r i b u t i o n i n T u r k e y : Bolu, Düzce, İstanbul, Ordu (ÖZDEMİR 2007, 2016; AYBERK et al. 2010).

***Rhoptria asperaria* (HÜBNER, 1817)**

E x e m i n e d m a t e r i a l : Kuruçışile, Curunlu 220m 2.VIII.2011 2♀♀ 3♂♂, 41°49'44"N 32°38'34"E; Karaman Village 130m 19.IX.2012 1♀ 1♂, 41°49'43"N 32°38'22"E.

D i s t r i b u t i o n i n T u r k e y : Bolu, Bursa, Çanakkale, Konya, Kahramanmaraş (KOÇAK & KEMAL 2009).

***Selenia dentaria* (FABRICIUS, 1775)**

E x e m i n e d m a t e r i a l : Amasra, Akkonak 50m 19.VII.2012 5♂♂, 41°47'25"N 32°30'12"E; Çakrazova 65m 21.V.2012 1♀, 41°46'38"N 32°30'17"E; Hatıpler 100m 2.VIII.2011 1♂, 41°46'39"N 32°30'18"E; Kuruçışile, Curunlu 220m 2.VIII.2011 1♀, 41°49'44"N 32°38'34"E; Karaman Village 130m 19.IX.2012 1♂, 41°49'43"N 32°38'22"E; Bartın center, Bahçecik 244m 19.VII.2012 1♂, 41°28'49"N 32°36'41"E; Güzelcehisar 220m 28.IX.2011 1♂, 41°39'13"N 32°13'21"E.

D i s t r i b u t i o n i n T u r k e y : Artvin, Bolu, Düzce Ordu (KOÇAK & KEMAL 2009; ÖZDEMİR 2016).

***Selenia lunularia* (HÜBNER, 1788)**

E x e m i n e d m a t e r i a l : Amasra, Akkonak 200m 2.VIII.2011 1♀ 7♂♂, 41°47'19"N 32°30'51"E; 50m 19.VII.2012 3♂♂, 41°47'25"N 32°30'12"E; Aliobası 150m 22.V.2012 1♂, 41°46'39"N 32°30'59"E; Hatipler 100m 2.VIII.2011 2♂♂, 41°46'39"N 32°30'18"E; Kurucasıle, Curunlu 220m 2.VIII.2011 1♀, 41°49'44"N 32°38'34"E; Uğurlu 210m 2.VIII.2011 4♂♂, 41°50'08"N 32°41'40"E; Bartın center, Bahçecik 244m 19.VII.2012 9♂♂, 41°28'49"N 32°36'41"E.

D i s t r i b u t i o n i n T u r k e y : Adana, Amasya, Ardahan, Bitlis, Bolu, Çanakkale, Düzce, İstanbul, Kocaeli, Ordu, Osmaniye, Rize, Sinop, Trabzon, Van (DOĞANLAR 2003; KOÇAK & KEMAL 2009; OKYAR 2010a, 2010b; BEŞKARDEŞ 2012; ÖZDEMİR 2016).

***Stegania dilectaria* (HÜBNER, 1790)**

E x e m i n e d m a t e r i a l : Amasra, Akkonak 200m 2.VIII.2011 2♀♀, 41°47'19"N 32°30'51"E; Aliobası 150m 22.V.2012 1♀ 1♂, 41°46'39"N 32°30'59"E; Çakrazova 65m 21.V.2012 1♀ 1♂, 41°46'38"N 32°30'17"E; Hatipler 100m 2.VIII.2011 1♀ 4♂♂, 41°46'39"N 32°30'18"E; Kazpinarı 60m 15.VI.2010 1♂, 41°42'05"N 32°21'54"E.

D i s t r i b u t i o n i n T u r k e y : Bitlis, Bolu, Düzce, Edirne, İstanbul, Konya, Kahramanmaraş, Siirt, Tokat, Van (KOÇAK & KEMAL 2009; SEVEN 2014; KEMAL & KOÇAK 2017).

***Synopsis sociaria* (HÜBNER, 1799)**

E x e m i n e d m a t e r i a l : Amasra, Çakrazova 65m 21.V.2012 1♂, 41°46'38"N 32°30'17"E; Hatipler 100m 2.VIII.2011 2♂♂, 41°46'39"N 32°30'18"E; Uğurlu 210m 2.VIII.2011 3♂♂, 41°50'08"N 32°41'40"E; Ulus 300m 29.IX.2011 5♀♀ 4♂♂, 41°34'41"N 32°38'12"E; Ariönü 300m 28.IX.2011 2♀♀, 41°37'52"N 32°12'04"E; Güzelcehisar 220m 28.IX.2011 12♀♀ 3♂♂, 41°39'13"N 32°13'21"E; Kaman Village 50m 21.V.2012 1♂, 41°40'25"N 32°21'56"E; Port area 130m 28.IX.2011 21♀ 1♂, 41°40'50"N 32°13'51"E; Şabankadı 250m 23.VI.2009 2♂♂, 41°29'38"N 32°17'42"E; Ulus, Aşağıemirce 240m 29.IX.2011 14♀♀ 20♂♂, 41°32'01"N 32°35'52"E; Gökpınar 730m 29.IX.2011 19♂♂, 41°38'35"N 32°50'34"E; Hasanören 640m 29.IX.2011 15♀♀ 4♂♂, 41°37'34"N 32°45'19"E.

D i s t r i b u t i o n i n T u r k e y : Bolu, Bursa, Çanakkale, Erzurum, İçel, İstanbul, Konya, Van (KOÇAK & KEMAL 2009).

Discussion

In this study, the Geometridae fauna of the Bartın province on the Black Sea coast was evaluated and records a total of 102 species were reached. Of these species, 9 are identified with previous published studies (TOPER KAYGIN et al. 2009; OKYAR 2012). Although the recordings were also found in present study on *Idaea degeneraria*, *Scopula imitaria*, *Aspitates ochrearia*, *Ematurga atomaria* and *Nychiodes waltheri* species, *Proteuchloris neriaria*, *Scopula rubiginata*, *Timandra comae* and *Cidaria fulvata* species were not found. 93 species were detected by this study for the first time.

Pachycnemia contains two species in the West Palaearctic region and in Northern Africa (SCOU & SIHVONEN 2015). Both of species (*Pachycnemia hippocastanaria* and *P. tibiaria*) were recorded from study area with this paper. *Pachycnemia tibiaria* is a new record for fauna of Turkey also. This species distributed in Europe only in the south from the Iberian Peninsula, southern France and Italy to Greece including Crete. In Corsica and Sardinia replaced by subsp. *benesignata*. Outside Europe the species has been found in Morocco

and Tunisia (subsp. *aindrahamaria*). In Europe a somewhat local and not common species that not often occurs in numbers. The caterpillar is recorded on *Erica scoparia*, *E. arborea*, *E. carnea* and *Calluna vulgaris*. In a variety of open land habitats with bushes and scattered trees (Mediterranean macchia) (SCOU & SIHVONEN 2015). *P. tibiaria* was recorded from Bulgaria where costal region of Black Sea were showed on map by this author. In this study, *P. tibiaria* was recorded from shore region of Black Sea where consists of local pseudomaquis habitats also. In the western Black Sea region where the study area is also located, pseudomaquis developed in ravaged forest areas can show spreading maquis elements such as *Erica arborea* (AYDINÖZÜ 2008), host of *P. tibiaria*. It is thought that *P. tibiaria* exhibits local spread in these areas.

References

- AKBULUT S., YÜKSEL B. & A. KETEN (2003): The Lepidoptera (Insecta) fauna of Düzce province, Turkey. – Turkish Journal of Zoology **27**: 257-268.
- ALТИПАРМАК Е. (2010): Determination of nocturnal Ichneumonidae, Braconidae (Hymenoptera), Geometridae (Lepidoptera) species and activation times in the research area. – Tirakya University, Institute of Science, (M. Sc. Thesis) 77p.
- Aydinözü D. (2008): An investigation on the distribution areas of the maquis formation in Turkey. – Kastamonu Education Journal **16** (1): 207-220.
- BEŞKARDEŞ V. (2012): Lepidoptera fauna of Yuvacık dam watershed in Kocaeli, Turkey. – African Journal of Agricultural Research **7** (11): 1749-1754.
- CAN F. (2008): The Geometrid Moths (Lepidoptera) from the Middle and Eastern Black Sea regions of Turkey. – Turkish Journal of Zoology **32**: 351-358.
- CAN F. (2010): Evaluation of morphological characters and male genitalia features of emerald moths (Lepidoptera: Geometridae, Geometrinae) from Turkey. – African Journal of Agricultural Research **5** (9): 867-873.
- DEMİRÖRS M. & F. KURT (2005): Contributions to the flora of the region among Zonguldak-Karabük and Bartın. – Kastamonu Education Journal **13** (2): 555-560.
- DOĞANLAR F. (2003): Systematic and Faunistic studies of Geometridae (Lepidoptera) in the East Mediterranean region of Turkey. – (Ph. D. Thesis) Cukurova University, Adana, Turkey, 275pp.
- DÖNMEZ Y. (2014): Studies on vegetation geography of Turkey. – Journal of Geography ISSN: 1302-7212, **29**: 1-27.
- GÖKTÜRK T. (2011): Lepidoptera Fauna in Artvin Province in North Black Sea Region of Turkey. – Journal of Animal and Veterinary Advances **10** (15): 1938-1946.
- HAUSMANN A. & J. VIIDALEPP (2012): Larentiinae I. In: Hausmann A. (ed.). The geometrid moths of Europe 3. – Stenstrup, Denmark, 743pp.
- HAUSMANN A. 2001: Introduction, Archiearinae, Orthostixinae, Desmobathrinae, Alsophilinae, Geometrinae. In: HAUSMANN A. (Ed.) The Geometrid Moths of Europe 1. Apollo Books, Stenstrup, 1-282.
- HAUSMANN A. (2004): Sterrhinae. In: HAUSMANN A. (ed.): The Geometrid Moths of Europe 2. – Apollo Books, Stenstrup, 600 pp., 24 pl., 237 text-figs, 420 b/w drawings, 198 maps.

- KEMAL M. & E. SEVEN (2013): Studies on the fauna and ecology of the Geometridae (Lepidoptera) in Şirvan district (Siirt Province, SE Turkey). – Priamus (Supp. 29), 1-48.
- KEMAL M. & A.Ö. KOÇAK (2015a): On the *Pterygota* fauna of Hatay Province (S. Turkey), based upon the info-system of the Cesa. – Cesa News **102**: 16-234.
- KEMAL M. & A.Ö. KOÇAK (2015b): Preliminary annotated list of the Lepidoptera of Sivas Province (East Turkey). – Cesa News **113**: 1-102.
- KEMAL M. & A.Ö. KOÇAK (2016c): On the Geometridae fauna of Bahçesaray, district, together with some morphological and eco-faunistical notes (Van Province, East Turkey) (Lepidoptera). – Priamus **14** (2):76-119, 69 figs. 32 maps.
- KEMAL M. & A.Ö. KOÇAK (2017): On the moths of Mutki district (Bitlis Province, East Turkey) (Lepidoptera). – Cesa News **150**: 1-47.
- KEMAL M., KOÇAK A.Ö. & M. AYDIN (2011): On the nocturnal spring moths of Kulp district (Diyarbakır Prov. SE Turkey) (Lepidoptera). – Cesa News **63**: 10-20.
- KEMAL M., KOÇAK A.Ö. & H. UÇAK (2017): On a collection of Lepidoptera from Dağlıca (South-East Turkey, Hakkari Province). – Cesa News **139**: 1-25.
- KOÇAK A.Ö. & S. SEVEN (1991): Faunistische Notizen über türkisch-Thrazien Lepidopteren. – Cent. ent. Stud., Misc. Pap. **10**: 4-12.
- KOÇAK A.Ö. & M. KEMAL (2009): Revised Checklist of the Lepidoptera of Turkey. – Priamus (Suppl. 17): 1-253.
- KOÇAK A.Ö. & M. KEMAL (2015): Annotated list of the Lepidoptera of Hakkari Province (SE Turkey). – Cesa News **116**: 1-147.
- KOYUNCU M.Ö. (2011): Researches on the Geometridae (Lepidoptera) fauna and systematic in Gaziantep province. – Gaziantep University, Department of Biology (M. Sc Thesis), 94pp.
- KOYUNCU M.Ö. & M. KÜTÜK (2011): Determination of Sterrhinae (Lepidoptera: Geometridae) fauna Gaziantep province. – Kafkas Univ. Fen Bil. Enst. Derg. **4** (1): 4-8.
- MANN J. (1859): Zur Lepidopterenfauna von Amasia. – Wien. ent. Monatschr. **3**: 78-106, 161-178.
- MANN J. (1861): Zur Lepidopterenfauna von Amasia . – Wien. ent. Monatschr. **5**: 155-162, 183-193.
- MIRONOV V. (2003): Larentiinae II (Perizomini and Eupitheciini). In: HAUSMANN A, editor. The geometrid moths of Europe 4. – Appollo Books, Stenstrup, 664pp.
- OKYAR Z. (2010a): Contribution to the Diurnal and Nocturnal Lepidoptera fauna of Turkish Thrace. – Trakya Univ J. Nat. Sci. **11** (2): 50-55.
- OKYAR Z. (2010b): A Study on Geometrid Moths (Lepidoptera) of the Eastern Black Sea Region (Ayder Plateau-Kaçkar Mountain-Turkey), with three new records. – Journal Of The Kansas Entomological Society **83** (3): 193–200 pp.
- OKYAR Z. & N. AKTAÇ (1999): Faunistic and taxonomic studies on the Geometridae species of Turkish Thrace. – Turkish Journal of Zoology 23, sup. 1: 99-132.
- OKYAR Z. (2012): Heteroceran (Lepidoptera) fauna of western black sea region. – Trakya Univ J Nat Sci. **13** (1): 1-14.
- OKYAR Z. & V. MIRONOV (2008): Checklist of the Geometridae of European Turkey, with new records (Lepidoptera). – Zootaxa **1789**: 1-56.
- OKYAR Z. (2012): Heteroceran (Lepidoptera) fauna of Western Black Sea Region. – Trakya Univ. J. Nat. Sci. **13** (1): 1-14.

- OKYAR Z., YURTSEVER S., AKTAÇ N. & G. ÇAKAN (2009): Some Aspects of the moth (Lepidoptera, Heterocera) species diversity in Western Black Sea Region of Turkey. – North-Western Journal of Zoology **5** (1): 104-120,
- ÖZDEMİR M. (2007): Studies on the Geometridae fauna of the provinces Bolu and Düzce in North-West Turkey (Lepidoptera). – Priamus (Suppl. 7): 1-154.
- ÖZDEMİR M. (2016): Contributions to the knowledge of Geometridae fauna (Lepidoptera) of Ordu province. – Plant Protection Bulletin, ISSN 0406-3597, **56** (2): 209–225.
- ÖZDEMİR M. (2017): Two new records for Turkish Geometridae (Lepidoptera) fauna from Ordu Province. – Turk J Zool. **41**: 185-188.
- SCHWINGENSCHUSS L. (1939a): Beitrag zur Fauna von Bithynien, besonders von Boli. – Ent. Rdsch. **56** (29): 325-328; (**32**): 358-360; (**34**): 383-384.
- SCHWINGENSCHUSS L. (1939b): Beitrag zur Fauna von Bithynien, besonders von Boli. – Ent. Z. Frankf. a. M. **49**: 273-274, 281-283.
- SCOBLE M.J. (1999): Geometrid moths of the world: a catalogue (Lepidoptera, Geometridae). – CSIRO Publishing Collingwood, Australia.
- SCOBLE M.J. & A. HAUSMANN (2007): Online list of valid available names of the Geometridae of the World. – http://www.lepbarcoding.org/geometridae/species_checklists.php. (Date of access: 09 March 2017).
- SEVEN E. & K. AKIN (2016): Contributions to the knowledge of Heterocera (Lepidoptera) fauna of Sivas province. – KSU J. Nat. Sci. **20** (4): 393-397.
- SEVEN E. (2014): Eco-faunistic studies on the Macroheterocera species in Şirvan district of Siirt (Lepidoptera). – (Ph. D. Thesis) Yüzüncüyil University, Department of Biology, Van, 427pp.
- SEVEN S. (1991): Bibliographic researches on Lepidoptera Fauna of Trace. – Priamus **6** (1/2): 1-95
- SEVEN S. (1995): Beiträge zur Schmetterlingsfauna von Thrazien. Cent. ent. Stud. – Misc. Pap. **23/24**: 1-13.
- SKOU P. & P. SIHVONEN (2015): Ennominae I. In: HAUSMANN, A., (Ed.) The Geometrid Moths of Europe 5. – Brill, Leiden: 1–657.
- TOPER KAYGIN A., YILDIZ Y. & M. Avcı (2009): Lepidoptera fauna in Bartın province, in western black sea region of Turkey. – African Journal of Agricultural Research **4** (9): 815-822.
- VARDIKJAN S.A. (1985): Atlas genital'nogo apparata pjadenits (Geometridae, Lepidoptera). Armjanskoy SSR. – Izd. AN Armjanskoy SSR, Erevan, 135pp.

Author's address:

Dr. Mustafa ÖZDEMİR
 Directorate of Plant Protection Central Research Institute,
 Gayret Mahallesi. Fatih Sultan Mehmet Bulvari. No: 66,
 06172 Yenimahalle / Ankara
 E-mail: ozdemir-mustafa@tarimorman.gov.tr

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Entomofauna](#)

Jahr/Year: 2019

Band/Volume: [0040](#)

Autor(en)/Author(s): Özdemir Mustafa

Artikel/Article: [The Geometridae \(Lepidoptera\) fauna of Bartın province in Black Sea Region, with a new record \(Pachycnemia tibaria \(Rambur, 1829\)\) for Turkey](#)
[391-413](#)