

Neoapolorryia gen. n. of the family Tydeidae from Egypt (Acari: Tydeoidea)¹⁾

MAHER E. EL-BAGOURY and FATEN M. MOMEN
(With 1 figure)

A b s t r a c t

Neoapolorryia aegyptica gen. n., sp. n. are described and illustrated from Tahrir provence region, in soil with roots of the tomato plant, **Lycopersicum esculentum** Mill.

I n t r o d u c t i o n

During our survey of mites associated with the rizosphere of root plants, **Neoapolorryia** gen. n. was found in soil with roots of the Tomato plant, **Lycopersicum esculentum** Mill. According to André 1980, 1985 and El-Bagoury 1986, the genus **Neoapolorryia** proved to be a new one. In the present work, **Neoapolorryia** gen. n. is placed near the genus **Apolorryia** André, 1980 and distinguished in having genital organotaxy (0-6-4) opposed to (0-3-3) in **Apolorryia**. In the description, André's terminology (1980, 1981) is adopted.

Genus *Neoapolorryia* gen. n.

Diagnosis: Prodorsum recurved, opisthosoma dorsal chaetotaxy: 9 (L2, h1 and h2 missing), genital organotaxy: (0-6-4); epimeral formulae: (3-1-4-2). Legs chaetotaxy: I (8-3-2-2-0) II (6-1-1-2-0) III (5-1-0-1-0) IV (5-1-0-1-0); eupathidia on tarsus I: (tc) and (P), solenidiotaxy: 2; femur IV undivided. Palp (6-1-2) + ω with a double eupathidium at the tip of the tarsus.

Type-species: **Neoapolorryia aegyptiaca** sp. n.

***Neoapolorryia aegyptiaca* sp. n.**

(Figs 1A-E)

Female: Prodorsum recurved, body length 225 µm, width 148,5 µm (fig. 1A, B), gnathosoma partially hidden, movable chela 8,8 µm shorter than the palp tarsus 15,4 µm, palpus setal pattern: (6-1-2) + ω (fig. 1E). Propodosoma with three reticulated areas separated by striations; P1 = 24,2 µm, P2 = 20,9 µm and S = 36,3 µm located in the largest area, while P3 = 24,2 µm

¹⁾ This work was carried out as parts of the project of biological control of root diseases by Mycorrhiza.




Fig. 1: *Neoapolorrya aegyptiaca* gen. n., sp. n., female. (A) Body dorsum; (B) Shape of dorsal setae and reticulation; (C) Genital region; (D) Chaerotaxy of legs; (E) Palp.

in the small one. All dorsal body setae lanceolate and serrated, except S which is filiform. Opisthosoma: dorsal chaetotaxy: 9 (L2, h1 and h2 missing), (d1-d5) = 22 µm, (L1, L4, L5) = 22 µm, ten hexagonal reticulated areas separated by striations. Genital organotaxy: (0-6-4) (fig. 1C); Epimeral formula: (3-1-4-2); coxal organ. Leg chaetotaxy (fig. 1D): I (8-3-2-2-0) II (6-1-1-2-0) III (5-1-0-1-0) IV (5-1-0-1-0); Solenidiotaxy: 2; femur IV undivided.

Male: unknown.

M a t e r i a l e x a m i n e d

Holotype: Female collected from sandy soil with roots of the tomato plant, *Lycopersicum esculentum* Mill; January 15, 1988, Tahrir Province, Egypt. Paratypes: Eight females with the above same data, two females from sandy soil with roots of the egg plant, *Solanum melongena* L.; February 10, 1988, South Sinai Province, Egypt. Holotype and 1 paratype, deposited in Zoologisches Institut und Zoologisches Museum der Universität Hamburg, Bundesrepublik Deutschland, the remaining paratypes in the National Research Centre, Plant Protection Department, Cairo, Egypt.

R e f e r e n c e s

- André, H. M., 1980: A generic revision of the family Tydeidae (Acari: Actinedida) IV. Generic descriptions, keys and conclusions. - Bull. Ann. Soc. r. belge Ent., **116**: 103-168. Bruxelles.
- André, H. M., 1981: A generic revision of the family Tydeidae (Acari: Actinedida) II. Organotaxy of the idiosoma and gnathosoma. - Acarologia, **22**: 31-46. Paris.
- André, H. M., 1985: Acari domum meliponinarum brasiliensium habitantes 10. **Melissotydeus macrosolenus** gen. n., sp. n. (Acari: Tydeidae). - Bull. Ann. Soc. r. belge Ent., **121**: 243-246. Bruxelles.
- El-Bagoury, M. E. & Abou-Awas, B. A., 1986: **Neonaudea** gen. n. of the family Tydeidae from Egypt. (Acari: Tydeoidea). - Acarologia, **27**: 121-123. Paris.

Address of the authors:

Dr. Maher E. El-Bagoury and Dr. Faten M. Momen, National Research Centre, Plant Protection Department, Dokki, Cairo, Egypt.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Entomologische Mitteilungen aus dem
Zoologischen Museum Hamburg](#)

Jahr/Year: 1990

Band/Volume: [10](#)

Autor(en)/Author(s): Momen Faten M., El-Bagoury Maher E.

Artikel/Article: [Neoapolorryia gen. n. of the family Tydeidae from
Egypt \(Acari: Tydeoidea\) 25-27](#)