

Coleoptera Europae nova

a **L. Fairmaire** descripta.

Cymindis Chaudoirii. — Long. $7\frac{1}{2}$ mill. — Oblongo-elongata, depressa, laevis, rufo-castanea, nitida, subtus testacea, antennis, ore pedibusque testaceis, capite summo obscuriore, elytris brunneis, limbo rufo-castaneo; capite lato, prothorace transverso, capite vix latiore, lateribus antice rotundatis, basi utrinque valde oblique truncato, angulis posticis obtusis, elytris apice oblique truncatis, striatis, striis laevibus. *C. canigoulensi* proxima, elytris nullo modo punctatis differt. — Sicilia.

Adelops ovoideus. — Long. 2 mill. — Breviter ovatus, convexus, rufo-testaceus, nitidus fulvo-sericans, capite prothoraceque tenuissime reticulatis, elytris postice attenuatis, tenuiter rugosulis, apice sat abrupte rotundatis, stria suturali nulla, prothorace antice late rotundato, lateribus antice arcuatim angustatis; antennis sat validis articulis 3—6 subaequalibus, gracilibus, articulis 5 ultimis sat incrassatis, articulo septimo nono longiore, articulis nono decimoque quadratis. — Gall. mer.

A. epuraeoides. — Long. $1\frac{1}{3}$ mill. — Oblongus, supra depressus, postice leviter attenuatus, testaceus, fulvo-sericans, tenuiter dense aspero-reticulatus, prothorace antice sat abrupte rotundato, elytris fere latiore, elytris oblongis, a basi leviter paulatim attenuatis, apice rotundato-subtruncatis, stria suturali antice posticeque abbreviata, antennarum articulis 3—6 aequalibus, articulo octavo transverso, parum angustiore, articulis nono decimoque transversis, undecimo oblongo, parallelo, apice acuminato. Ab *A. Aubei* differt statura majore corpore magis depresso, prothorace minus ampliato, elytris apice magis truncatis, stria suturali vix impressa. — Gall. mer.

A. subalpinus. — Long. 2 mill. — Ovato-oblongus, supra depressus, postice leviter attenuatus, testaceus, fulvo-sericans, tenuissime aspero-reticulatus, prothorace antice angustato, lateribus arcuato, elytris fere latiore, elytris oblongis, vix postice attenuatis, apice rotundatis, stria suturali medio vix impressa et utrinque stria brevi obsoletissima, antennarum articulis 3—6 parum gracilibus, clavam versus leviter incrassatis, clava haud abrupta, articulo octavo breviora at vix angustiore, nono decimoque transversim subquadratis. Ab *A. Aubei* differt statura majore, corpore magis depresso, prothorace minus amplo, antice magis angustato; a praecedente

differt statura majore, prothorace antice magis angustato et angulo suturali magis rotundato. — Alp. Gall.

Anommatus planicollis. — Long. $1\frac{1}{2}$ mill. — Oblongus, parallelus, nitidus, fulvo-testaceus, supra depressiusculus, prothorace sat dense sat fortiter punctato, elytris grosse punctato-substriatis. *A. duodecimstriato* valde affinis, sed prothorace spatio medio elevato laevi nullo differt, prothorace latiore, elytris magis fortiter striatis et statura brevior. — Gall. mer.

Amaurorhinus crassiusculus. — Long. 3 mill. — Oblongus, antice attenuatus, minus convexus, piceo-rufescens, nitidus, capite rostroque tenuiter punctatis, prothorace dense sat fortiter punctato, antice sensim attenuato, postice leviter angustato, elytris ovato-oblongis, apice rotundatis et sat abrupte declivibus, lineato-punctatis, interstitiis tenuiter vage punctulatis, interstitio secundo convexiusculo, postice magis elevato. *A. narbonensi* affinis, sed major, minus convexus, gracilior, prothorace aequaliter et fortius punctato, spatio medio laevi nullo elytrisque minus dense punctato-lineatis, interstitio secundo convexiusculo, distinguendus. — Italia.

Rhytirhinus alpicola. — Long. 5 mill. — Oblongus, sat convexus, fuscus, indumento cinereo tectus, parce luteo-setosus, rostro late sulcato, inter oculos latius, prothorace antice haud dilatato, medio late canaliculato, utrinque ad basin leviter sulcato et antice profunde impresso, elytris convexis, grosse punctato-substriatis, interstitiis alternatim magis elevatis. *R. stableaui* valde affinis, differt elytris multo minus costatis, punctis striarum multo minoribus, intervallis omnibus elevatis, rostro minus punctato, prothorace latius sulcato et in maribus angustiore, lateribus fere rectis. — Alp. Gall.

Brachyderes ophthalmicus. — Long. 10—11 mill. — Elongatus, convexus, fusco-niger, squamulis farinosis cinereo-subaureis indutus, lineolis marginalibus capitis, prothoracis elytrorumque et vittula basali, in utroque elytro, densius squamosis; capite dense punctato, oculis valde prominentibus, antennis piceis, prothorace brevi, haud impresso, ocellato-punctato, elytris ellipticis, punctato-substriatis, apice obtusis. — Andalus a.

Orthochaetes discoidalis. — Long. $2\frac{1}{3}$ mill. — Oblongus, testaceo-ruber, setis crassis albidis sparsutus, prothorace albido-pubescente, elytris disco nigricantibus; rostro nigro arcuato, tenuiter tricostulato, prothorace angusto, antice leviter attenuato, grosse punctato, elytris oblongo-ovatis, sat late punctato-striatis, intervallis aequaliter convexis, striarum punctis interdum confluentibus et obsoletis. *O. rubricato*

affinis, sed rostro magis arcuato, prothorace antice attenuato et elytrorum intervallis aequaliter convexis facile distinguendus.

— Alp. Gall.

Dichotrachelus maculosus. — Long. $4\frac{1}{2}$ mill. — Fuscus, luteo variegatus, rostro antice setoso, inter oculos penicillato, prothorace oblongo, medio valde canaliculato, margine antico et canaliculi lateribus dense pileatis, elytris striatis, striis tenuiter punctis intervallis convexiusculis, alternatim magis elevatis, setis pileatis, lateribus et postice magis dense obsitis. *D. Stierlini* affinis, sed magis oblongus, maculis evidentioribus, prothorace multo longiore, elytris basi posticeque magis attenuatis. — Alp. Gall.

Cebrio pubicornis. — Long. 16 mill. — Oblongus, convexus, ater, nitidus, obscure fulvo-pilosus, subtus cum femoribus pallide testaceus, tibiis fuscis, tarsis obscure testaceis, antennis fuscis, margine interno testaceo, dense fulvo-pubescente; capite rugoso, antice leviter impresso, antennis corpore medio multo longioribus, articulo quarto primo multo longiore, tertio secundo latiore, reliquis apice intus productis, oculis valde prominulis, prothorace brevi, dense sat tenuiter punctato, utrinque oblique impresso, angulis posticis acutis, extus haud productis, scutello oblongo, elytris dense sat tenuiter punctatis, leviter sulcatulis, interstitiis convexiusculis. *C. Fabricii* valde affinis, sed minus gibbosus, antennarum articulis 2 et 3 brevioribus, prothoracis angulis haud divaricatis, pube obscuriore, capiteque antice transversim impresso. — Lusitania.

Zwei neue Dermapteren aus Nordaustralien,

beschrieben

von **Dr. H. Dohrn.**

1. *Pygidierana Daemeli* n. sp.

Castaneo fusca, capite cum primo antennarum articulo testaceo, labro et clypeo fuscis, occipite nigro bistriato, pronoto testaceo, fusco bifasciato, scutello, alis, pedibus, pectore testaceis unicoloribus, elytris fuscis, testaceo lateribus fasciatis supra oblonge maculatis; breviter undique pilosa.

Corp. long. 19, lat. $3\frac{1}{2}$, fore. long. 4 mill. ♀.

Habitat ad Cape York Australiae borealis (Daemel).

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Entomologische Zeitung Stettin](#)

Jahr/Year: 1869

Band/Volume: [30](#)

Autor(en)/Author(s): Fairmaire Léon Marc Herminie

Artikel/Article: [Coleoptera Europae nova 231-233](#)