

FLORA.

N^o. 39.

Regensburg.

21. Oktober.

1861.

Inhalt. ORIGINAL-ABHANDLUNG. Dr. J. K. Hasskarl, Horti malabarici (Continuatio). — W. Uloth, Beiträge zur Flora der Laubmoose und Flechten in Kurhessen (Forts.). — LITTERATUR. Dr. E. Hallier, Die Vegetation auf Helgoland. M. Seubert, Lehrbuch der gesammten Pflanzenkunde. F. A. W. Miquel, Journal de Botanique Neerlandaise.

Horti malabarici clavis nova.

Autore Dr. J. K. Hasskarl.

(Continuatio cf. p. 584).

Pag.	Tab.	Tom. VIII.	Miq. Fl. Ind. Bat.
51	27	Kudici Valli = <i>Convolvulus flavus</i> Willd. DC. Prdr. IX. 415, 110, (R. S. IV. 263, 28) qui autem differt: foliis repandi-sublobatis, dentibus obtusis, nec 3-, sub 5-lobis, lobis acutis, intermedio sublobato, peduncolis folio suo multo — (nec parum) longioribus. C. agrestis Mart. DC. l. c. 405., 41 aliquot accedit; — <i>Convolvulacea dubia</i> sec. Endl. Gen. Sppl. I. 3807. Observ.; Msn. Gen. Comm. 366. — <i>Merremia convolvulacea</i> Dunst.	
53	28	Ulinja (Vlinja in Ic.) = <i>Cardiospermum Hellicacabum</i> L. (2900) β microcarpum Bl. Rmph. III. 185, 1. β . Wlp. Ann. II. 208, 1. β .	I. II. 578.
55	29	Naga-Valli s. Mandaru-Valli = <i>Bauhinia debilis</i> Hsckl. Cat. p. 287 not. Wlp. Rprt. V. 572, 1; — <i>Phanera debilis</i> Miq. Wlp. Ann. IV. 606, 24.	I. I. 69. 20.

Flora 1861.

39

Pag.	Tab.		Miq. Fl. Ind. Bat.
57	30—31	Naga-Mu-Valli = <i>Baohinia scandens</i> L. Sp. I. (2947, ubi perperam tab. 29 citatur); — <i>Bauh. anguina</i> Rxb. DC. Prdr. II. 516, 38; — <i>Lasiobema anguinum</i> Krth. Miq.	I. 1.71. 1.
59	32—34	Perim-Kaku-Valli (cf. Tom. IX, t. 77) = <i>Mimosa scandens</i> L. (7665); — <i>Entada Pursaetha</i> DC. Prdr. II. 425, 2	I. 1.45. 1.
61	35	Nai-Corana = <i>Dolichos pruriens</i> L. (5335); <i>Mucuna pruriens</i> DC. Prdr. II. 405, 4 ex parte; — <i>Mucuna prurita</i> Hook. Wlp. Rprt. I. 767, 7.	I. 1.211. 1.
63	36	Kaku Valli = <i>Dolichos urens</i> L. (5338, ubi Tom. X. p. 63 citatur); <i>D. altissimus</i> L. (5336); — <i>Mucuna gigantea</i> DC. Prdr. II. 405, 5.	I. 1.213. 5.
67	37	Pustja- (Putsja in Ic.) Paeru = sec. Dnnst. <i>Dolichos pilosus</i> Willd.?, qui autem ex Miq. differt legumine 4—5 poll. longo, quod in icone 1 poll., in descriptione nostra 2—3 poll. longum dicitur; praeterea carina in icone nostra evidenter gamopetala est, et 2-petala secundum descriptionem. — An <i>Phaseolus trinervis</i> Heyne huc pertineat?	I. 1.200.15?
69	38	Schanga Cuspi = <i>Clitoria Ternatea</i> L. DC. Prdr. II. 233, 2.	I. 1.226. 1.
71	39	Konui = <i>Abrus precatorius</i> L. (5168) DC. Prdr. II. 381, 1.	I. 1.159. 1.
73	40	Ana-Mullu = <i>Amerimnum horridum</i> Dennst. Genus <i>Amerimnum</i> P. Br. (Endl. Gen. 6701 Msn. Gen. p. 95) autem a nostra planta differt: calyce subbilabiato, legumine 1-spermo, sutura superiore subalata subrecta, inferiore valde convexa, 2-valvi, trunco inermi et foliis simplicibus. — E descriptione nostra legumen rectum linear-oblongum probabiliter indehiscens 1—3 spermium adparet, uti calyx in icone evidenter bilabiatus est. Genus <i>Dalbergia</i> maxime affine videtur, quocum et inflorescentia, fructus et habitus conveniunt.	
75	41	Paeru = <i>Dolichos Catjang</i> L. (5352, ubi sphaemate: Peru); — <i>D. sinensis</i> L. β ? Wlp. Rprt. I. 780, 21? β ., quae Vig a sinensis Savi β ex Miq.	I. 1.187. 1.

Pag.	Tab.		Miq. Fl. Ind. Bat.
79	42	Catu Paeru (Katu- in Icon.) = sec. Dennst. Phaseolus farinosus L. ?, quae species autem (cf. DC. Prdr. II. 395, 40) differt leguminibus seminibusque teretibus nec compressis, foliisque haud lobatis. — Phas. Xuaresii Zucc. Miq. I.1.194, 2. aliquot accedere videtur.	
83	43	Catu- (Katu- in Ic.) Tsjandi = Canavallia obtusifolia DC. Prdr. II. 404, 1.	I.1.215. 1.
85	44	Bara-Mareca = Canavallia ensiformis DC. Prdr. II. 404, 4. α; Hsskl. Cat. p. 276.	
87	45	Catu Bara-Mareca = Canavallia gladiata DC. β machaeroides DC. Prdr. II. 404, 3, β; Canavallia virosa W. A.	I.1.216. 3.
89	46	Tsjeria - Cametti - Valli = Derris uliginosa Bnth.	I.1.141.1.
91	47—48	Penar - Valli femina = Zanonina indica L. (7432) ♀ DC. Prdr. III. 298, 1. — Röm. Pep. 117, 1, ubi perperam p. 31—39 citatur, uti apud Miquelium similis error p. 31, tab. 39, tab. 47, 48 reperitur.	I.1.682. 1.
93	49	Penar - Valli mascula = est planta masculina praecedentis. — In contextu Rheedii perperam p. 39 loco 93 designata est.	— —
95	50	Katu Ulinu (-Ulinu. in Ic.) = Phaseolus radiatus L.; — Dolichos pubescens Willd. sec. Dennst.	I.1.197. 8.
97	51	Mu-Kelengu = Dioscorea sativa L. (7458) Bl. En. 23, 12, Kunth. En. V. 340, 17; — Miquelius sphalmate Mokelengu habet.	III. 570.
Tom. IX (de herbis).			
1	1	Tsjovanna-Areli (-Aleri in titulo paginae) s. Fula mestica incarnata = Nerium oleander L. (1715); — Nerium odorum Solnd. fl. pleno DC. Prdr. VIII. 420, 2 β (ubi uti apud Miquelium perperam Tsjovanna citatur) — fl. roseo.	II. 443.
3	2	Belutta-Arelli s. Fula mestica alba = Nerium Oleander L. (1715, ubi Areli tantum citatur); — β odoratum L. Fl. Zeyl (1715); — Nerium odorum Solnd. fl. simpl. DC. Prdr. VIII. 402, 2; — fl. albo.	— —

Pag.	Tab.		Miq. Fl. Ind. Bat.
5	3-4	Nelem-Pala (Nelam in Ic. tab. 3) cf. Tom. X. t. 49; R. et S. syst. Veg. IV. 415, 5 Observ. Wrightiam sec RBr. dubiose habent nostram; an W. dubia Sprng. DC. Prdr. VIII. 407, 12? — W. pubescens RBr. sat quadrat, sed DC. (l. c. 405, 5) hanc inter albifloras species enumerat, dum nostra floribus flavo rubentibus extus viridescentibus gaudet.	II. 432. 2.
7	5-6	Belutta-Kaka-Kodi = Chonemorpha macrophylla G. Don DC. Prdr. VIII. 430. 1.	II. 444.
9	7	Ada-Kodien = Holostemma Adakedien R. et S. VI. 95, 1, quod H. Rheedii Sprng. DC. Prdr. VIII. 532, 1. — Periploca tunicata Willd. sec. Dnnst.	II. 480. 2.
11	8	Kaka-Kodi = Marsdenia tinctoria RBr. DC. Prdr. VIII. 615, 12 (ubi Kada — scribitur)	
13	9	Kudici-Kodi = Heligme javanica Bl. Hsckl. Flor. 1845 p. 248, ibid. 1857 p. 105, 21. Bonplandia VII. p. 259. — Periploca albo-flavescens Dnnst.	II. 429. 1.
15	10	Wallja Pal-Valli = Heligme Rheedii Wight. Hsckl. Flor. 1857 p. 105, 22. Bonpl. VII. p. 260. — Periploca cordata Lam. sec. Dnnst.	II. 430. 3.
17	11	Katu-Pal Valli = Periploca mauritiana Poir. R. et S. Syst. VI. 127, 2, quae Camptocarpus mauritianus Descn. DC. Prdr. VIII. 493, 1, ubi Rheedei planta est exclusa; — Cryptolepis javanica Bl., quam ob foliorum texturam crassiusculam huc retuli; cf. Hsckl. Flor. 1857 p. 97; Bonpl. VII. 263.	II. 467. 1.
19	12	Pal - Valli = Periploca Palvalli Dnnst. — Echites malabarica Lam. DC. Prdr. VIII. 477, 166. — Chonemorpha G. Don Dichl. IV. 76	
21	13	Nansjera Patsja = Hoya pendula W. A. α Rheedei Decsn DC. Prdr. VIII. 635, 9.	
23	14	Kametti-Valli = Aganosma caryophyllata G. Don Dichl. IV. 77. DC. Prdr. VIII. 432, 1. sec. Rxb. (ubi perperam Kametti citatur.) — Ellertonia Rheedei Wight. Wlp. Ann. III. 40. 1; — Echites costata Frst. sec. Dnnst., quae haud est, cf. R. S. IV. 416, 4. DC. Prdr. VIII. 409. 4.	

Pag.	Tab.		Miq. Fl. Ind. Bat.
25	15	Wattakaka-Kodi = <i>Hoya viridiflora</i> DC. Prdr. VIII. 639, 38. — <i>Wattakaka viridiflora</i> Hsskl. Flor. 1857 p. 99. Bonplnd. VII. p. 267; <i>Cynanchum capense</i> Donst.	II. 496.
27	16	Njota - Njoden - Valli (Njodem in Titulo, — Njodien in Ic.) = <i>Ceropegia Candelabrum</i> L. (1743, ubi Niota Niodem citatur), DC. Prdr. VIII. 643, 21.	
29	17	Parparam = <i>Asclepias tetrapetala</i> Dnnst. (cf DC Prdr. VIII. 536, 1); flores tetrameri! — <i>Leuconotis</i> Jck. DC. Prdr. VIII. 33) corolla et bacca ovoidea 1—3-sperma differt.	
31	18	Neli-Tali = <i>Aeschynomene indica</i> L. (5490), DC. Prdr. II. 320, 6. Wlp. Rprt. I. 733, 8.	I. 274.1.
33	19	Todda-Vaddi (in Contextu: Valli scribitur) = <i>Oxalis Reinwardtii</i> Zucc. Wlp. Rprt. I. 476, 3. — <i>O. sensitiva</i> L. (3375). Hsskl. p. 251. <i>Biophytum sensitivum</i> DC. Prdr. I. 690, 1.	
35	20	Niti-Todda Vaddi = <i>Aeschynomene pumila</i> L. (5492, ubi — Valicitatur). <i>Mimosa virgata</i> L. (7669); <i>Desmanthus natans</i> Willd. DC. Prdr. II. 444, 2, qui est <i>Neptunia oleacea</i> Lour.	
37	21	Malam-Todda Vaddi = <i>Aeschynomene pumila</i> L. DC. Prdr. II. 321, 10. Hssk. Cat. p. 273?, quae est varietas β Ae. indicae L. Wlp. Rprt. I. 733, 8; Wlp. Ann. IV. 535. 1, β .	
39	22	Man-Todda Vaddi = sec. Dennst. <i>Aeschynomene sensitiva</i> Sw.?, quae jam foliis 16—20-jugis diversa. Nostra <i>Ormocarpus senoides</i> DC. esse videtur cf. DC. Prdr. II. 315, 3; Hsskl. Hort. Bog. I. 172.	
41	23	Aria-Veela = <i>Cleome viscosa</i> L. (4898), Schult. Syst. VII. 46, 77? Obs. — Hsskl. Cat. p. 183; sub <i>Polanisia</i> DC. Prdr. I. 242, 6, α .	
43	24	Cara Veela = <i>Cleome pentaphylla</i> L. (4894) Schlt. Syst. VII. 24, 3 Obs. — <i>Gynandropsis pentaphylla</i> DC. Prdr. I. 238, 3	
45	25	Tandale Cotti = <i>Crotalaria retusa</i> L. (5253) DC. Prdr. II. 125, 14; Wlp. Rprt. I. 586, 26.	

Pag.	Tab.		Miq. Fl. Ind. Bat.
47	26	Tandale - Cotti II = <i>Crotalaria juncea</i> L. (5251); — <i>C. sericea</i> Rtz. DC. Prdr. II. 125. 19. Wlp. Rprt. I. 586, 24.	I. 1.330.14.
49	27	Nellia-Tandale Cotti (Nella in Icon.) = <i>Crotalaria laburnifolia</i> L. (5260), DC. Prdr. II. 130, 73; Wlp. Rprt. I. 589, 53; Hsskl. Cat. p. 270.	I. 1.347.52.
51	28	Wellia-Tandale Cotti = <i>Crotalaria quinquefolia</i> L. (5263) DC. Prdr. 135, 130.	I. 1.331.17.
53	29	Pee Tandale Cotti = <i>Crotalaria verrucosa</i> L. (5256); Miquelius Tom. XX citat.	
55	30	Kattu - (Katu in Ic.) <i>Tagera</i> = <i>Indigofera hirsuta</i> L. (5553 ubi <i>perperam</i> Tom. I citatur).	I. 1.304.1.
57	31	Kondam - Pallu (-Pullu in Ic.) = <i>Impatiens oppositifolia</i> L. (6794 ?); <i>Balsamina oppositifolia</i> DC. Prdr. I. 686, 4 ?	
59	32	Nir - Murri = <i>Alysicarpus glumaceus</i> DC.; <i>Indigofera simplicifolia</i> Dnnst.	
61	33	Cupa - Vela (-Veela in Ic.) = <i>Vinca pusilla</i> Murr. DC. Prdr. VIII. 382, 3; <i>Cataranthus pusillus</i> G. Don. Diehl. IV. 95, 2; — <i>Vinca parviflora</i> Ait. sec. Dunst.	
63	34	Tajeru - Veela (Vela in Ic.) = <i>Cleome monophylla</i> L. (4907) α . <i>malabarica</i> DC. Prdr. I. 239, 17 α .	
65	35	Vallia-Capo-Molago = <i>Capsicum annum</i> L. (1496) ubi uti et in R. S. syst. Veg. IV. 559 I: Tom. II citatur; est autem <i>Capsicum longum</i> DC. γ <i>incrassatum brevipes</i> Dun. DC. Prdr. XIII. I. 424 373.	
67	36	Nir-Pullari (-Polari in Ic.) = <i>Indigofera hedyaroides</i> Lam. DC. Prdr. II. 232, 112 = (<i>I. trita</i> L. Wlp. Rprt. I. 669, 105.)	(I. 1.305.2.)
69	37	Manneli (Mannelli apud L.) = <i>Aspalathus indica</i> L. (5231); <i>Indigofera aspalathoides</i> Vhl. DC. Prdr. II. 231, 95, ubi tab. 3 cit.	I. 1.314.16.
71	38	Tajovanna - Manneli = sec. Dnnst: <i>Smithia sensitiva</i> Ait. ? quae autem ob fructus haud articulatos huc haud pertinet; — ni flores papilionacei adessent <i>Desmantho</i> , si fructus articulati essent <i>Aeschynomene</i> accederet.	
73	39	Tajeru - Manneli = <i>Dentella repens</i> Frst. (icon mala sec. Miquel.); <i>Heymia rigida</i> Dnnst.	II. 196.

Pag.	Tab.		Miq. Fl. Ind. Bat.
75	40	Suendadi-Pollu = <i>Trifolium Melilotus indica</i> L. (5638, ubi Swendadi-pulla).	
77	41	Coletta - Veetla = <i>Prionitis Hystrix</i> Miq.; <i>Barleria</i> L. (4622) DC. Prdr. XI. 237, 47. ubi pariter ac apud Miquelium Veetla cit.	II. 809.
79	42	Vada-Kodi = <i>Gendarussa vulgaris</i> Nees DC. Prdr. XI. 410, 1.	II. 831.
81	43	Adel-Adagam = <i>Justicia bivalvis</i> L. (127); <i>Adhatoda Vasica</i> Nees DC. Prdr. XI. 387, 11? Adel-Adagam apud DC., Adil-Adagam apud Miquel. citatur.	II. 829.
83	44	Katu-Karivi = <i>Andrographis Wightiana</i> Arn. Nees DC. Prdr. XI. 517, 6; — <i>Justicia atropurpurea</i> Dnnst.	
85	45	Valli-Upu-Dati = <i>Asystasia coromandelina</i> Nees β DC. Prdr. XI. 165, 7, β ; <i>Ruellia intrusa</i> Vhl. sec. Dnnst.	II. 792. 3.
87	46	Pee Tumba = <i>Andrographis echinoides</i> Nees DC. Prdr. XI. 518, 11; — <i>Justicia</i> L. (121)	II. 856.
89	47	Onapu = <i>Impatiens fasciculata</i> Lam. Wlp. Rprt. I. 474, 64; <i>Balsamina</i> DC. Prdr. I. 686, 5.	
91	48	Valli-Onapu = <i>Impatiens latifolia</i> L. (6793); <i>Balsamina</i> DC. Prdr. I. 686, 9.	
93	49	Tsjeria Onapu = <i>Impatiens Rheedei</i> W. A. Wlp. Rprt. 474, 62 <i>Balsamina Tilo</i> DC. Prdr. I. 686, 6. — <i>Imp. fasciculata</i> Lam. sec. Dnnst.	
95	50	Man-Onapu = <i>Impatiens Kleinii</i> W. A. Wlp. Rprt. I. 475. 69? <i>Balsamina minor</i> DC. Prdr. I. 686, 7. — <i>Imp. fasciculata</i> Lam. sec. Dnnst.	
97	51	Bellutta-Onapu = eadem ac praecedens.	
101	52	Tilo-Onapu s. Notenga = <i>Impatiens Balsamina</i> L. (6796, ubi — Nolengu citatur), Haskl. Cat. p. 252, ubi perperam Tom. X scriptus est; — <i>Balsamina hortensis</i> Desp.? DC. Prdr. I. 685. 1?	
103	53	Kaka - Pu = <i>Torenia asiatica</i> L. (4475), Wlp. Rprt. III. 294, 3; — mihi potius ad <i>T. hirsutam</i> Lam. DC. Prdr. X. 410, 10 spectare videtur.	
105	54	Schit Elu = <i>Sesamum indicum</i> DC. γ sub indivisum DC. Prdr. IX. 250, 1. γ ; — <i>Sesamum orientale</i> L. (4602).	II. 760.

Pag.	Tab.		Miq. Fl. Ind. Bat.
107	55	Car-Elu = forsán eadem ac praecedens sec. DC. l. c.; — <i>S. indicum</i> L. sec. Dnnst.	
109	56	Cara - Caniram (-Carinam in titulo paginae) = <i>Justicia sexangularis</i> L. (122); — <i>Andrographis paniculata</i> DC. Prdr. XI. 515, 1.	II. 855.
111	57	Tsjanga-Puspam = <i>Gratiola rotundifolia</i> L. (133, ubi Tsianga), quae Bonnaya Bnth. Wlp. Rpt. III. 290, 12; <i>Ilysanthes</i> Bnth. DC. Prdr. X. 420, 7.	
113	58	Katu-Pee-Tsjanga-Puspam = <i>Vandellia crustacea</i> Bnth. Wlp. Rpt. III. 292, 4. cf. <i>Ruellia antipoda</i> L. (4617, Observ.) — <i>Gratiola lucida</i> ? Dnnst.	
115	59	Pee-Tsjanga-Puspam (-Tjanga-Pulpani in titulo paginae et a Linnaeo et Wlp. citatur!) = <i>Ruellia antipoda</i> L. (4617). Bonnaya <i>veronicaefolia</i> Sprng. Wlp. Rpt. III. 288, 3; <i>B. brachiata</i> Luk. Otto Wlp. Rpt. III. 287, 1.	
117	60	Nelam - Parenda = <i>Viola enneasperma</i> L. (6785). <i>Jonidium</i> Vnt. α <i>malabaricum</i> DC. Prdr. 308, 7 α.	I. II. 115.1.
119	61	Katu Vistna-Clandi (Elenli in Ic.) = <i>Polygala</i> (<i>Blepharidium</i> ?) <i>Rheedei</i> Hsckl. „Herbariis teretibus tenuibus lignosis, viridibus pilosis, petiolis brevibus, foliis ovalibus s. obovatis sparse pilosis, mollibus flaccidis viridifuscis; floribus axillaribus solitariis, breviter pedunculatis bracteatis, petalis 2 rosaceo-purpureis, tertio unguiculato coeruleo cristato, crista e pilis purpureo-coeruleis“ Caet. desunt. — Habit in arenosis Malabaricae.	
121	62	Manja-Kurini = <i>Crossandra infundibuliformis</i> Nees DC. Prdr. XI. 280, 1; <i>Justicia</i> L. (112); — <i>Miquelius perperam</i> p. 191 citat.	II. 821.
123	63	Tali-Pullu = <i>Tradescantia malabarica</i> L. (2285); <i>Aneilema nudiflorum</i> R. Br. α <i>Miq. Knth. En. III. 66, 4. Hsckl. plnt. Jav. p. 95. 40.</i>	III. 537.

Cont. seq.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Flora oder Allgemeine Botanische Zeitung](#)

Jahr/Year: 1861

Band/Volume: [44](#)

Autor(en)/Author(s): Hasskarl C.

Artikel/Article: [Horti malabarici clavis nova 608-616](#)