

FLORA.

56. Jahrgang.

Nº 31.

Regensburg, 1. November

1873.

Inhalt. Dr. Karl Müller: Sechs neue Laubmose Nordamerika's. — Prof. C. Haussknecht: Beitrag zur Kenntniss der Arten von Fumaria. Fortsetz.

Sechs neue Laubmose Nordamerika's.

Beschrieben von Dr. Karl Müller Hal.

1. *Mnium (Eumnium) Neevii* C. Müll. n. sp.; *Synoicum*; cespites laxi inferne fusco-tomentosi subhumiles; *caulis simplex*, inferne fusco-radiculosus gracilis, *apice clavato solum imbricato-foliosus virescens*; *folia caulina erecto-conferta regulariter se tegentia*, crispatula, madore reflexo patula et carnosula, e basi decurrente serie cellularum angustarum unica formata breviter cuneatâ late ovata vel orbiculata, acumine brevi acuto plerumque reflexo terminata, *denticulis elongatis plus minus erectis vel reflexis ciliiformibus fimbriata*, nervo in aristulam excedente, cellulis ubique parvis hexagonis regularibus viridissimis chlorophyllosis, limbo lato flavidio; perich. pauca parva lanceolato-acuminata integerrima, limbo e cellularum serie unica efformato apice solum cincta, e cellulis elongatis laxis apice hexagonis paucis densioribus reticulata; *thecae in pedunculis flavidis flexuosis geminatae pendulae cylindrico-oblongae basi incrassatae*. Peristomium normale. Opercula et calyptrae ignotae.

Patria Oregon, Portland, vere 1873: Rever. Dr. R. Neevius; mis. C. Mohr 1873 ex Mobile Alabamae.

Notis accuratius designatis primo visu cognoscendum. *Mn. cuspidatum* affine facile distinguitur: ramis sterilibus reptantibus, fertilibus laxe et valde patule foliosis, foliis tenuissime limbatis breviter dentatis fructibusque turgide ovalibus.

2. *Catharinea (Atrichum) xanthopelma* C. Müll. n. sp.; dioica, laxe et late cespitosa *lutescens* *gracilis* crispatissima elongata, tenera; folia caulinata madore patentissima, angusta brevia lanceolata *obtusata*, lamellis 4—6 regularibus humilibus, margine undulato, limbo angustissimo apic evanido, dentibus pro more simplicibus nec duplicatis brevissimis rarius aculeiformibus, dorso nervi dentibus similibus paucissimis in pagina folii crebrioribus, cellulis parvis quadratis apicem versus hexagono-rotundis; perich. similia; theca in pedunculo elongato flavido nunquam rubente longiuscule cylindrica angustata parum curvata badia, operculo e basi cupulata longe oblique rostrato, calyptra angustissima apice dentibus ciliiformibus erectis subappressis longiuscule hispida; peristomii dentibus elongatis angustis.

Patria. Texas, Dallas County: J. Boll 1870.

A. *C. angustata* proxima caracteribus accuratius laudatis differt et species elegans. Planta mascula ignota.

3. *Bartramia (Philonotis) Mohriana* C. Müll. n. sp.; dioica; *caulis robustiusculus* ubique densifolius, inferne fusco-radiculosus, *ramulis crassiuseulis* inaequalibus plerumque brevibus et strictis nonnullis coronatus, breviusculus; folia caulinata undique inserta, madore et siccitate patenti-patula strictissima, sordide lutescentia, latiuscule perfecte lanceolata longe acuminata aristatula, longitudinaliter indistincte plicata, basi saepius cavernosula igitur irregulariter concava, margine angustissime revoluta, apice dense serrulato-denticulata inferne ob papillas prominentes nodosula, nervo profunde canaliculato in aristam excedente furcata, e cellulis angustis elongatis linearibus pellucidis laxis reticulata igitur tenuiter membranacea et flaccidula, valde punctato-papillosa; perich. caulinis similia sed basi latiore magis ovata laxissime reticulata, papillis ad parietes cellularum transversales indistinctis grossis singulis obtecta; theca in ped. flexuoso rubro robusto plantulam longitudine aequante parum obliqua magna globosa plicatula, operculo cupulato umbonato minuto; peristomium duplex normale: dentes externi late lanceolati valde et robuste trabeculati fusi striatuli, interni magis lutei asperuli breviores.

Patria. Louisiana, ad truncos putridos in sylvis opacis prope Donaldsonville, mense Decembris: C. Mohr legit et misit.

Species excellentissima, statura humiliore robusta, foliis strictissimis nunquam secundis lanceolatis laxe reticulatis maxime papillosum prima scrutatione a *B. fontana* et *calcarea* distinguenda, cum alia nunquam confundenda, *B. Schlumbergeri mexicanae*

quoad habitum aliquantulum affinis. *B. radicalis* gracilitate maxima aliisque notis multis ac habitatione terrestri palustri toto coelo differt.

4. *Barbula (Senophyllum) cancellata* C. Müll. n. sp.; dioica, late cespitosa humilis crispatissima simpliciuscula e viridi lutescens gracilis; folia caulinata madore erecto-conserata, lineali-lanceolata obtusa sed brevissime apiculata, margine infero angustissime reflexa, ubique integerrima, nervo crasso profunde canaliculato rubro dorso aspero percursa concava, basi e cellulis parvis firmiusculis regularibus rectangularibus pallidis superne minoribus quadratis obscure viridibus papillosis areolata; perich. e basi vaginata lanceolato-acuminata angustata, e cellulis elongatis laxiusculis apice magis ellipticis areolata; theca in pedunculo mediocri rubro flexuoso tenui recto cylindrica badia exannulata, operculo capsulam longitudine subaequante conico obliquiusculo, calyptra asperrima, peristomio fontinaloideo-cancellato elegantissimo tenerimo levissimo purpureo ad basin fere usque fissio.

Patria. Texas, Dallas county: J. Boll cum *Angstromia varia* associatam legit 1870.

A. *B. Crügeriana* proxima peristomii dentibus conum tenerimum fontinaloideum sistentibus appendiculatis jam longe differt.

5. *Grimmia (Dryptodon) Neevii* C. Müll. n. sp.; dioica; laxe cespitosa atro-viridis vel nigricans; *caulis* longiusculus *gracilis*, *inferne foliis diffractis nudiusculus*, per innovationem parce et breviter ramulosus, *ramulis supremis incurvulis foliosis*; *folia caulinata* appresso-imbricata vel parum patula, e basi parum decurrente oblonga *late ligulacea obtusata*, valde concava, *margine infero plus minus revoluto integerrima, apicali dentibus nonnullis remotis brevibus angulate-eroso-denticulata*, nervo latiusculo canaliculato in strias veluti dissoluto ante apicem evanido percursa, e cellulis carnosulis viridissimis parietibus undulatis instructis quadratis papillosis chlorophyllosis areolata; perich. semivaginantia oblongo-lanceolata acuminata integerrima, plicatula, exteriora evanidinervia, interiora enervia, omnia valde concava e cellulis tenuibus pellucidis rectangularibus angustis reticulata; *theca in pedunculo brevissimo curvulo majuscula ovalis* badia submacrostoma, *dentibus robustis rubris levibus in crura duo cohaerentia fassis crasse articulatis carnosulis*. Caetera ignota.

Patria, Portland Oregon, vere 1873: Rever. Dr. Neevius legit, C. Mohr ex Mobile Alabamae misit.

Planta mascula femineae simillima, androeciis nonnullis turgidis coloratis secus caulem dispositis, antheridiis magnis clavatis eparaphysatis, foliis perigonialibus e basi late ovata laxe reticulata aurantiaca breviter oblongis vel ligulaceo-obtusatis angulate denticulatis, interioribus brevissime acuminatis obtusiusculis integris, omnibus evanidinervibus. — Quoad foliorum formam *Grimmia aciculari* affinissima, sed haec species primo intuitu differt: pedunculo elongato cellulisque foliorum dolioliforme rotundis.

6. *Hypnum homalostegium* C. Müll. n. sp.; monoicum; cespites lati humiles sordide virides densiusculi rigiduli; caulis repens sed ramulis creberrimis brevibus erectis vel curvulis dense aggregatis tenuibus firmiusculis simplicibus pseudo-pinnatus; folia caulina dense imbricata indistincte secunda, madore caulem sub-julaceum turgescem tenuem sistentia, e basi cordata latiuscule ovata breviter acuminata concava, margine erecto obsolete denticulata, nervis binis brevibus, cellulis minutis tenuibus indistinctis angustissime ellipticis pallidis tenerime papillosum, alaribus ad alas solum impresso-revolutas quadratis massa grisea repletis; perich. anguste lanceolato-acuminata longiuscula secunda, concava, longitudinaliter plicatula, enervia, leviuscula lutescentia; theca in ped. ascendentem flavidо-rubro mediocri horizontalis minuta asymmetrico-cylindracea, medio valde constricta, basi gibbosa vel strumosa, macrostoma badia, operculo minuto e basi planiuscula brevissime apiculato recto; perist. dentes valde prominentes callo angustissimum sistentes madore conniventes, externi rubri valde cristati, interni lutei, laeves latiusculi carinati imperforati, ciliis singulis latis brevioribus pallidioribus interjectis.

Patria. Alabama, West-Fowl-River secus littora maritima ad corticem arborum: C. Mohr vere 1871 legit, misit 1873.

Hypno adnato proximum, sed haecce species primo momento differt: ramis multo tenuioribus nunquam turgide julaceis, foliis caulinis integerrimis, cellulis multo majoribus distincte ellipticis levissimis coloratis, perich. late ovatis acuminatis, theca longius pedunculata multo majore robustiore e collo angustiore curvato-oblonga nec gibbosa nec strumosa medio constricta et operculo magno e basi turgide conica rostellato. — Primo intuitu formis gracilioribus *Hypni cespitosi* simile, sed cellulis alaribus vesiculiformibus flavidis carens.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Flora oder Allgemeine Botanische Zeitung](#)

Jahr/Year: 1873

Band/Volume: [56](#)

Autor(en)/Author(s): Müller Karl

Artikel/Article: [Sechs neue Laubmose Nordamerika's 481-484](#)