in figure 1.

SHORT NOTE

SHORT NOTE

Table 1: Sampling localities and codes as used

81

## Diversity of 12S mitochondrial DNA sequences in Iberian and North-west African water frogs across predicted geographic barriers

Taxonomy of Palearctic water frogs has been historically controversial. North African frogs have been considered as a distinct species, Rana saharica BOULENGER, 1913 which inhabits the entire Maghreb (e.g., GÜNTHER 1991), as synonymous with Iberian R. perezi SEOANE, 1885 (STEINWARZ & SCHNEIDER 1991), or synonymous with *R*. perezi only in the part of Morocco north of the Sahara (BENHACHEM 1988). Most authors now accept R. perezi to occur in the Iberian Peninsula, R. saharica in northwestern Africa (e.g., ARANO et al. 1998), and the hypothesis that the separation of these taxa was due to the opening of the Strait of Gibraltar (BUSACK 1986). BONS & GENIEZ (1996) suggest that R. saharica might be a species complex, with different forms in northern and southern Morocco, but BUCKLEY et al. (1996) found little genetic differentiation among populations from Morocco. Based on allozyme and morphometric data ARANO et al. (1998) distinguished two subspecies of R. saharica, R. s. saharica from Algeria and R. s. riodeoroi from Morocco. They also suggest that the north-east Moroccan Moulouya river basin was the probable cause of discontinuity. Thus the major barriers to gene flow would be the Strait of Gibraltar and the Moulouya river basin. Both of these have been considered barriers for many other species of amphibians and reptiles (e.g., BUSACK 1986; ALVAREZ et al. 2000). However recent molecular studies suggest that the opening of the Strait of Gibraltar was not directly related to genetic divergences in the wall lizards of the Podarcis hispanica species complex (HARRIS et al. 2002), nor was the Moulouya river basin in Testudo graeca (LINNAEUS, 1758) (HARRIS et al. in press).

The aim of this study was to examine genetic diversity within part of the 12S rRNA mtDNA gene of water frogs across both predicted geographic barriers, and also from northern and southern populations in Morocco. This should give insight into if these are real barriers to gene flow, and if *R*.

Species	Code	Locality
Rana perezi	RP1	Laroya, SE Spain
	RP2	Vilar Pouca de Aguiar, NW Portugal
	RP3	Vilar Pouca de Aguiar, NW Portugal
Rana saharica	RS1	Kenitra, Morocco
	RS2	Debdou, Morocco
	RS3	Taza, Morocco
	RS4	Ouarzazate, Morocco
	RS5	Ketama, Morocco
	RS6	Jebel Sirwah, Morocco
	RS7	Bou Ghanem, Tunisia
	RS8	Kesra, Tunisia

saharica has genetically distinct units that could imply it is a species complex.

Genomic DNA was extracted following standard high-salt protocols. The 12S rRNA fragment was amplified by PCR using the primers published in KOCHER et al. (1989) and conditions described in HARRIS (2001). Sampling localities are given in table 1. The amplified products were sequenced on an automated sequencer (ABI 310 by Amersham Biosciences). New sequences were deposited on Genbank, accession numbers AY332762 - AY332766. Sequences were aligned including those previously published - two R. saharica from Tunis and El Fahs, Tunisia (PLÖTNER 1998), and three R. perezi (location not mentioned - MARMAYOU et al. 2000; Sierra de la Peña, Spain - PLÖTNER 1998). Three samples of R. bedriagae CAME-RANO, 1882 (Ansari Mountains, Syria; Ceyhan, Turkey; Amman, Jordan - PLÖTNER et al. 2001) were included as outgroups. Alignment was facile, as only two single base pair insertions were needed. In total, twenty sequences of 346 base pairs were included.

Phylogenetic analysis was performed using PAUP\* ver. 4.0b10 software package (SwOFFORD 2003). Using maximum parsimony of the 346 characters, 20 were informative. A 10 replicate heuristic search was performed, and node support estimated by bootstrapping (FELSENSTEIN 1985) with 1000 replicates (fig. 1). An uncorrected neighbour joining analysis gave the same estimate of relationships.

Our results support the genetic distinction between *R. perezi* and *R. saharica* 





Fig. 1: Maximum parsimony 50% Bootstrap consensus tree of relationships between sampled taxa.
Bootstrap values (1000 replicates) are given above nodes. Sequences from: \* - PLÖTNER (1998),
\*\* - MARMAYOU et al. (2000), \*\*\* - PLÖTNER et al. (2001), AY04035 - unpublished sequence from Genbank. For the localities associated with the sample codes see table 1.

as predicted by allozyme data (ARANO et al. 1998). They further show that, to a lesser extent, that there is a genetic differentiation between Moroccan and Tunisian samples of R. saharica that does not conflict with the subspecific differentiation proposed by ARANO et al. (1998). Within Morocco, however, there is almost no genetic differentiation between populations. This therefore does not support the hypothesis that R. saharica might be a species complex, and is thus similar to the data derived from protein electrophoretic analysis (BUCKLEY et al. 1996). The sample from Debdou, on the east side of the Moulouya river basin (RS2 in fig. 1) is identical to specimens from the west side using this piece of the mtDNA. Thus while there is a difference between Tunisian and Moroccan specimens, the present barrier between the two forms is not the Moulouya river basin. Extensive sampling across Algeria will be necessary to determine where the barrier does occur.

Within *R. perezi*, the sample from MARMAYOU et al. (2000) is quite distinct from the others. However, examination of the sequence shows that is has four unique mutations within the first nine base pairs of the sequence. This could be an indication that these are sequencing errors rather than true differences; similar errors have been reported in other amphibian sequences on Genbank (HARRIS 2001). Otherwise there SHORT NOTE

HERPETOZOA 16 (1/2) Wien, 30. Juli 2003

SHORT NOTE

83

is little genetic differentiation within *R*. *perezi* across the Iberian Peninsula.

To conclude, our results support the views of ARANO et al. (1998) and PLÖTNER (1998) in separating *R. perezi* from *R. saharica*. Within *R. saharica* there are two distinct clades as suggested by ARANO et al. (1998), but the geographic limit of the two taxa is not the Moulouya river as they predicted. Our results give no indication that *R. saharica* is a species complex within Morocco.

ACKNOWLEDGEMENTS: This project was supported by grants from Fundação para a Ciência e Tecnologia POCTI/41912/BSE/2001 and SFRH/BPD/ 5702/2001 (to DJH). Thanks to C. PINHo and D. BAR-BOSA (CIBIO/UP - Centro de Investigação em Biodiversidade e Recursos Genéticos / Universidade do Porto) for participation during fieldwork in North Africa, and to C. SOARES and F. SEQUEIRA (CIBIO/UP) for the samples from Vilar Pouca de Aguiar.

EFERENCES: Álvarez, Y. & Mateo, J. A. & Andreu, A. C. & Díaz-Paniagua, C. & Diez, A. & BAUTISTA, J. M. (2000): Mitochondrial DNA haplotyping of Testudo graeca on both continental sides of the Straits of Gibraltar.- J. Heredity, Oxford; 91: 39-41. ARANO, B. & LLORENTE, G. A. & MONTORI, A. & BUCK-LEY, D. & HERRERO, P. (1998): Diversification in northwest African water frogs: molecular and morphological evidence.- Herpetol. J., London; 8: 57-64. BENHACHEM, L. M. (1988): Taxinomie et charactérisation de deux espècies d'amphibiens anoures: Rana ridibunda PAL-LAS, 1771 et Bufo viridis LAURENTI, 1768, au Maroc. Doctoral Thesis, Faculty of Sciences, Rabat; 220 pp. Bons, J. & GENIEZ, P. (1996): Amphibiens et reptiles du Maroc (Sahara Occidental compris). Atlas biogéographique. Barcelona (Asociación Herpeto-lógica Española), 320 pp. BUCKLEY, D. & ARANO, B. & HERRERO, P. & LLORENTE, G. (1996): Population structure of Moroccan water frogs: genetic cohesion despite a fragmented distribution.- J. Zool. Syst. Evol. Res., Berlin; 34: 173-179. BUSACK, S. D. (1986): Biogeographic analysis of the herpetofauna separated by the formation of the Strait of Gibraltar.- National Geographic Research J., Washington; 2: 17-36. FELSENSTEIN, J. (1985): Confidence limits on phylogenies: an approach using the bootstrap.- Evolution, Lawrence; 39: 783-791. GUNTHER, R. (1991): Europäische Wasserfrösche (Anura, Ranidae) und biolo-gisches Artenkonzept.- Mitt. Zool. Mus., Berlin; 67: 39-53. HARRIS, D. J. (2001): Re-evaluation of 16S ribosomal RNA variation in Bufo (Anura: Amphibia).-Molec. Phylogen. Evol., Chicago; 19: 326-329. Harris, D. J. & Carranza, S. & Arnold, E. N. & PINHO, C. & FERRAND, N. (2002): Complex biogeo-graphic distribution of genetic variation within Podarcis Wall lizards across the Strait of Gibraltar.- J. Biogeogr., Oxford; 29(9): 1257-1262. HARRIS, D. J. & ZNARI, M. & MACE, J.-C. & CARRETERO, M. A. (in press): Genetic variation in *Testudo graeca* from Morocco estimated using 12S rRNA DNA sequencing.- Rev. Española Herpetol., Barcelona; 17. Kocher, T. D. & THOMAS, W. K. & MEYER, A. & EDWARDS, S. V. & PÄÄBO, S. & VILLABLANCA, F. X. & WILSON, A.

C. (1989): Dynamics of mitochondrial evolution in animals: Amplification and sequencing with conserved primers.- Proc. Natn. Acad. Sci. USA, Washington; 86: 6196-6200. MARMAYOU, J. & DUBOIS, A. & OHLER, A. & PASQUET, E. & TILLIER, A. (2000): Phylogenetic relationships in the Ranidae. Independent origin of direct development in the genera Philautus and Taylorana .-R. Acad. Sci., Paris; (ser. III, Sci. Vie) 323 (3), 287-297. PLÖTNER, J. (1998): Genetic diversity in mitochondrial 12S rRNA of western Palearctic water frogs (Anura, Ranidae) and implications for their systematics.- J. Zool. Syst. Evol. Res., Berlin; 36: 191-201. PLÖTNER, J. & OHST, O. & BÖHME, W. & SCHREIBER, R. (2001): Divergence in mitchondrial DNA of Near Eastern water frogs with special reference to the systematic status of Cypriote and Anatolian populations (Anura, Ranidae). Amphibia-Reptilia, Leiden; 22: 97-412. STEINWARZ, D. & SCHNEIDER, H. (1991): Distrib-ution and bioacustics of *R. perezi* SEOANE, 1885 (Amphibia: Anura: Ranidae) in Tunisia.- Bonner Zool. Beitr., Bonn; 42: 283-297. Swofford, D. L. (2003): PAUP\*: Phylogenetic Analysis Using Parsimony (and other methods) 4.0.b10. Sinauer Associates, Sunderland, Massachusetts, USA.

KEY WORDS: Amphibia: Anura: Ranidae: Rana saharica, Rana perezi, systematics, distribution, Morocco, North Africa, Iberian Peninsula

SUBMITTED: March 18, 2003

AUTHORS: D. James HARRIS, Vasco BATISTA, Miguel A. CARRETERO, Centro de Investigação em Biodiversidade e Recursos Genéticos (CIBIO\UP), ICETA, Campus Agrario de Vairão, 4485-661 Vila do Conde, Portugal. < james@mail.icav.up.pt >

## New records and natural history notes for *Pristidactylus nigroiugulus* CEI, SCOLARO & VIDELA, 2001 from Río Negro and Chubut provinces, Argentina

Lizards of the genus Pristidactylus are endemic to Argentina and Chile with ten species now recognized (CEI et al. 2001). Several of these species are poorly known, rare, and possibly endangered. Two such species, P. casuhatiensis (GALLARDO, 1968) and P. volcanensis LAMBOROT & DÍAZ, 1987, are restricted to small geographic areas of Argentina and Chile respectively, while others, such as P. fasciatus (D'ORBIGNY & BIBRON, 1837) and P. torquatus (PHILIPPI, 1861) have a large geographic range (LAMBOROT & DIAZ 1987; CEI 1986, 1993; AVILA et al. 2000). In recent years, geographically significant records were made for several species of Pristidactylus (AVILA 1994; ETHERIDGE & ESPINOZA 1997; CRUZ et al., 1999; AVILA et al. 2001) showing that the

## **ZOBODAT - www.zobodat.at**

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Herpetozoa

Jahr/Year: 2003

Band/Volume: 16\_1\_2

Autor(en)/Author(s): Harris James, Batista Vasco, Carretero Miguel A.

Artikel/Article: <u>Diversity of 12S mitochondrial DANN sequences in Iberian</u> and North-west African water frogs across predicted geographic barriers <u>81-83</u>