

SMITH 1943). These snakes could therefore pose a threat to the autochthonous small vertebrate fauna of the islands where they have been introduced. Although it is presently unknown whether *L. capucinus* will colonize undisturbed habitats on these various islands, the potential consequences of the introduction of such exotic predators on island ecosystems should not be underestimated (FRITTS & RODDA 1998).

ACKNOWLEDGMENTS: We are grateful to Frank Bambang YUWONO (Jakarta), Bulquiny BAADIL-LA (formerly of Paso, Ambon) and Bapak ELI (Labuan, Buru) for help with logistics and assistance in the field.

REFERENCES: DE HAAS, C. P. J. (1950): Checklist of the snakes of the Indo-Australian archipelago (Reptiles – Ophidia).- Treubia, Bogor; 20: 511-625. DE ROOIJ, N. (1917): The reptiles of the Indo-Australian Archipelago; II. Ophidia. Leiden (E. J. Brill), 334 pp. DUNN, E. R. (1927): Results of the Douglas Burden expedition to the islands off Komodo. II. Snakes from the East Indies.- American Museum Novitates, New York City; 287: 1-7. EDGAR, P. W. & LILLEY, R. P. H. (1993): Herpetofauna survey of Manusela National Park; pp. 131-141. In: EDWARDS, I. D. & MACDONALD, A. A. & PROCTOR, J. (Eds.): The natural history of Seram; Andover, Hampshire, UK (Intercept Ltd.). FRITTS, T. H. & RODDA, G. H. (1998): The role of introduced species in the degradation of island ecosystems: a case history of Guam.- Annual Review of Ecology and Systematics, Palo Alto; 29: 113-140. HOW, R. A. & KITCHENER, D. J. (1997): Biogeography of Indonesian snakes.- Journal of Biogeography, Oxford; 24: 725-735. HOW, R. A. & SCHMITT, L. H. & SUYANTO, A. (1996): Geographical variation in the morphology of four snake species from the Banda Arcs, eastern Indonesia.- Biological Journal of the Linnean Society, London; 59: 439-456. IN DEN BOSCH, H. A. J. (1985): Snakes of Sulawesi: a checklist, keys and additional biogeographic remarks.- Zoologische Verhandelingen, Leiden; 217: 1-50. ISKANDAR, D. T. & COLIJN, E. (2001): A Checklist of Southeast Asian and New Guinean Reptiles – Part I. Serpentes. Jakarta (Biodiversity Conservation Project, Indonesian Institute of Sciences, Japan International Cooperation Agency, The Ministry of Forestry, The Gibbon Foundation and Institute of Technology Bandung), 195 pp. ISKANDAR, D. T. & TJAN, K. N. (1996): The amphibians and reptiles of Sulawesi, with notes on the distribution and chromosomal number of frogs; pp. 39-46. In: KITCHENER, D. J. & SUYANTO, A. (Eds.): Proceedings of the first international conference on eastern Indonesian-Australian vertebrate fauna, Manado, Indonesia, November 22-26, 1994. Perth (Western Australian Museum). LEVITON, A. E. (1965): Contributions to a review of Philippine snakes, VIII. The snakes of the genus *Lycodon* H. BOIE.- Philippine Journal of Science, Manila; 94: 117-140. MANTHEY, U. & GROSSMANN, W. (1997): Amphibien & Reptilien Südostasiens. Münster (Natur und Tier-Verlag), 512 pp. MERTENS, R. (1930): Die Amphibien und Reptilien der Inseln Bali, Lombok, Sumbawa und Flores.- Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft, Frankfurt am Main; 42: 117-344. MONK, K. A. & DE FRETES, Y. & REKSODIHARJO-LILLEY, G.

(1997): The ecology of Nusa Tenggara and Maluku. Singapore (Periplus), 923 pp. SMITH, L. A. (1988): *Lycodon aulicus capucinus* a colubrid snake introduced to Christmas Island, Indian Ocean.- Records of the Western Australian Museum, Perth; 14: 251-252. SMITH, M. A. (1943): The Fauna of British India, Ceylon and Burma, including the whole of the Indo-Chinese Sub-Region; Reptilia and Amphibia Vol. III Serpentes. Reprint 1981. New Delhi (Today's & Tomorrow's Printers and Publishers), xii, 583 pp. STUEBING, R. B. & INGER, R. F. (1999): A field guide to the snakes of Borneo, Kota Kinabalu (Natural History Publications [Borneo]), 254 pp.

KEY WORDS: Squamata: Serpentes: Colubridae: *Lycodon capucinus*; geographical distribution; new island records; biogeography; invasive species; Ambon; Buru; Seram; Maluku; Indonesia

SUBMITTED: May 7, 2004

AUTHORS: Ulrich KUCH, Sektion Herpetologie, Forschungsinstitut und Naturmuseum Senckenberg, Senckenberganlage 25, D-60325 Frankfurt am Main, Germany < u.kuch@em.uni-frankfurt.de >; Jimmy A. MCGUIRE, Museum of Vertebrate Zoology, 3101 Valley Life Science Building, University of California, Berkeley, CA 94720-3160, USA < mcguirej@socrates.berkeley.EDU >.

Record of the Black-striped Snake, *Rhinoplocephalus nigrostriatus* (KREFFT, 1864), from Papua, Indonesia

The genus *Rhinoplocephalus* MÜLLER, 1885 contains six species of small to moderately sized elapid snakes found in the coastal and adjacent regions of Australia (COGGER 2002). Two of these, the Carpenteria Snake *Rhinoplocephalus boschmai* (BRONGERSMA & KNAAP-VAN MEEUWEN, 1964) and the Black-striped Snake *Rhinoplocephalus nigrostriatus* (KREFFT, 1864) also occur in southern New Guinea: The type locality of *R. boschmai* is Merauke in the southeast of the Province of Papua, Indonesia, and this species is suspected to also occur in adjacent parts of Papua New Guinea (O'SHEA 1996), in addition to its more extensive range in eastern Queensland, Australia (COGGER 2002). *Rhinoplocephalus nigrostriatus*, on the other hand, inhabits the coast and adjacent ranges of northeastern Australia from Cape York Peninsula to southern Queensland and has been collected in southern Papua New Guinea (Western Province: southern Trans-Fly, Orinomo River to Bensbach River: O'SHEA 1996), but has so far not been recorded from the Indonesian Province of Papua (DE ROOIJ

1917; DE HAAS 1950; SUPRIATNA 1995; ISKANDAR & COLIJN 2001).

In April and August 1998, two live *R. nigrostriatus* were collected in the vicinity of Merauke, Province of Papua, Indonesia, by native collectors. In their external phenotype, both snakes closely resembled Australian and Papua New Guinean specimens of *R. nigrostriatus* as described and illustrated in the literature (COGGER 2002; O'SHEA 1996). One of the specimens was preserved and deposited in the herpetological collection of the Forschungsinstitut und Naturmuseum Senckenberg, Frankfurt am Main, Germany (SMF 81869). It documents a range extension of approximately 100 km NW from collecting localities in Papua New Guinea (O'SHEA 1996: 213) and represents a new province and country record for Papua and Indonesia, respectively.

Rhinoplocephalus nigrostriatus is an extremely slender, cylindrical snake with a narrow, flattened head which is only slightly distinct from the neck, and a long tail. The maximum total length of *R. nigrostriatus* is reported to be 615 mm (O'SHEA 1996: 138), and based on their small size and the unavailability of information on their venom and toxins or cases of snakebite, these snakes are believed to be unlikely to effectively envenom humans (O'SHEA 1996), and not considered to belong to the dangerous species in Australia (SUTHERLAND & TIBBALLS 2001).

Rhinoplocephalus nigrostriatus has been described as being a secretive, nocturnal snake which feeds principally on lizards and is usually found sheltering under logs or ground litter in sclerophyll forests and woodlands (COGGER 2002). Maintained in captivity under a western Javan climate regime, the specimen from Merauke exhibited considerable activity during the day which was reminiscent of that of a diurnal forager. It also rapidly hunted down and swallowed unidentified small skinks during the day, and defended itself by swift strikes when cornered. Under identical conditions, specimens of *R. boschmai* from the same area were strictly nocturnal in agreement with the literature (COGGER 2002), and inoffensive.

ACKNOWLEDGMENTS: I am grateful to Frank Bambang YUWONO (P.T. Vivaria Indonesia, Jakarta) for providing laboratory facilities and help with logistics.

REFERENCES: COGGER, H. G. (2002): Reptiles and amphibians of Australia; 6th ed. Sydney (Reed New

Holland), 808 pp. DE HAAS, C. P. J. (1950): Checklist of the snakes of the Indo-Australian archipelago (Reptiles – Ophidia).- Treubia, Bogor; 20: 511-625. DE ROOU, N. (1917): The reptiles of the Indo-Australian Archipelago; II. Ophidia. Leiden (E. J. Brill), 334 pp. ISKANDAR, D. T. & COLIJN, E. (2001): A checklist of Southeast Asian and New Guinean reptiles – Part I. Serpentes. Jakarta (Biodiversity Conservation Project, Indonesian Institute of Sciences, Japan International Cooperation Agency, The Ministry of Forestry, The Gibbon Foundation and Institute of Technology Bandung), 195 pp. O'SHEA, M. T. (1996): A guide to the snakes of Papua New Guinea. Port Moresby (Independent Publishing), 239 pp. SUPRIATNA, J. (1995): Ular berbisa di Indonesia. Jakarta (Penerbit Bhratara), 75 pp. SUTHERLAND, S. K. & TIBBALLS, J. (2001): Australian animal toxins. Melbourne (Oxford University Press), 856 pp.

KEY WORDS: Reptilia: Squamata: Serpentes: Elapidae: *Rhinoplocephalus nigrostriatus*; venomous snakes; geographical distribution; new country record; Merauke, Papua, Indonesia; Papua New Guinea; New Guinea; Australia

SUBMITTED: May 7, 2004

AUTHOR: Ulrich KUCH, Sektion Herpetologie, Forschungsinstitut und Naturmuseum Senckenberg, Senckenberganlage 25, D-60325 Frankfurt am Main, Germany <u.kuch@em.uni-frankfurt.de>.

New distributional records of amphibians in the Andes of Ecuador

During herpetological field studies in the Andes of Ecuador, research associates of the Universidad San Francisco de Quito discovered new locations where poorly-known amphibians occurred. Little information is available for Ecuadorian amphibians and the present contribution is part of a series that intends to enhance our knowledge about the amphibians in order to promote their conservation.

Two specimens of *Eleutherodactylus sobetes* LYNCH, 1980 (DFCH- USFQ 0902, 0912) were collected at the Bosque Protector Río Guajalito, km 56 of the road Quito-San Juan-Chiriboga-Las Palmas, province of Pichincha (00°14'S / 78° 49'W, 1.900 m, February 2000) by D. F. CISNEROS-HEREDIA and A. LEÓN. These specimens represent vouchers for the second locality of the species (previously known just from the type locality, LYNCH & DUELLMAN 1997), extending its range ca. 15 km to the west. Both specimens were collected near a ravine in an old secondary evergreen low montane forest at

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Herpetozoa](#)

Jahr/Year: 2004

Band/Volume: [17_3_4](#)

Autor(en)/Author(s): Kuch Ulrich

Artikel/Article: [Record of the Black-striped Snake, *Rhinoplocephalus nigrostriatus* \(KREFFT, 1864\), from Papua, Indonesia 193-194](#)