Discovery of a population of the critically endangered Vipera darevskii VEDMEDERJA, ORLOV & TUNIYEV, 1986 in Turkey, with new elements on its identification (Reptilia: Squamata: Viperidae)

Entdeckung einer Population der stark bedrohten *Vipera darevskii* VEDMEDERJA, ORLOV & TUNIYEV, 1986 in der Türkei, mit neuen diagnostischen Einzelheiten (Reptilia: Squamata: Viperidae)

Philippe Geniez & Alexandre Teynié

KURZFASSUNG

Vipera darevskii VEDMEDERJA, ORLOV & TUNIYEV, 1986, ist eine kleine Gebirgsviper, die bisher nur von einem einzigen Fundort bekannt war, dem Berg Legli an der armenisch-georgischen Grenze. Deshalb wird die Schlange auch als in ihrem Bestand stark gefährdet eingestuft. Wir berichten über ein weiteres Vorkommen in der Nordost-Türkei, dessen Entdeckung Anlaß zur Hoffnung gibt, daß die Art tatsächlich noch weiter verbreitet ist. Die Ähnlichkeit von V. darevskii mit der häufigeren V. [renardi] eriwanensis (REUSS, 1933) und der Mangel publizierter Daten über ihre diagnostischen Merkmale mag zum Teil für das Fehlen von Nachweisen zwischen den nummehr bekannten Fundorten verantwortlich sein. Neue bestimmungswichtige Merkmalsausprägungen werden beschrieben.

ABSTRACT

Darevsky's Viper Vipera darevskii VEDMEDERJA, ORLOV & TUNIYEV, 1986 is a small mountain viper previously known from a single locality, Mt Legli, on the Armeno-Georgian border. Consequently it is classified as "critically endangered". A new locality is reported here in north-eastern Turkey, giving hope that the species is actually more widespread. Similarity of V. darevskii with the more common V. [renardi] eriwanensis (REUSS, 1933) and scarcity of published information on its diagnostic characters may be partly responsible for the lack of records between these two localities. New elements on the identification of this species are given here.

KEY WORDS

Reptilia: Squamata: Serpentes: Viperidae; Vipera darevskii, Turkey, Zekeriya, chorology

INTRODUCTION

The poorly known *Vipera darevskii* VEDMEDERJA, ORLOV & TUNIYEV, 1986 is one of the rarest of the true vipers. It is presently known from a single locality in north-western Armenia, Mount Legli in the south-eastern part of the Javakheti range, on subalpine and alpine meadows between 2600 and 3000 m a.s.l. (VEDMEDERJA et al. 1986; ORLOV & TUNIYEV 1990), probably extending into the adjacent Georgian territory. Consequently, the species is classified as critically endangered by the IUCN (http:// www.redlist.org/search/details.php?species= 23000).

Published information on this species is scarce. The Mount Legli population was first documented by DAREVSKY (1956) under the name Vipera kaznakovi dinniki NIKOLSKY,

1913. Based on nine specimens, three of which are pictured in black and white photographs, VEDMEDERJA et al. (1986) described these animals as a new species. They provided comparisons of the principal features of this viper with V. kaznakovi NIKOLSKY, 1909 and V. dinniki NIKOLSKY, 1913 but not with the taxa of *V. ursinii* (BONAPARTE, 1835) sensu lato, as they considered the species to be part of the V. kaznakovi complex. ORLOV & TUNIYEV's (1990) synopsis on Caucasian vipers, combines material previously published by VEDMEDERJA et al. (1986) and additional information including a picture of a live specimen. HÖGGREN et al. (1993) provide little new information, while the voluminous synthesis of NILSON & ANDRÉN (2001) on the V. ursinii complex does not really cover *V. darevskii* as these authors do not include it in the *ursinii* complex.

For a long time, the phylogenetic affinities of *V. darevskii* have been unclear. It was usually considered a member of the kaznakovi complex, but often regarded as "intermediate" between the "ursinii" and the "kaznakovi" complexes. What was meant by "intermediate" is unclear, but a hybrid origin has been postulated (ORLOV & TUNIYEV 1990). "Intermediate" probably reflected the fact that its characters do not allow a precise allocation. NILSON & ANDRÉN (2001), for example, noted that its "external morphology evolved as typical for many alpine taxa within the *ursinii* complex, but shows clear similarities also with members of the Vipera kaznakovi group". The genetic results (JOGER et al. 2002; KALYABINA-HAUF 2004) suggest that the "ursinii complex" and the "kaznakovi complex" do not form natural groups in mitochondrial phylogeny, but that the taxa included in these assemblages constitute a monophyletic clade relative to other vipers; in their study, V. darevskii appears as a well differentiated species within this kaznakovi – ursinii complex.

The viper fauna of the Anatolian-Caucasian region is characterized by its high species richness: the *kaznakovi – ursinii* complex is represented by *anatolica* EISELT & BARAN, 1970, known from one or two localities in the calcareous mountain of Elmadağ (south-western Turkey), *eriwanensis* REUSS, 1933 from the basaltic high plateaux of north-eastern Turkey and Armenia, extending to adjacent Azerbaijan, lotievi Nilson, Tuniyev, Orlov, Höggren & ANDRÉN, 1995 from high altitude habitats along the ridge of the Greater Caucasus Mountains, *renardi* (CHRISTOPH, 1861) with a wide distribution in Eurasian steppes, reaching the low altitude steppe areas of the northern slopes of the Caucasus, ebneri KNOEPFFLER & SOCHUREK, 1955 from the high altitude areas of the Elbruz mountain range in NW Iran and SE Azerbaijan (NILSON et al. 2001), kaznakovi from forested habitats in the eastern Black Sea coast and western Caucasus, dinniki from open habitats at higher altitudes in the western and central Caucasus, and orlovi TUNIYEV & OSTROVSKIKH, 2001 and magnifica TUNIYEV & OSTROVSKIKH, 2001 from the western Black Sea coast (TUNIYEV & OSTROVSKIKH 2001). The other Vipera species in the area, Vipera [berus] barani BÖHME & JOGER, 1983 along the Black Sea coast of Turkey and V. ammodytes transcaucasiana Bou-LENGER, 1913 in northern Turkey, Georgia and NW Azerbaijan, do not belong to the kaznakovi – ursinii clade. The enigmatic V. pontica Billing, Nilson & Sattler, 1990, known from two specimens from NE Turkey, was recently suggested to be closely related to, or even identical with, barani (BARAN et al. 2001). However, we strongly disagree with this classification because the holotype of this taxon shows clearly morphological features of V. kaznakovi and V. a. transcaucasiana, which are both known to occur in the same area, but not of Vipera berus (LINNAEUS, 1758) s.l.

RESULTS

The authors discovered a population of vipers of the *ursinii-kaznakovi* complex in a new site situated 2 km east of Zekeriya village, ca 26 km south-east of Ardanuç, in the Artvin province, at 1970-2070 m. Six vipers were found on 18.9.2000 and two on 1.6.2002, in total three males, four females and one newborn. Three specimens were collected and deposited in the Muséum national d'Histoire naturelle of Paris (MNHN 2002.410, male), in the "Alcide d'Orbigny" collection (ASHNAO.00102, female) and in the Laboratoire de Biogéographie et Ecologie des Vertébrés de l'Ecole Pratique des Hautes Etudes, in Montpellier (BEV.8369, juvenile). All specimens were photographed and the pictures are kept in the picture collection of PG (numbers PGe 449-456).

Description

The Zekeriya vipers were easily assigned to the *kaznakovi – ursinii* complex by the general habitus and colour pattern (see figures), rounded snout and presence of large unfragmented frontal and parietal

26

Another population of the critically endangered Vipera darevskii



Fig. 1: Adult male *Vipera darevskii* VEDMEDERJA, ORLOV & TUNIYEV, 1986 from Zekeriya, Turkey (MNHN 2002.410). Photograph: Ph. GENIEZ & A. TEYNIÉ.
Abb. 1: Adulte männliche *Vipera darevskii* VEDMEDERJA, ORLOV & TUNIYEV, 1986 aus Zekeriya, Türkei (MNHN 2002.410). Photo: Ph. GENIEZ & A. TEYNIÉ.



Fig. 2: Adult female *Vipera darevskii* VEDMEDERJA, ORLOV & TUNIYEV, 1986 from Zekeriya, Turkey (ASHNAO.00102). Photograph: Ph. GENIEZ & A. TEYNIÉ.
Abb. 2: Adulte weibliche *Vipera darevskii* VEDMEDERJA, ORLOV & TUNIYEV, 1986 aus Zekeriya, Türkei (ASHNAO.00102). Photo: Ph. GENIEZ & A. TEYNIÉ.

Table. 1: Main morphological features of our eight Turkish specimens of *Vipera darevskii* VEDME-DERJA, ORLOV & TUNIYEV, 1986. SVL - snout-vent length (mm); TL - tail length (mm); Dors - number of dorsal scales rows at midbody; Vent - number of ventral plates; SC - number of pairs of subcaudal plates; Apic - number of apical scales; Supl - number of supralabial scales on one side; Circ - number of circumocular scales on one side; * - tail incomplete.

Tab. 1: Morphologische Hauptmerkmale unserer acht türkischen Exemplare von Vipera darevskii VEDMEDERJA, ORLOV & TUNIYEV, 1986. SVL - Kopf-Rumpf-Länge (mm); TL - Schwanzlänge (mm); Dors -Anzahl Rückenschuppenlängsreihen in Rumpfmitte; Vent - Anzahl Bauchschienen; SC - Anzahl Unterschwanzschildpaare; Apic - Anzahl Apicalia; Supl -Anzahl Supralabialia einer Seite; Circ - Anzahl Circumorbitalia einer Seite; * - Schwanz unvollständig.

SVL	TL	Dors	Vent	SC	Apic	Supl	Circ
345	60	19	138	38	2	9	9
353	40	19	144	33	1	9	10
390	50	19	143	31	2	7	10
360	50	19	136	33	2	9	9
330	35	19	133	26	2	8	10
109	22	21	137	39	2	8	10
390	60*	21	129	29*	2	9	9
340	40	21	139	31	2	8	10

scales on the pileus. The only vipers of this complex known from NE Turkey at the time were *eriwanensis* and *kaznakovi*. The observed vipers nevertheless looked different from these taxa, which we had seen in the field several times before (see below). The principal features of these vipers are the small size (max. 45 cm, cf. table 1), 19 (in 5 specimens) or 21 (in 3 specimens) longitudinal rows of dorsal scales at midbody, a relatively big head with rather angular canthus rostralis, one series of large oval vertical dark blotches on each flank, often two bright yellowish dorso-lateral stripes (greyer in some females), sometimes with clearcut anterior edge as in Vipera kaznakovi and V. dinniki (figs. 1 to 3), a dark to very dark pileus, pale labials with dark sutures in all but one specimen, and a mostly dark belly, dark-grey with numerous black dots and whitish scale edges. The general tint of the body is yellowish in males with a dark vertebral zigzag band, greyer in females with a brown zigzag band, sometimes connected with the dark headed-pattern (on 3 specimens). The newborn specimen was very small (total length 13.5 cm). The venom of all specimens was yellow. Additional scale characters are given in table 1.

Comparisons

In order to establish the identity of these vipers, we compared their characters with the information available on the morphology of the other viper taxa of the region, and with our own observations of specimens of these taxa in captivity, in the field, and from published and unpublished color photographs.

Most of the other vipers of the kaznakovi – ursinii complex known from this region could be easily excluded by these characters: kaznakovi and the closely related orlovi and magnifica live in very different habitats, have a very different color pattern, a relatively larger head, a different head shape with more pronounced canthus rostralis, more perioculars (11-12 [NILSON et al. 1995] versus 9-10), dinniki has a gold-edged iris (NILSON et al. 1995) and a head shape rather similar to kaznakovi (ORLOV & TUNIYEV 1990; pers. obs.), with a very pro-

- Mount Legli locality in Armenia; * the new site near Zekeriya, Turkey.
 - Abb. 4: Geographische Verbreitung von Vipera darevskii.

Fig. 5: Female Vipera eriwanensis (REUSS, 1933) from Arpaçay, NE Turkey, (PGe.439).

Abb. 5: Weibchen von Vipera eriwanensis (REUSS, 1933) von Arpaçay, NE Türkei, (PGe.439).

Figs. 3 - 6 (opposite page) / Abb. 3 - 6 (gegenüberliegende Seite): Vipera darevskii VEDMEDERJA, ORLOV & TUNIYEV, 1986 and Vipera eriwanensis (REUSS, 1933). Photos: PH. GENIEZ & A. TEYNIÉ.

Fig. 3: Adult female Vipera darevskii from Zekeriya, Turkey (PGe.451).

Abb. 3: Adulte weibliche Vipera darevskii von Zekeriya, Türkei (PGe.451).

Fig. 4: Geographical distribution of Vipera darevskii.

Mount Legli Fundort in Armenien; * - der neue Fundort bei Zekeriya, Türkei.

Fig. 6: Male Vipera darevskii from Zekeriya, Turkey (PGe.450).

Abb. 6: Männchen von Vipera darevskii von Zekeriya, Türkei (PGe.450).

Another population of the critically endangered Vipera darevskii

5









minent canthus rostralis above the nostril (pers. obs.). In addition, the Zekeriya vipers are smaller than all these taxa (compare table 1 with data in ORLOV & TUNIYEV 1990) and the specimens with pale pileus have a fronto-parietal ocellated spot, which is absent in the other taxa (see NILSON & ANDRÉN 2001).

In *lotievi*, *anatolica* and *ebneri* there is almost always one apical scale (NILSON & ANDRÉN 2001) versus two in six of the Zekeriya vipers (table 1), the upper preocular is generally in contact with the nasal (NIL-SON & ANDRÉN 2001; no contact in our eight specimens), the belly is white, the body coloration grey even in males (yellowish in males of the Zekeriya vipers), the pileus is generally pale (dark in four Zekeriya specimens), there is usually no dark suture on the labials and the dark dorsal zigzag band is never in contact with the dark parts of the pileus (present in three of our specimens). These latter two characters distinguish the Zekeriya vipers from *renardi*, which on its part has clear marks on labials and throat, a single apical scale and in general the upper preocular and nasal in contact with each other. When following the key in NILSON & ANDRÉN (2001), the specimens characterized by having 21 dorsal scale rows are eriwanensis.

Thus, correct determination would have to decide between the two taxa whose distribution areas come closest to this new locality: *eriwanensis* (known from this region, but not from this locality) and *darevskii*, whose known range lies just some 130 km from Zekeriya (fig. 4). In the field the general impression of a snake similar to *Vipera ursinii* s.l. but with a larger head, more robust body and yellowish coloration in males strongly spoke in favour of *Vipera darevskii*. Unfortunately, we were unable to obtain or look at any specimens of *darevskii* for comparison from the collections of the Zoological Institute, Russian Academy of Sciences, St Petersburg and Naturhistoriska Museet, Göteborg, so identification had to be based on published information and pictures.

There is little published information on the characters which separate eriwanensis from darevskii. Nilson & Andrén (2001) mention that in their key *darevskii* will fall out as eriwanensis (as was the case in the Zekeriya vipers). ORLOV & TUNIYEV (1990) mention that *darevskii* has, compared with *eriwanensis*, (1) a relatively high head with a much less pointed upper anterior snout edge, (2) yellowish general coloration, (3) clear contrasted pattern and (4)different pholidosis, but no information is given on these pholidotic characters. Compared to eriwanensis, our Zekeriya vipers have clearly a larger head, with a wider and more rounded snout, and males have a yellowish coloration with a more contrasted zigzag band (pure black) in males. This would thus correspond to characters of darevskii.

In addition, differences in color pattern are also apparent: in the majority of the Zekeriya vipers the flanks have a series of large dark blotches (usually two series of smaller dots in *eriwanensis*, fig. 5) (pers. obs.) and the neck pattern is more contrasted, with clear-cut dark lines and a wide intrusion of the pale flank coloration onto the sides of the neck (compare both taxa in figs. 6 and 7). In some specimens, the dark vertebral zigzag band meets the dark coloration of the pileus, a condition that we

Figs. 7 - 10 (opposite page) / Abb. 7 - 10 (gegenüberliegende Seite): Vipera darevskii VEDMEDERJA, ORLOV & TUNIYEV, 1986 and Vipera eriwanensis (REUSS, 1933).

Photos: 7, 8, 10 - PH. GENIEZ & A. TEYNIÉ; 9 - P.-A. CROCHET.

Fig. 7: Male Vipera eriwanensis from Arpaçay, NE Turkey (PGe.438). Abb. 7: Männchen von Vipera eriwanensis von Arpaçay, NE Türkei (PGe.438).

Fig. 8: Ventral view of a male Vipera darevskii from Zekeriva, Turkey (PGe.450).

Abb. 8: Ventralansicht eines Männchens von Vipera darevskii von Zekeriya, Türkei (PGe.450).

Fig. 9: Ventral view of a male Vipera eriwanensis

from 4 km north of Ölçek on the road to Posof, NE Turkey (PGe.440).

Abb. 9: Ventralansicht eines Männchens von Vipera eriwanensis

von 4 km nördlich Ölçek, an der Straße nach Posof, NE Türkei (PGe.440).

Fig. 10: Habitat of Vipera darevskii in Zekeriya, Turkey.

Abb. 10: Lebensraum von Vipera darevskii in Zekeriya, Türkei.

Another population of the critically endangered Vipera darevskii







never observed in *eriwanensis*. Some specimens (figs. 1, 2, 6) have a coloration that is strikingly different from any *eriwanensis*, while others (fig. 3) look more similar but with a more contrasted head and neck pattern. The ventral coloration of the Zekeriya vipers is grey to dark grey with numerous black dots and whitish scale edges (fig. 8). The impression is of a dark belly, very different from the mostly pale belly of *eriwanensis* (fig. 9). Last, the corners of the dorsal band windings are less rounded in the Zekeriya vipers than in *eriwanensis*.

Although these differences are not listed in the few published accounts of *darevskii*, they correspond well with the published pictures. The dark belly is shown in figures 10 and 11 of ORLOV & TUNIYEV (1990), as well as the series of large dark blotches on the flanks and the contrasted, more geometric neck pattern. The contact of the vertebral zigzag band with the dark pileus is shown in fig. 12 of the above publication. Note also striking similarity between the *darevskii* specimen in NILSON & ANDRÉN (2001, fig. 64) and our figure 1.

Pholidotic features of the Zekeriya vipers correspond well also with the published data for *darevskii* (ORLOV & TUNIYEV 1990): 19 or 21 dorsal scales, two apical scales in the majority of specimens, frontal usually larger than parietals, 129-144 (average 137) ventrals, nostril usually piercing the lower part of the nasal. The main difference concerns the contact between the upper pre-ocular and the nasal, present in the majority of *darevskii* from Mount Legli but absent in the eight Zekeriya specimens.

DISCUSSION AND CONCLUSIONS

In summary, the morphological features of the Zekeriya vipers separate them from all known viper taxa except *darevskii* while their features correspond well with what is known about *darevskii*. The known records of *darevskii* lie only 130 km from Zekeriya, much closer than those of the other taxa except *eriwanensis*, from which they clearly differ in body proportions and coloration. We therefore conclude that the Zekeriya vipers constitute an additional population of *Vipera darveskii*.

As in Armenia, the Zekeriya population of *darevskii* lies on the northern edge of the distribution of *eriwanensis*. The closest known site of V. eriwanensis in Turkey is north of Kömürlü (vilayet of Erzurum), 25 km SE of Zekeriya (BRUNO 1985, pers. obs.). The habitat at Zekeriya differs from those in all known Turkish localities of eriwanensis, which are on basalt substratum (TEYNIÉ 1987, pers. obs.) by being calcareous, with short, heavily grazed alpine meadows and numerous piles of stones, above the upper altitudinal limit of the forest (fig. 10). This corresponds well to the habitat at Mount Legli as described in ORLOV & TUNIYEV (1990) although the Mount Legli habitats are also on volcanic substratum. The herpetological community observed at the Zekeriya locality was rather poor: Rana

macrocnemis camerani BOULENGER, 1886 (four individuals) and *Anguis fragilis colchicus* (NORDMANN, 1840) (three individuals).

In our opinion, the morphological characters of *darevskii* do not link it with the kaznakovi – dinniki subgroup, in contrast to what has been previously published. Our observations fit well with the recent genetic results (JOGER et al. 2002: KALYABINA-HAUF 2004). To the characteristics discussed above, we can add that our juvenile specimen (apparently recently born) was very small (13.5 cm total length), much smaller than is typical for newborn kaznakovi which are from 160 mm (pers. obs.) to 145 mm (MALLOW et al. 2003). The new characters to identify V. darevskii are: (1) a rather angular canthus rostralis, (2) one series if large oval vertical dark blotches on each flank, (3) the two contrasted light dorsolateral stripes, (4) the clear-cut anterior edge of the vertebral zigzag, (5) the very little size of the newborn and (6) the yellow coloured venom.

Discovery of the Zekeriya population has important consequences for the conservation of *Vipera darevskii*, since the species was previously known from a single locality only. The considerable range extension associated with this new site raises the possibility that the species occurs elsewhere in the mountain ranges between Mount Legli

32

and Zekeriya. Future investigations should be directed at high altitude meadows, especially dry alpine meadows with short, grazed vegetation and bare grounds, probably a habitat probably rarely visited by herpetologists and snake enthusiasts in NE Turkey. Although this discovery provides evidence that the area inhabited by *Vipera darevskii* is larger than previously realized, we wish to stress that collecting for whatever purpose should be avoided given the present uncertainties on the species' prospects.

33

ACKNOWLEDGEMENTS

We want thank the Société d'Histoire Naturelle "Alcide d'Orbigny", Michel BUCHARD (Ets E. Leclerc, Clermont-Ferrand, France) for financial support, Frederic DURAND (Société d'Histoire Naturelle "Alcide d'Orbigny", Clermont-Ferrand, France), Pierre-André CROCHET (CEFE-CNRS, Montpellier, France) and Gilles POTTIER (Nature Midi-Pyrénées, Odos, France) who accompanied us during the second trip.

REFERENCES

BARAN, I. & JOGER, U. & KUTRUP, B. & TÜR-KOZAN, O. (2001): On new specimens of *Vipera barani* BÖHME & JOGER, 1983, from northeastern Anatolia, and implications for the validity of *Vipera pontica* BILLING, NILSSON & SATTLER, 1990 (Reptilia, Viperidae).- Zoology in the Middle East, Heidelberg; 23: 47-53.

BRUNO, S. (1985): Le vipere d'Italia ed'Europa. Bologna (Edagricole), 269 pp.

HÖGGREN, M. & NILSON, G. & ANDRÉN, C. & ORLOV, N. & TUNIYEV, N. L. (1993): Vipers of the Caucasus: natural history and systematic review.- Herpetological Natural History; Phoenix; 1: 11-19.

JOGER, U. & KALYABINA-HAUF, S. A. & SCHWEIGER, S. & MAYER, W. & ORLOV, N. L. & WINK, M. (2002): Phylogeny of Eurasian Vipera (subgenus Pelias). Abstract of Internationale Tagung der DGHT-AG Feldherpetologie und der AG Amphibien und Reptilienschutz in Hessen (AGAR), Freitag 22. bis Sonntag 24. November 2002, Darmstadt; p. 77.

KALYABINA-HAUF, S. A. & SCHWEIGER, S. & JOGER, U. & MAYER, W. & ORLOV, N. L. & WINK, M. (2004): Phylogeny of Eurasian *Vipera* (subgenus *Pelias*); pp. 7-16. In: JOGER, U. & WOLLESEN, R. (eds.): Verbreitung, Ökologie und Schutz der Kreuzotter (*Vipera berus* [LINNAEUS, 1758]), Mertensiella, Rheinbach; 15.

MALLOW, D. & LUDWIG, D. & NILSON, G. (2003): True Vipers. Natural history and toxinology of

Old World Vipers. Malabar (Krieger Publishing Company), 359 pp.

NILSON, G. & ANDRÉN, C. (2001): The meadow and steppe vipers of Europe and Asia. The Vipera (Acridophaga) ursinii complex.- Acta Zoologica Academiae Scientiarum Hungaricae, Budapest; 47 (2-3): 87-267.

NILSON, G. & TUNIYEV, B. S. & ORLOV, N. & HÖGGREN, M. & ANDRÉN, C. (1995): Systematics of the vipers of the Caucasus: polymorphism or sibling species?- Asiatic Herpetological Research, Berkeley; 6: 1-26.

ORLOV, N. L. & TUNIYEV, B. (1990): Three species in the *Vipera kaznakowi* complex (Eurosiberian group) in the Caucasus: their present distribution, possible genesis, and phylogeny. Asiatic Herpetological Research, Berkeley; 3: 1-36.

Research, Berkeley, 3: 1-36. TEYNIÉ, A. (1987): Observations herpétologiques en Turquie. l^{ère} partie.- Bulletin de la Société Herpétologique de France, Paris; 43: 9-18.

TUNIYEV, B. S. & OSTROVSKIKH, S. V. (2001): Two new species of vipers of "*kaznakovi*" complex (Ophidia, Viperinae) from the Western Caucasus. -Russian Journal of Herpetology, Moscou; 8: 117-126.

VEDMEDERJA, V. J. & ORLOV, N. L. & TUNIYEV, B. S. (1986): On taxonomy of three viper species of the *Vipera kaznakovi* complex; pp. 55-61. In: ANANJEVA, N. & BORKIN, L. (eds.): Systematics and ecology of amphibians and reptiles.- (USSR Acad. Sci.) Proc. Zool. Inst., Leningrad; 157. [in Russian].

DATE OF SUBMISSION: April 19, 2004

Corresponding editor: Heinz Grillitsch

AUTHORS: Philippe GENIEZ, Ecole Pratique des Hautes Etudes, Laboratoire de Biogéographie et Ecologie des Vertébrés, cc.94, UM2, F-34095 Montpellier-cedex 5, France < geniez@univ-montp2.fr >; Alexandre TEYNIÉ, Société d'Histoire Naturelle "Alcide d'Orbigny", Route de Verneuge, 63970, Aydat, France. < ateynie@clermont. inra.fr >

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Herpetozoa

Jahr/Year: 2005

Band/Volume: <u>18_1_2</u>

Autor(en)/Author(s): Geniez Philippe, Teynié Alenxandre

Artikel/Article: Discovery of a population of the critically endangered Vipera darevskii VEDMEDERJA, ORLOV & TUNIYEV, 1986 in Turkey, with new elements on ist identification 25-33