

and *T. v. graecus*, must be determined and further examined to explain the apparent absence of *T. v. vulgaris*.

ACKNOWLEDGEMENTS: Johann TÖPFER (Wien) and Michael DUDA (Perchtoldsdorf) were the first to indicate the presence of conspicuous Smooth Newts in the vicinity of Kaltenleutgeben and Perchtoldsdorf. Our sincere thanks to them. We are also grateful to Günter MERKL for his information about the history of the quarry water bodies on the Parapluiberg and Großer Flössberg mountains. For almost 40 years until 2000, he was the manager of the cement plant, using limestone from the quarries of Kaltenleutgeben; we owe it to his vision that these water bodies were created in these inactive quarries. Thanks also to Marcia K. NELSON (Fulton, Missouri, USA) for translating the German original into English. The office of NÖ Landesregierung, Gruppe Raumordnung, Umwelt und Verkehr, Abteilung Naturschutz issued the special collecting permits RU5-SB-033/009-2001, RU5-SB-033/012-2002, RU5-SB-033/013-2003, RU5-BE-64/001-2004.

REFERENCES: ARNTZEN, J. W. (2003): *Triturus cristatus* Superspecies – Kammolch-Artenkreis; pp. 421-514. In: GROSSENBACHER, K. & THIESMEIER, B. (ed.): Handbuch der Reptilien und Amphibien Europas; Vol. 4/II A Schwanzlurche (Urodela) IIA, Salamandridae II; *Triturus* I; Wiebelsheim (Aula Verlag). BLAB, J. & BRÜGEMANN, P. & SAUER, H. (1991): Tierwelt in der Zivilisationslandschaft; Teil II Raumeinbindung und Biotopnutzung bei Reptilien und Amphibien im Drachenfels-Ländchen; Bonn-Bad Godesberg (Kilda), pp. 94. CABELA, A. & GRILLITSCH, H. & TIEDEMANN, F. (2001): Atlas zur Verbreitung und Ökologie der Amphibien und Reptilien in Österreich: Auswertung der Herpetofaunistischen Datenbank der Herpetologischen Sammlung des Naturhistorischen Museums in Wien; Wien (Umweltbundesamt, Hrsg.), pp. 880. RAXWORTHY, C. J. (1990): A review of the smooth newt (*Triturus vulgaris*) subspecies, including an identification key. - Herpetological Journal, London; 1 (11): 481-492. SCHÄFER, H.-J. & KNEITZ, G. (1993): Entwicklung und Ausbreitung von Amphibienpopulationen in der Agrarlandschaft – ein E + E-Vorhaben. Natur und Landschaft, Bonn-Bad Godesberg; 68 (7/8): 376-385. SCHIMDTLER, J. J. & SCHIMDTLER, J. F. (1983): Verbreitung, Ökologie und innerartliche Gliederung von *Triturus vulgaris* in den adriatischen Küstengebieten. - Spixiana, München; 6 (3): 229-249. SCHIMDTLER, J. F. & FRANZEN, M. (2004): *Triturus vulgaris* (LINNAEUS, 1758) – Teichmolch; pp. 847-967. In: THIESMEIER, B. & GROSSENBACHER, K. (ed.): Handbuch der Reptilien und Amphibien Europas; Vol. 4/II B Schwanzlurche (Urodela), Salamandridae III; Wiebelsheim (Aula Verlag).

SUBMITTED: December 28, 2004

AUTHORS: Antonia CABELA, Heinz GRILLITSCH, Herpetological Collection, First Zoological Department, Natural History Museum Vienna, Burgring 7, A-1010 Vienna, Austria; Günter SCHULTSCHIK, Sachsenweg 6, Haus 12, A-2391 Kaltenleutgeben, Austria; Franz TIEDEMANN, Herpetological Collection, First Zoological Department, Natural History Museum Vienna, Burgring 7, A-1010 Vienna, Austria. <franz.tiedemann@nhm-wien.ac.at>

Triturus italicus (PERACCA, 1898) in the Lepini Mountains: new north-western range limit

Triturus italicus (PERACCA, 1898) is a newt endemic to southern Italy. The northern limits of its distribution are located in the province of Ancona (region of Marche) and in the Ausoni Mountains (region of Lazio) at the Adriatic and the Tyrrhenian side of the Apennines, respectively (BOLOGNA 2000) (fig. 1). The first sighting of *T. italicus* for Lazio was reported in 1981 (cf. BONIFAZI & CARPANETO 1990). To date about 50 breeding sites have been recorded for southern Lazio (CORSETTI, in press) while its presence has never been recorded for the province of Rome (RM) (cf. BOLOGNA 2000; CORSETTI 1994, 2000; CORSETTI & CAPULA 1992). In this note we report the first data on the presence of *T. italicus* in an area north of the Ausoni Mts.

Between 2003 and 2004, we found the newt in five localities between 700 and 1,020 m a.s.l. in the Lepini Mts.: One in the district of Castro dei Volsci (province of Frosinone), three in the district of Segni (RM) and one in the district of Montelanico (RM). We report here also on a record locality in the district of Carpineto Romano (RM) where the newt was found in spring 2004 (MARCO A. BOLOGNA, pers. comm.). In some of these localities more than one breeding site was present. In total we found eleven breeding sites where we observed adult males and females, as well as neotenic individuals. Our observations move the known north-western limit of the range of the Italian Newt about 35 km in a northward direction along the Tyrrhenian side of Apennines (fig. 1). Moreover, the site at 1,020 m a.s.l. in the Lepini Mts. represents the new altitudinal record for this species in the region of Latium where it was not known from above 900 m a.s.l. before (BOLOGNA 2000). Among the reported breeding sites, ten were stony wells and only one was a natural pond; they were all located in pastures, with or without shrubs.

T. italicus is protected by regional law number 18 of 1988 and the newt's conservation is usually implemented by preserving




Fig. 1: The new stations of *Triturus italicus* (PERACCA, 1898) in the Lepini Mts. move the known north-western limit of the range of the Italian Newt about 35 km in a northward direction along the Tyrrhenian side of the Apennines (black dashed line). Gray dashed line - limit of distribution of the Italian Newt as previously known; ■ - locality of *T. italicus* in the province of Frosinone; ★ - localities in the province of Rome; FR - Frosinone; LT - Latina.

natural breeding sites. However, 60 % of *T. italicus* breeding sites in southern Lazio are artificial water bodies such as fountains, livestock watering ponds, stony wells and tanks (CORSETTI 2000, in press). During the last decades, the functionality and integrity of these artificial structures has been more and more compromised (i.e. collapses, filling processes, water drying up) by changes in the traditional methods of local livestock husbandry (CORSETTI in press). For this reason conservation problems should be expected to arise for the newt populations living in this area. Our data confirm the dependence, at a local level, of *T. italicus* populations from the integrity of artificial water bodies. Accordingly, we suggest the promotion of traditional management of the artificial water bodies used in local livestock husbandry as a more effective strategy for the conservation of *T. italicus* populations living in Southern Lazio.

ACKNOWLEDGEMENTS: We are grateful to Marco A. BOLOGNA, Gianpaolo MONTINARO, Monica PITZALIS and Leonardo VIGNOLI for the informations concerning the breeding site in Carpineto Romano; Renzo DE ANGELIS, Antonio RUGGIERO and Tiziana TAMBURU for their useful contribution to the draft of this note.

REFERENCES: BOLOGNA, M. A. (2000): *Triturus italicus*; pp. 44-45. In: BOLOGNA, M. A. & CAPULA, M. & CARPANETO, G. M. (eds.): Anfibi e rettili del Lazio; Roma (Fratelli Palombi). BONIFAZI, A. & CARPANETO, G. M. (1990): Indagine preliminare sugli anfibi e sui rettili dei Monti Ausoni-Aurunci (Lazio Meridionale).- Centro reg. doc. beni cult. amb. Lazio, Assessore Cultura Reg. Lazio, Roma, 47 pp. CORSETTI, L. (1994): Anfibi e rettili dei Monti Lepini.- Quaderni del Museo di Storia Naturale di Pätrica, Pätrica (FR), 5, 190 pp. CORSETTI, L. (2000): Caratteristiche ambientali dei siti riproduttivi di *Triturus italicus* (PERACCA, 1898) nel Lazio (Italia centrale).- Rivista di Idrobiologia, Perugia; 38 (1999) 1/2/3: 449-455. CORSETTI, L. (in press): Distribuzione e preferenze ambientali degli anfibi urodeli nel Lazio meridionale (Italia centrale).- 5th Congresso Nazionale Societas Herpetologica Italica, Calci (PI), 29 settembre - 3 ottobre 2004, Riassunti. CORSETTI, L. & CAPULA, M. (1992): The Amphibians and reptiles of the Lepini Mts. (Latium, Central Italy): Checklist and provisional atlas.- British Herpetological Society Bulletin, London; 39: 8-16.

KEY WORDS: *Triturus italicus*, limit of the range, new records, regional distribution, Lepini Mts., Italy

SUBMITTED: June 28, 2004

AUTHORS: Luigi CORSETTI, Via Adige 45, 04100 Latina, Italy < info@edizionibelvedere.it >; Roberto RAGNO, Via Triboniano 16, 04100 Latina, Italy; Antonio ROMANO, Dipartimento di Biologia, Università degli Studi di Roma "Tor Vergata", Via della Ricerca Scientifica 00133 Roma, Italy < antonioromanol@libero.it >.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Herpetozoa](#)

Jahr/Year: 2005

Band/Volume: [18_1_2](#)

Autor(en)/Author(s): Romano Antonio, Corsetti Luigi, Ragno Roberto

Artikel/Article: [Triturus italicus \(PERACCA, 1898\) in the Lepini Mountains:
new north-western range limit 87-88](#)