

REFERENCES: PÉREZ-SANTOS, C. & MORENO, A. G. (1991): Serpientes de Ecuador.- Monografía di Museo Regionale di Scienze Naturali, Torino; 11: 1–538. SAVAGE, J. M. (2002): The amphibians and reptiles of Costa Rica, a herpetofauna between two continents, between two seas; Chicago (The University of Chicago Press), pp. 934.

KEY WORDS: Reptilia: Serpentes: Colubridae: *Lampropeltis triangulum micropholis*; geographic range, altitudinal distribution, new country records, conservation status, Ecuador

SUBMITTED: February 22, 2006

AUTHOR: Diego F. CISNEROS-HEREDIA, College of Biological and Environmental Sciences, Universidad San Francisco de Quito, Casilla Postal 17-12-841, Quito, Ecuador < diegofrancisco_cisneros@yahoo.com >

Notes on some *Osteocephalus* treefrogs from Amazonian Ecuador

Forty-nine species of treefrogs (family Hylidae) have been recorded at the Tiputini Biodiversity Station (TBS), Orellana province, Amazonian Ecuador (0°37'5"S, 76°10'19"W, 190–270 m a.s.l.) (CISNEROS-HEREDIA 2001, 2003); including eight species of the genus *Osteocephalus*, namely *O. buckleyi* (BOULENGER, 1882), *O. cabrerai* (COCHRAN & GOIN, 1970), *O. planiceps* COPE, 1874, *O. taurinus* STEINDACHNER, 1862, *O. deridens* JUNGFER, RON, SEIPP & ALMENDÁRIZ, 2000, *O. fuscifacies* JUNGFER, RON, SEIPP & ALMENDÁRIZ, 2000, *O. mutabor* JUNGFER & HÖDL, 2002, and *O. yasuni* RON & PRAMUK, 1999 (CISNEROS-HEREDIA 2003). Several *Osteocephalus* species have been recently described and our knowledge on them is still at a basic level. Herein, I present some notes on the distribution and natural history of two poorly-known *Osteocephalus*.

Two recently described species, *O. yasuni* and *O. mutabor*, were collected at the Tiputini Biodiversity Station, being their easternmost location known from Ecuador. These records extend the ranges of *O. yasuni* ca. 31 km E from the type locality and of *O. mutabor* ca. 54 km SE from the nearest known record (San Pablo de Kantesiaya, JUNGFER & HÖDL 2002). *Osteocephalus mutabor* was also collected at the Reserva de Producción Faunística Cuyabeno (76°12'

54"W, 00°05'02"S, 290 m a.s.l., 20 July 2000), in terra firme forest, extending its range ca. 30 km E from the nearest known locality in northern Napo River bank (San Pablo de Kantesiaya, JUNGFER & HÖDL 2002).

Adults of *Osteocephalus yasuni* were reported in the original description as having white bones, brown webbing and yellow venter with the color intensifying towards the groin (RON & PRAMUK 1999); however juveniles were found to have green bones (ontogenetic change also reported for *Osteocephalus lepreurii*, JUNGFER & HÖDL 2002) and intense yellow-orange color on webbing and on the entire venter and throat. *Osteocephalus yasuni* was found at primary terra firme (mainly females and non-reproductive males) and flooded forests near stream edges (mainly reproductive males) in TBS.

ACKNOWLEDGMENTS: Thanks to S. CÁRDENAS, M. BRANDT, C. PONCE and A. LEÓN for field support; to K.-H. JUNGFER, J. SIMMONS, R. W. MCDIARMID, and S. RON for sharing literature and information; to L. COLOMA and K.-H. JUNGFER for confirming the identifications; to Tiputini Biodiversity Station, and Maria Elena HEREDIA and Laura HEREDIA who provided financial and moral support.

REFERENCES: CISNEROS-HEREDIA, D. F. (2001): Anfíbios y reptiles de la Estación de Biodiversidad Tiputini (EBT), Ecuador. Ver. 2.1 (August 2001). Laboratorio de Anfíbios y Reptiles, Universidad San Francisco de Quito. < <http://www.cisneros-heredia.org> > [last accessed: 24 February 2004]. CISNEROS-HEREDIA, D. F. (2003): Herpetofauna de la Estación de Biodiversidad Tiputini; pp. 1–21. In: DE LA TORRE, S. & RECK, G. (eds.): Ecología y ambiente en el Ecuador. Mem. I. Congr. Ecología y Ambiente. CD. Universidad San Francisco de Quito, Ecuador. JUNGFER, K.-H. & HÖDL, W. (2002): A new species of *Osteocephalus* from Ecuador and a redescription of *O. lepreurii* (DUMÉRIL & BIBRON, 1841) (Anura: Hylidae).- Amphibia-Reptilia, Leiden; 23: 21–46. RON, S. & PRAMUK, J. B. (1999): A new species of *Osteocephalus* (Anura: Hylidae) from Amazonian Ecuador and Peru.- Herpetologica, Emporia; 55 (4): 433–446.

KEY WORDS: Amphibia: Anura: Hylidae: *Osteocephalus yasuni*, *Osteocephalus mutabor*, distribution, natural history, ontogenetic chromatic change, new records, Ecuador

SUBMITTED: December 06, 2005

AUTHOR: Diego F. CISNEROS-HEREDIA, College of Biological and Environmental Sciences, Universidad San Francisco de Quito, Ave. Interocéánica y calle Diego de Robles, Campus Cumbayá, Edif. Maxwell. Casilla Postal 17-12-841 Quito, Ecuador < diegofrancisco_cisneros@yahoo.com >

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Herpetozoa](#)

Jahr/Year: 2007

Band/Volume: [19_3_4](#)

Autor(en)/Author(s): Cisneros-Heredia Diego F.

Artikel/Article: [Notes on some Osteocephalus treefrogs from Amazonian Ecuador 183](#)