

**ACKNOWLEDGMENTS:** We appreciate the assistance provided by Steve GOTTE, Ron HEYER, Roy McDIARMID, and George ZUG while studying *Phyllonastes* at the National Museum of Natural History. Special thanks are extended to Roy McDIARMID, Michael HARVEY, Edgar LEHR, Karl-Heinz JUNGFER, John LYNCH, and Ana ALMENDÁRIZ for sharing their thoughts on *Phyllonastes* or providing access to their research specimens, and to Adrian FORSYTH, Susana CÁRDENAS, Andrés LEÓN, Javier ROBAYO and Vlastimil ZAK for field companionship and support. Research by R. P. REYNOLDS was supported by Conservation International's Rapid Assessment Program. Research by D. F. CISNEROS-HEREDIA was supported by the 2002 Research Training Program, National Museum of Natural History, Smithsonian Institution, the Smithsonian Women's Committee, Universidad San Francisco de Quito, Vlastimil ZAK (Bosque Protector Río Guajalito), and María Elena HÉREDA and Laura HERÉDIA.

**REFERENCES:** ALMENDARIZ, A. (1997): Overview of the herpetofauna of the western slopes of the Cordillera del Cóndor; pp. 80–82, 199–201. In: SCHULENBERG, T. S. & AWBREY, K. (Eds.): The Cordillera del Cóndor region of Ecuador and Peru: A biological assessment. Rapid Assessment Working Paper No. 7. Washington, D.C. (Conservation International). ALONSO, A. & DALLMEIER, F. (Eds) (1999): Biodiversity assessment and monitoring of the lower Urubamba region, Peru: Pagoreni well site assessment and training.- SI/MAP Series No. 3. Washington, D.C. (Smithsonian Institution / Man and Biosphere Biodiversity Program). COLOMA, L. (2005–2006): Anfibios de Ecuador. Ver. 2.0., web document accessible at <<http://www.puce.edu.ec/zoologia/vertebrados/amphibia/webec/index.html>> Quito (Museo de Zoología, Pontificia Universidad Católica del Ecuador). DOAN, T. M. & ARIZABAL, W. (2002): Microgeographic variation in species composition of the herpetofaunal communities of Tambopata region, Peru.- Biotropica, Connecticut; 34: 101–117. DUELLMAN, W. E. & LYNCH, J. D. (1988): Anuran amphibians from the Cordillera de Cutucú, Ecuador.- Proc. Acad. Nat. Sci. Philadelphia, Philadelphia; 140 (2): 125–142. FROST, D. R. (2004): Amphibian species of the world: an online reference. Version 3.0 (22 August, 2004) <<http://research.amnh.org/herpetology/amphibia/index.html>> New York (American Museum of Natural History). HEYER, W. R. (1977): Taxonomic notes on frogs from the Madeira and Purus rivers, Brazil.- Papéis Avulsos de Zoologia, São Paulo; 31: 141–162. HOOGMOED, M. S. & LESCURE, J. (1984): A new genus and two new species of minute leptodactylid frogs from Northern South America, with comments upon *Phyzelaphryne* (Amphibia: Anura: Leptodactylidae).- Zool. Mededel., Leiden; 58: 85–115. LEHR, E. & AGUILAR, C. & LUNDBERG, M. (2004): A new species of *Phyllonastes* from Peru (Amphibia, Anura, Leptodactylidae).- Journal of Herpetology, Salt Lake City; 38 (2): 214–218. LYNCH, J. D. (1976): Two new species of frogs of the genus *Euparkerella* (Amphibia: Leptodactylidae) from Ecuador and Perú.- Herpetologica, Emporia; 32: 48–53. LYNCH, J. D. (1986): New species of minute leptodactylid frogs from the Andes of Ecuador and Peru.- Journal of Herpetology, Salt Lake City; 20 (3): 423–431. LYNCH, J. D. (2005): Discovery of the richest frog fauna in the world – An exploration of the forests to the North of Leticia.- Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales, Bogota; 113: 581–588. LYNCH, J. D. & DUELLMAN, W. E. (1997): Frogs of the

genus *Eleutherodactylus* (Leptodactylidae) in western Ecuador: systematics, ecology, and biogeography.- Nat. Hist. Mus., Univ. Kansas, Spec. Publ., Lawrence; (23): 1–236. REICHLE, S. & AGUAYO, R. & CORTEZ, C. (2004): Geographic distribution: *Phyllonastes myrmecoides*.- Herpetological Review, New Haven; 35 (3): 283. REYNOLDS, R. P. & ICOCHEA, M. J. (1997): Amphibians and reptiles of the upper Río Comainas, Cordillera del Cóndor; pp. 82–86, 202–206. In: SCHULENBERG, T. S. & AWBREY, K. (Eds.): The Cordillera del Cóndor region of Ecuador and Peru: A biological assessment. Rapid Assessment Working Paper No. 7, Washington, D. C. (Conservation International). RODRÍGUEZ, L. O. & CADLE, J. E. (1990): A preliminary overview of the herpetofauna of Cocha Cashu, Manu National Park, Peru, pp. 410–425. In: GENTRY, A. H. (Ed.): Four Neotropical rainforests. New Haven (Yale University Press), pp. xiii + 627. RODRÍGUEZ, L. O. & CORDOVA, J. H. & ICOCHEA, J. (1993): Lista preliminar de los anfibios del Perú.- Pub. Mus. His. Nat. Univ. Mayor de San Marcos, Lima; 45: 1–22. SIERRA, R. (Ed.) (1999): Propuesta preliminar de un sistema de clasificación de la vegetación para el Ecuador continental. Quito (Proyecto INEFAN/GEF-BIRF y EcoCiencia), pp. 194.

**KEY WORDS:** Amphibia: Anura: Brachycephalidae: *Phyllonastes myrmecoides*, *Phyllonastes lochites*, new country records, geographic range, distribution, Ecuador, Peru

SUBMITTED: April 23, 2006

AUTHORS: Diego F. CISNEROS-HEREDIA, College of Biological and Environmental Sciences, Universidad San Francisco de Quito, Casilla 17-12-841, Quito, Ecuador, <[diegofrancisco\\_cisneros@yahoo.com](mailto:diegofrancisco_cisneros@yahoo.com)>; Robert P. REYNOLDS, US Geological Survey, Patuxent Wildlife Research Center, National Museum of Natural History, MRC 111, P.O. Box 37012 Washington, DC 20013-7012, USA, <[rpreynolds@usgs.gov](mailto:rpreynolds@usgs.gov)>

## First record of the coral snake *Micrurus margaritiferus* ROZE, 1967 in Ecuador

*Micrurus margaritiferus* ROZE, 1967 is known from lower montane wet forests and tropical rainforests on the Amazonian slopes of the Peruvian Andes near the confluence of the rivers Santiago and Marañón (CAMPBELL & LAMAR 2004, 1989; CARILLO DE ESPINOZA & ICOCHEA 1995). Until recently, speculations on the presence of this species in Ecuador (CAMPBELL & LAMAR 2004) remained unsubstantiated.

The first record of *M. margaritiferus* in the territory of Ecuador (Gustavo ORCÉS Herpetological Foundation reference collection – FHGO 4836) was made on 6 July, 2005, near the banks of Río Makuma (02° 08'6"S/77°42'25"W, 720 m a.s.l.) in the Shuar-Makuma community located in the Morona Santiago Province, northeastern



Fig. 1: *Micrurus margaritiferus* ROZE, 1967 (FHGO 4836), dorsal aspect. First record from Ecuador.

portion of the Cordillera de Cutucú (see BORGTOFT et al. 1998).

**Morphological data:** Total length 75.5 cm (snout-vent-length 63.3 cm, tail length 12.2 cm); 1+1 temporals, 201 ventrals, 52 subcaudals. Dorsally the specimen is tricolored (black-white-brown/reddish-white-black, Fig. 1), which seems to be rare in this species (CAMPBELL & LAMAR 2004). The white rings are irregularly and frequently interrupted in the different parts of the body. The dark (black plus brown/reddish) pattern

is disposed in 111 rings across the trunk (each ring being on the average three scales wide middorsally), and 13 on the tail. The brown/reddish color is clearly visible in the anterior and becomes black in the posterior part of the body. The venter is tricolored with black, brown/reddish and white rings (Fig. 2). The head is black with irregular white dots on the posterior supra- and infra-labials, occipitals and temporals. Mentals, some supralabials and first and second ventrals are black.

The snake was found during the day, in a grassland area next to the evangelic mission. The habitat was relatively flat secondary forest, in an advanced stage of succession at the confluence of various small streams. At this site, the forest is not continuous due to the presence of clearing gaps and cultivated areas (VALENCIA et al. 2005).

Fieldwork has demonstrated that this area is a region of great importance for Amazonian Ecuador in terms of reptile diversity (VALENCIA et al. 2005).

**ACKNOWLEDGMENTS:** I would like to thank Dwain HOLMES and Lloyd Holmes (Centro Shuar Makuma), Walter SCHARGEL (University of Texas at Arlington), and Manuel MORALES (Amphibians and Reptiles Laboratory FHGO) for their help and María Elena BARRAGÁN and Katy GARZÓN (FHGO - Gustavo ORCÉS Herpetological Foundation) for comments and correcting the English manuscript.

**REFERENCES:** BORGTOFT, H. & SKOV, F. & FJELDSÅ, J. & SCHJELLERUP, I. & ØLLGAARD, B. (eds.) (1998): People and biodiversity. Two cases studies from the Andean foothills of Ecuador. Kalø, Denmark. (National Environmental Research Institute; The Danish Environmental Research Programme); Centre for Research on Cultural and Biological Diversity of Andean Rainforests (DIVA) Technical Report no. 3; pp. 190. CAMPBELL, J. A. & LAMAR, W. W. (1989): The venomous reptiles of Latin America. Ithaca, New York (Cornell University Press), pp. 425. CAMPBELL, J. A. & LAMAR, W. W. (2004): The venomous reptiles of the Western Hemisphere. Vols. 1 & 2. Ithaca, New York (Cornell University Press), pp. xviii + 898. CARRILLO DE ESPINOZA, N. & ICOCHEA, J. (1995): Lista taxonómica preliminar de los reptiles vivientes del Perú.- Publicaciones del Museo de Historia Natural / Universidad Nacional Mayor de San Marcos (UNMSM), Lima; (Ser. A) 49: 1-27. VALENCIA, J. H. & MORALES-MITE, M. & HOLMES, D. (2005): Anfibios y reptiles de las provincias de Pastaza y Morona Santiago, Amazonía del Ecuador. Memorias del Simposio Nacional Avances en la Herpetología Neotropical. Instituto Clodomiro Picado. San José Costa Rica, 9 November, 2005, (CD-ROM).

**KEY WORDS:** Reptilia; Squamata: Elapidae, *Micrurus margaritiferus*, distribution, new record, Morona Santiago, Ecuador

SUBMITTED: March 17, 2006

AUTHOR: Jorge H. VALENCIA, Fundación Herpetologica Gustavo Orcés, Av. Amazonas 3008 y Rumipamba, Quito, Ecuador <jvalencia@fhgo.org.ec>



Fig. 2: *Micrurus margaritiferus* ROZE, 1967 (FHGO 4836), ventral aspect. First record from Ecuador.

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Herpetozoa](#)

Jahr/Year: 2007

Band/Volume: [19\\_3\\_4](#)

Autor(en)/Author(s): Valencia Jorge H.

Artikel/Article: [First record of the coral snake Micrurus margaritiferus Roze, 1967 in Ecuador 186-187](#)