

the Conservation of Nature for the logistic support. The work of the last author (JM) was supported by the Grant Agency of the Czech Republic, project no. 206/05/ 2334, and by the Ministry of Culture of the Czech Republic, projects nos. MK00002327201 and DE06P04OMG008.

REFERENCES: ABED, A. M. (2002): An overview of geology of Jordan; pp. 21-24. In: DISI, A. (Ed.): Jordan country study on biological diversity: The herpetofauna of Jordan; United Nations Environment Programme, Amman. ABU BAKER, M. & QARQAZ, M. & RIFAI, L. & HAMIDAN, N. & AL OMARI, K. & MODRÝ, D. & AMR, Z. (2004): Results of herpetofaunal inventory of Wadi Ramm Protected Area, with notes on some relict species.- Russian Journal of Herpetology, Moskva; 11 (1): 1-5. BAHA EL DIN, S. (2005): An overview of Egyptian species of *Hemidactylus* (Gekkonidae), with the description of a new species from the high mountains of South Sinai.- Zoology in the Middle East, Heidelberg; 34: 11-26. CARRANZA, S. & ARNOLD, E. N. (2006): Systematics, biogeography, and evolution of *Hemidactylus* geckos (Reptilia: Gekkonidae) elucidated using mitochondrial DNA sequences.- Molecular Phylogenetics and Evolution, San Diego; 38: 531-545. DISI, A. M. (1996): A contribution to the knowledge of the herpetofauna of Jordan. VI. The Jordanian herpetofauna as a zoogeographic indicator.- Herpetozoa, Wien; 9 (1/2): 71-81. DISI, A. M. & AMR, Z. S. (1998): Distribution and ecology of lizards in Jordan (Reptilia: Sauria); pp. 43-66. In: FRITZ, U. & OBST, F. J. & ANDREAS, B. (Eds.): Contribution to a "Herpetologia arabica", Faunistische Abhandlungen Staatliches Museum für Tierkunde Dresden, 21, Suppl., 182 pp. DISI, A. M. & AMR, Z. S. & MARTENS, H. (2004): On a collection of amphibians and reptiles made by J. KLAPPERICH in Jordan.- Herpetozoa, Wien; 16 (3/4): 141-150. DISI, A. M. & MODRÝ, D. & NEČAS, P. & RIFAI L. (2001): Amphibians and reptiles of the Hashemite Kingdom of Jordan. An atlas and field guide. Frankfurt a. M. (Chimaira), 408 pp. DISI, A. M. & MODRÝ, D. & BUNIAN, F. & AL-ORAN, R. M. & AMR, Z. S. (1999): Amphibians and reptiles of the Badia region of Jordan.- Herpetozoa, Wien; 12 (3/4): 135-146. MORAVEC, J. & BÖHME, W. (1997): A new subspecies of the Mediterranean gecko *Hemidactylus turcicus* from the Syrian lava desert (Squamata: Sauria: Gekkonidae).- Herpetozoa, Wien; 10 (3/4): 121-128.

KEY WORDS: Reptilia: Squamata: Sauria: Gekkonidae, *Hemidactylus mindiae*, *Hemidactylus turcicus*, new country record, Jordan

SUBMITTED: September 08, 2006

AUTHORS: Zuhair S. AMR, Department of Biology, Faculty of Sciences, Jordan University for Science and Technology, P.O. Box 3030, 22110 Irbid, Jordan < amrz@just.edu.jo >; David MODRÝ, Department of Parasitology, University of Veterinary and Pharmaceutical Sciences, 612 42 Brno, Czech Republic; Mohammad ABU BAKER, Dept. Biological Sciences (MC066), 845 West Taylor, St. University of Illinois at Chicago, Chicago, IL 60607 USA; Mayas QARQAS, The Royal Society for the Conservation of Nature, Amman, Jordan; Jamaal AL Zaidanyen, Ramm Protected Nature Reserve, Aqaba Special Economic Zone Authority; Jiří MORAVEC, Department of Zoology, National Museum, 115 79 Prague 1, Czech Republic < jiri.moravec@nm.cz >.

Cochranella punctulata RUIZ-CARRANZA & LYNCH, 1995 – new record from the department of Antioquia (Colombia), with comments on the conservation status

During a herpetological survey carried out in the Magdalena River valley in October 2005, the authors found an individual of the poorly-known glassfrog *Cochranella punctulata* RUIZ-CARRANZA & LYNCH, 1995; Fig. 1) at Hacienda Santa Bárbara (ca. 06° 32'N / 74°38'W, between 515-575 m above sea level; 1 in Fig. 2), vereda Las Brisas, Maceo municipality, Department of Antioquia. This location is ca 70 km northeast of the nearest known record of this species (vereda La Granja, Cocorná municipality, Department of Antioquia, 06°3'N / 75°11' W, 900-930 m a.s.l.; RUIZ-CARRANZA & LYNCH 1995; 2 in Fig. 2). The female found (Museo de Herpetología Universidad de Antioquia, MHUA 4071; coll. P. D. GUTIÉRREZ-C.) was observed at night (2042 h) on herbaceous vegetation attached to rock boulders near to a small waterfall inside a secondary forest.

This glassfrog species has been listed as vulnerable (IUCN category VU B1ab(iii)), based on its small range area, fragmented distribution and the decline of its forest habitat quality (GIL & LYNCH 2004). The first two arguments are valid considering that until now there are documented records only from three disjunctive localities as indicated in the original description (departments of Antioquia, Caldas and Tolima – RUIZ-CARRANZA & LYNCH 1995; 2-4 in Fig. 2). However, we consider that the previous arguments are only an artifact of lack of surveys, since our new record suggests that there may be other unknown northern populations along the eastern flank of the Cordillera Central, to which *C. punctulata* is apparently restricted (ACOSTA-GALVIS et al. 2006; FROST 2006). This new locality indicates that this species has a more extended distribution range. On the other hand, the adverse effect of declining habitat quality on the frog's distribution range reported by Gil & Lynch (2004) is well substantiated since in some localities of *C. punctulata* such as Samaná (Department of Caldas) and Cocorná



Figure 1: Female *Cochranella punctulata* RUIZ-CARRANZA & LYNCH, 1995 (MHUA 4071) from Vereda Las Brisas, municipio Maceo, department of Antioquia, Colombia. Photo: M. RIVERA-CORREA.




Fig. 2: Localities (municipalities and departments) reported for *Cochranella punctulata* RUIZ-CARRANZA & LYNCH, 1995. 1 - Maceo (Antioquia), the new record; 2 - Cocorná (Antioquia); 3 - Samaná (Caldas); 4 - Falán (Tolima). Localities 2-4 according to RUIZ-CARRANZA & LYNCH (1995).

(Antioquia) there is an impoverishment of the habitat caused mainly by timber logging, dam constructions (i.e., La Miel hydroelectric project in Samaná) and extensive cattle grazing.

With this note, we want to demonstrate that the discovery of new localities for this and other species considered threatened by knowing only their type localities, is essential to differentiate a declining population from a species that is ecologically rare.

ACKNOWLEDGMENTS: We thank to John LYNCH (Instituto de Ciencias Naturales, Universidad Nacional de Colombia, Bogotá) and Juan Manuel GUAYASAMIN (Department of Ecology and Evolutionary Biology, The University of Kansas, Lawrence, KS) for the revision of the specimens. Maritza CUARTAS (Unidad Municipal de Asistencia Técnica Agropecuaria – UMATA of Municipio Maceo) provided permits and lodging facilities at study site. We also thank to students of herpetology class 2005-1 from Universidad de Antioquia for field assistance. Juan M. DAZA (Department of Biology, University of Central Florida, Orlando, FL) gave useful suggestions on an earlier draft of the manuscript. Universidad de Antioquia at Medellín provided funding for fieldwork.

REFERENCES: ACOSTA-GALVIS, A. R. & HUERTAS-SALGADO, C. & RADA, M. (2006): Aproximación al conocimiento de los anfibios en una localidad del Magdalena Medio (Departamento de Caldas, Colombia). - Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales, Bogotá; 30: 291-303. FROST, D. R. (2006). Amphibian species of the world: an on-line reference. Version 4.0 [17 August 2006]. New York. < <http://research.amnh.org/herpetology/amphibia/index.php> > (last accessed: 28 August 2006). GIL, A. Q. & LYNCH, J. (2004): *Cochranella punctulata*. In: IUCN 2006 Red list of threatened species. < <http://www.iucnredlist.org> > (last accessed: 23 August 2006). RUIZ-CARRANZA, P. M. & LYNCH, J. D. (1995): Ranas Centroleniidae de Colombia V. Cuatro nuevas especies de *Cochranella* de la Cordillera Central.- Lozania, Acta Zoológica Colombiana, Bogotá; 62: 1-24.

KEY WORDS: Amphibia: Anura: Centroleniidae, *Cochranella punctulata*, geographic distribution, conservation, new record from Magdalena river valley, Colombia

SUBMITTED: September 08, 2006

AUTHORS: Paul D. GUTIÉRREZ-C., Mauricio RIVERA-CORREA, Grupo Herpetológico de Antioquia and Museo de Herpetología Universidad de Antioquia (MHUA), Instituto de Biología, Universidad de Antioquia, oficina 7-106, A. A. 1226, Medellín, Colombia. Current address of PDG: Departamento de Ciencias Biológicas, Universidad de Caldas, Calle 65 # 26-10, A. A. 275. < pdgutierrez2@yahoo.com > < mauriciorivera79@yahoo.com.ar >

Leposoma southi RUTHVEN & GAIGE, 1924, a new record to the fauna of Antioquia (Colombia)

Leposoma southi RUTHVEN & GAIGE, 1924, the Northern Spectacled Lizard (Fig. 1), is a species distributed from Costa Rica to Colombia (ORTLEB & HEATWOLE, 1965; SAVAGE 2002; CASTAÑO-M. et al. 2004). In Colombia, it is fairly widespread in the western Pacific lowlands, having been registered in several municipalities such as Bahía Solano, Istmina and Nuquí (Department of Chocó); Pueblo Rico (Department of Risaralda); Buenaventura (Department of Valle del Cauca); and Guapi (exactly in Isla Gorgona, Department of Cauca) (SÁNCHEZ-C. et al. 1995; CASTAÑO-M. et al. 2004; Fig. 2).

In a study from Quibdó and Lloró municipalities in the Department of Chocó, RENGIFO et al. (2003) reported *Leposoma rugiceps* (COPE, 1869), but we consider that this report rather refers to *L. southi*, since the former species occurs in the Caribbean lowlands, the Magdalena River valley and the Guiana shield (AYALA & CASTRO unpublished document; PETERS & DONOSO-BARROS 1970; SÁNCHEZ-C. et al. 1995; ÁVILA-PIRES 2005) and not in western Colombia (CASTAÑO-M. et al. 2004). However, since RENGIFO et al. (2003) did not report a voucher specimen to corroborate the actual identity of the lizard and its origin, we did not take into account the localities reported by them in this note. CASTAÑO-M. et al. (2004) mentioned also "Department of Antioquia" for the species, but they did neither specify an exact locality nor a museum specimen from this site. Therefore, we consider that *L. southi* has never been previously recorded from both the Department of Antioquia and the Magdalena River valley in Colombia.

On 26-27 October 2005, J. C. ARREDONDO and V. PÁEZ captured two individuals of the lizard *L. southi* at the Hacienda Santa Bárbara (ca. 06°32'N / 74°38'W, between 515-575 m above sea level), vereda Las Brisas, Maceo municipality, Department of Antioquia. Both specimens were collected on litterfall in the morning, inside secondary forest, then deposited at the Museo de Herpetología Universidad de Antioquia

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Herpetozoa](#)

Jahr/Year: 2007

Band/Volume: [20_1_2](#)

Autor(en)/Author(s): Gutierrez-C. Paul D., Rivera-Correa Mauricio

Artikel/Article: [Cochranella punctulata Ruiz-Carranza & Lynch, 1995 - new record from the department of Antioquia \(Colombia\), with comments on the conservation status 75-77](#)