

graphic region as suggested by HERNÁNDEZ-CAMACHO et al. (1992). In the locality reported here, *L. southi* is syntopic with the congeneric species *L. rugiceps*.

**ACKNOWLEDGMENTS:** We thank to the students of the herpetology class 2005-1 from Universidad de Antioquia for field assistance. We also thank to Maritza CUARTAS (Unidad Municipal de Asistencia Técnica Agropecuaria – UMATA of Municipio Maceo) who provided permits and lodging facilities at study site. Universidad de Antioquia at Medellín provided funding for fieldwork.

**REFERENCES:** ÁVILA PIRES, T. C. S. (2005): Reptiles; pp. 25-40. In: HOLLOWELL, T. & REYNOLDS, R. P. (eds.): Checklist of the terrestrial vertebrates of the Guiana Shield. Bulletin of the Biological Society of Washington, no. 13, Washington. AYALA, S. & CASTRO, F. (unpublished): Lagartos de Colombia / Lizards of Colombia. CASTAÑO-M., O. V. & CÁRDENAS-A., G. & HERNÁNDEZ-R., E. J. & CASTRO-H., F. (2004): Reptiles en el Chocó biogeográfico; pp. 599-631. In: RANGEL-CH., J. O. (ed.): Colombia. Diversidad biótica IV. El Chocó biogeográfico y costa pacífica; Bogotá, Colombia (Instituto de Ciencias Naturales, Universidad Nacional de Colombia). HERNÁNDEZ-CAMACHO, J. & HURTADO-GUERRA, A. & ORTIZ QUIJANO, R. & WALSBURGER, T. (1992): Unidades biogeográficas de Colombia; pp. 104-151. In: HALFFTER, G. (ed.): La diversidad biológica de Iberoamérica; Ciudad de México (Acta Zoológica Mexicana, Xalapa; N.S. 9). ORTEB, E. & HEATWOLE, H. (1965): Comments on some Panamanian lizards with a key to the species from Barro Colorado Island, C. Z. and vicinity.- Caribbean Journal of Science, Mayaguez; 5: 141-147. PETERS, J. & DONOSO-BARROS, R. (1970): Catalogue of Neotropical squamata Part II. Lizards and amphisbaenians; Washington (Smithsonian Institutional Bulletin 297), pp. 293. RENGIFO M., J. T. & JIMÉNEZ ORTEGA, A. M. & ASPIRILLA PEREA, J. & RENJIFO, J. M. & ROA GARCÍA, Y. & MORENO MOSQUERA, F. (2003): Distribución vertical y por sustrato de reptiles en un bosque pluvial tropical (bp-T) del Chocó.- Revista Institucional de la Universidad Tecnológica del Chocó D. L. C., Quibdó; 18: 43-49. SÁNCHEZ-C., H. & CASTAÑO-M., O. & CÁRDENAS-A., G. (1995): Diversidad de los reptiles en Colombia; pp. 277-325. In: RANGEL-CH., J. O. (ed.): Colombia. Diversidad biótica I; Bogotá, Colombia (Instituto de Ciencias Naturales, Universidad Nacional de Colombia). SAVAGE, J. M. (2002): The amphibians and reptiles of Costa Rica. A herpetofauna between two continents, between two seas; Chicago (The University of Chicago Press), pp. 934.

**KEY WORDS:** Reptilia: Squamata: Gymnophthalmidae: *Leposoma southi*, biogeographic regions, distribution, inter-Andean valley, Colombia, new record

**SUBMITTED:** September 08, 2006

**AUTHORS** Paul D. GUTIÉRREZ-C., Juan Camilo ARREDONDO-S., Grupo Herpetológico de Antioquia and Museo de Herpetología, Universidad de Antioquia (MHUA), Instituto de Biología, Universidad de Antioquia, oficina 7-121, A.A. 1226, Medellín, Colombia <pdgutierrez2@yahoo.com> <jcas36@gmail.com>. Present address of Paul D. GUTIÉRREZ-C. - Departamento de Ciencias Biológicas, Universidad de Caldas, Calle 65 # 26-10, Manizales, Caldas

## Presence of *Echinantera undulata* (WIED-NEUWIED, 1824) in Ecuador

Until recently, members of the genus *Echinantera* COPE, 1824, were assigned to the genera *Liophis*, *Taeniophallus*, and *Rhadinaea* (PETERS & OREJAS-MIRANDA 1986). Today, the genus *Echinantera* is considered a monophyletic and well defined group that includes six species: *Echinantera amoena* (JAN, 1863), *E. cephalomaculata* DI-BERNARDO, 1994, *E. cephalostriata* DI-BERNARDO, 1996, *E. cyanopleura* (COPE, 1885), *E. melanostigma* (WAGLER, 1824) and *E. undulata* (WIED-NEUWIED, 1824) (DI-BERNARDO 1992; SCHARGEL et al. 2005).

The presence of *Echinantera undulata* in Ecuador, as the only representative of this genus, was uncertain, even though PETERS & OREJAS-MIRANDA (1986) included Ecuador in the group of countries where this species should occur. *Echinantera undulata* was excluded from the lists of reptiles of Ecuador, later to the above publication, due to the lack of evidence from reference specimens in museums (ALMENDARIZ 1991; PEREZ-SANTOS & MORENO 1991; COLOMA et al. 2006). Nevertheless, monitoring in the central-east of this region, confirmed the presence of this species in the Amazon region of Ecuador (VALENCIA et al. 2005) (Fig. 1).

*Echinantera undulata* was found in the lowlands wet forest of the Pastaza province, Amazon region of Ecuador, in two localities: Moretecocha (FHGO 584) (01° 35' S, 77° 24' W, ca. 427 m a.s.l.) and Shell (FHGO 759) (01° 30' S, 78° 03' W, ca. 1042 m a.s.l.), in tropical rain forest, with the distance between the localities being about 75 km (Fig. 2). This nocturnal snake inhabits leaf litter in primary and secondary forest. Time when specimens were collected was between 19:00 to 21:00 hrs.

Body color pattern is similar in both individuals: Dorsum grayish, including lateral ends of ventrals; a conspicuous longitudinal black postocular stripe to the angle of the mouth, an inverted V constituting a black nuchal band, dark dorsal blotches, approximately 19 in number in both sides along the third and fourth longitudinal scale rows of the anterior midbody (not present in



Fig. 1: *Echinantera undulata* (WIED-NEUWIED, 1824) (FHGO 759) from Shell, Pastaza, Ecuador. Dorsal (above) and ventral (above) aspect.

the specimen from Shell); a black longitudinal stripe extending posteriorly along the third third of the body to the tail. Ventral, mental scales and throat spotless beige in color, small brown blotches in the mental region.

The morphology of the specimens will be described in the following order: FHGO 759 / FHGO 584.

Snout-vent length 24.4 cm / 27.8 cm; tail 5.4 cm / 6.6 cm; dorsal longitudinal scale rows 17-17-15 in both specimens, ventrals 134 / 150; subcaudals 39 / 48; temporals 1+2, supralabials 7, infralabials 8, preoculars 2 and postoculars 2 in both specimens each.

In the lowland tropical rain forests in the Pastaza province of Ecuador *E. undulata* is sympatric with *Liophis typhlus* (LINNAEUS, 1758) and *Liophis reginae* (LINNAEUS, 1758) and can be confounded easily with these species. *Liophis typhlus* differs from *E. undulata* in the number of 19-19-15



Fig. 2: The new record localities (▲) of *Echinantera undulata* (WIED-NEUWIED, 1824) in the Province of Pastaza, Amazonian Ecuador.

dorsal scales, variable dorsal color pattern between grayish and greenish, and lateral blotches in chevron shape; *Liophis reginae* presents black ventral blotches alternating with yellowish ventral blotches and the dorsum without any black longitudinal stripe.

The provinces of Pastaza and Morona Santiago as part of the Amazon region show the highest reptile diversity within Ecuador. Makuma a Shuar Center in Morona Santiago for example, houses more than 90% of the snake species on record for the checklist of the Ecuadorian Squamata (VALENCIA 2007 in press).

**ACKNOWLEDGMENTS:** I would like to express my sincere greetings to Dr. James DIXON (University of Texas, A&M), for his kind and generous spending of time in checking some of the specimens, Walter SCHARGEL (University of Texas, Arlington) for sending me information, Katy GARZON, María Elena BARRAGÁN, María José BARRAGÁN, Jean-Marc TOUZET, Raquel BETANCOURT (FHGO - Fundación Herpetológica Gustavo Orcés, Lima), for their support, advice and comments to this manuscript, and for the logistic support and Fernando NOGALES-SORNOZA and Dwain HOLMES for their help with a map of localities. The research was supported by FHGO.

**REFERENCES:** ALMENDARIZ, A. (1991): Lista de vertebrados del Ecuador. Anfibios y Reptiles.- Politécnica, Quito; 16(3): 86-165. COLOMA, L. A. & QUI-

GUANGO, A. & RON, S. (2006): Reptiles del Ecuador: lista de especies y distribución. Crocodylia, Serpentes y Testudines. Version 1.1 25 mayo 2000. Museo de Zoología, Pontificia Universidad Católica del Ecuador, Quito WWW document < <http://www.puce.edu.ec/Zoologia/repecua.htm> > (last accessed 12 June 2006). Di-BERNARDO, M. (1992): Revalidation of the genus *Echinantera* COPE, 1894, and its conceptual amplification (Serpentes, Colubridae).- Comunicações do Museu de Ciências da Pontifícia Universidade Católica do Rio Grande do Sul (PUCRS), sér. zool., Pôrto Alegre; 5 (13): 225-256. PEREZ-SANTOS, C. & MORENO, Y. A. (1991): Serpientes del Ecuador.- Museo Regionale di Scienze Naturali di Torino. Monografie 11; pp. 538. PIETERS, J. A. & OREJAS-MIRANDA, B. (1986): Catalogue of the Neotropical Squamata: Part I Snakes. Bulletin of the U. S. National Museum, Washington DC; 297: 1-347. SCHARGEL, W. & FUENMAYOR, G. R. & MYERS, C. W. (2005): An enigmatic new species from cloud forest for the Península de Paria, Venezuela (Colubridae: genus *Taeniophallus*?).- American Museum Novitates, New York; 3484: 1-24. VALENCIA, J. H. (2007 in press): Reptiles del Ecuador; pp. 1-74. In: TORRACCHI, S. (ed.): Fauna del Ecuador. Universidad Técnica Particular del Loja. Loja, Ecuador. VALENCIA, J. H. & MORALES-MITE, M. & HOLMES, D. (2005): Anfibios y reptiles de las provincias de Pastaza y Morona Santiago, Amazonía del Ecuador.- Memorias del Simposio Avances en la Herpetología Neotropical. Instituto Clodomiro Picado. 9 November 2005. CD Room, San José, Costa Rica.

**KEY WORDS:** Reptilia: Squamata: Colubridae: *Echinantera undulata*, new country record, morphology, similar species, Ecuador

SUBMITTED: November 22, 2006

AUTHOR: Jorge H. VALENCIA, Fundación Herpetológica Gustavo Orcés, Av. Amazonas 3008 y Rumipamba, Quito, Ecuador <[jvalencia@fhgo.org.ec](mailto:jvalencia@fhgo.org.ec)>

### First record of *Malpolon monspessulanus* (HERMANN, 1804) from the island of Corsica (France)

In southern Europe, *Malpolon monspessulanus* (HERMANN, 1804) exhibits a wide geographic distribution which is comprised of three main portions (HAAN 1999): (1) western portion (Iberian Peninsula through SE France to NW Italy); (2) middle portion (Balkan Peninsula from about Trieste/NE Italy to Burgas/Bulgaria and Istanbul/Turkey); and (3) eastern portion (eastern Caucasus and western part of Caspian lowlands). This snake has not been recorded from the French Island of Corsica yet (GASC et al. 1997; HAAN 1999).

In early August 2001, GH found a skin shedding of a large snake in a bush alongside a paved road in northern Corsica (Fig. 1), ca. 1.5 km E Cervione (GPS data for Cervione: 42°19'54"N, 9°29'28"E), France. The snake slough (SMF 86351) was deposited in the collection of the Forschungs-




Fig. 1: Habitat of *Malpolon monspessulanus* ca. 1.5 km E Cervione, Corsica, France.  
The arrow indicates the place where the skin shedding (SMF 86351) was found.

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Herpetozoa](#)

Jahr/Year: 2007

Band/Volume: [20\\_1\\_2](#)

Autor(en)/Author(s): Valencia Jorge H.

Artikel/Article: [Presence of Echinantera undulata \(Wied-Neuwied, 1824\) in Ecuador 79-81](#)