

Distribution of *Siphlophis compressus* (DAUDIN, 1803) in Venezuela with a remarkable geographic extension

Siphlophis compressus (DAUDIN, 1803), is the most common member of the genus in northern South America and southern Central America. Two species of the genus *Siphlophis* are known from Venezuela, one of which, *S. cervinus* (LAURENTI, 1768), is apparently widely distributed south of the Orinoco River, although only two localities are known (BARRIO-AMORÓS et al. 1998). On the other hand, *S. compressus*, has been mentioned only in generalized Venezuelan herpetological literature (ROZE 1966; LANCINI 1979; McDIARMID & PAOLILLO 1988; LANCINI & KORNACKER 1989; RODRIGUEZ & FUENTES 1995-1996); and only two recent reports are known: KAISER et al. (2003) and BARRIO-AMORÓS & DUELLMAN (2009). The latter species is widespread south of the Orinoco River, and through eastern Venezuela. Although considered by SAVAGE (2002) and SOLORIZANO (2004) as an uncommon species in Central America, in some areas south of the Orinoco river it can be quite common (like in Río Uei, Sierra de Lema, Estado Bolívar, Venezuela; BARRIO-AMORÓS & DUELLMAN 2009). During an expedition to this last place (BARRIO-AMORÓS & DUELLMAN 2009), three out of six snakes observed in 14 days belonged to this species.

Currently, this snake is known from Costa Rica and Panama south to Bolivia (SAVAGE 2002), with an apparent gap between the Darien area and the area south of the Orinoco River, though it is well known in the Colombian Chocó (CASTRO-HERRERA & VARGAS-SALINAS 2008). All known Venezuelan localities (Fig. 1) were south of the Orinoco River and in the east of the country (BEEBE 1946; KAISER et al. 2003; BARRIO-AMORÓS & DUELLMAN 2009). But because of the species' presence in Central America (e.g. SAVAGE 2002), it is reasonable to assume a wider distribution along the northeastern Andean piedmont, as well as along the piedmont of the Sierra de Perijá, where it has not yet been found. This distribution pattern is followed by other ophidian species like *Drymobius rhombifer* (GÜNTHER, 1860), *Oxybelis fulgidus* (DAUDIN, 1803), and *Xen-*



Fig. 1: Known distribution (white dots) of *Siphlophis compressus* (DAUDIN, 1803) in Venezuela south and east of the Orinoco River based on data in BEEBE (1946), ROZE (1966), LANCINI (1986), McDIARMID & PAOLILLO (1988), LANCINI & KORNACKER (1989), RODRÍGUEZ & FUENTES (1995-1996), KAISER et al. (2003), BARRO-AJAMÓS & DUELLMAN (2009) and new records from the foothills of the Cordillera de Mérida, western Venezuela (red dots). Upper red symbol - 08°54'56"N, 70°26'50"W (BARRO-AJAMÓS & MOLINA 2010), lower red symbols - 07° 54'51"N, 71°43'28"W and 07°47'59"N, 71°45'38"W (this paper).



Fig. 2: *Siphlophis compressus* (DAUDIN, 1803) from near Calderas, Municipio Bolívar, Estado Barinas, Venezuela (08°54'56"N, 70°26'50"W, 900 m a.s.l.).

odon rhabdocephalus (WIED, 1824) (CALCAÑO & BARRIO 2002; SAVAGE 2002, SOLÓRZANO 2004, BARRIO-AMORÓS & MOLINA 2010).

Recently, some members of the Programa Andes Tropicales, a NGO based in Mérida city (Venezuelan Andes), found two specimens of *S. compressus* at the eastern (Orinoquian) versant foothills of the Cordillera de Mérida. One of the animals was photographed (Fig. 2) but not collected. The locality is at the vicinity of the town of Calderas, Municipio Bolívar, Estado Barinas, 08°54'56"N, 70°26'50"W, 900 m a.s.l. (BARRIO-AMORÓS & MOLINA 2010). A second locality, further to the southwest is at Segundo Desarrollo Complejo Hidroeléctrico Uribante-Caparo, Sector Las Cuevas, Municipio Uribante, Estado Táchira, 07°54'51"N, 71°43'28"W, 1200 m, with a voucher specimen at Colección Biológica del Desarrollo Uribante Caparo, Táchira, Venezuela (CBDESURCA-00563). Finally, two more specimens from the same locality, but at slightly lower elevation (07°47'59"N, 71°45'38"W, 790 m) were deposited at Museo Científico de la Universidad Nacional Experimental del Táchira, Venezuela (CVUNET AN-0067 and 0153). All these sites are located within areas classified as premontane humid forest and premontane very humid forest, according to HOLD-RIDGE's system (EWEL & MADRIZ 1968).

The remarkable new records extend the known range of *S. compressus* in Venezuela about 400 km to the NNW. This corroborates once again the Amazonian corridor along the eastern piedmont of the Venezuelan Andes. The senior author and colleagues have already reported many Amazonian species, otherwise known only from the south of the Orinoco River in Venezuela and sometimes with extension to the eastern part of the country (BARRIO-AMORÓS 1998). These include species such as the frogs *Leptodactylus* (*Lithodytes*) *lineatus* (SCHNEIDER, 1799) (BARRIO 1999a), *Hypsiboas boans* (LINNAEUS, 1758) (BARRIO 1999b, 2001), and the snakes *Anilius scytale* LINNAEUS, 1758 (BARRIO et al. 2002), *Micrurus lemniscatus* LINNAEUS, 1758 (BARRIO-AMORÓS & CALCAÑO 2003), and *Epicrates cenchria* (LINNAEUS, 1758) (BARRIO-AMORÓS & DÍAZ DE PASCUAL 2008).

ACKNOWLEDGMENTS: We are grateful to José ALIRIO NAVA who provided the photo and data of the specimen from Calderas, Barinas. We benefited also making the map from data at EBRG (thanks to Francisco BISBAL); MHNLS (thanks to Celsa SEÑARIS), and individual reports from Javier MESA, Alberto BLANCO and Oswaldo FUENTES. Field work in Táchira State was funded by two projects (DIUNET04-005-2006 and DIUNET 04-002-2008) to Andrés CHACÓN.

REFERENCES: BARRIO-AMORÓS, C. L. (1998): Sistemática y biogeografía de los anfibios (Amphibia) de Venezuela.— Acta Biológica Venezolana, Caracas; 18 (2): 1–93. BARRIO, C. L. (1999a): Geographic distribution: Anura: *Lithodytes lineatus*.— Herpetological Review, New Heaven; 30 (1): 50. BARRIO AMORÓS, C. L. (1999b): Geographic distribution. Anura: *Hyla boans*.— Herpetological Review, New Heaven; 30 (4): 230. BARRIO-AMORÓS, C. L. & CALCAÑO, D. (2003): First record of *Micrurus lemniscatus* (LINNAEUS, 1758) from Western Venezuela with comments on coral snakes from the eastern Andean piedmont.— Herpetozoa, Vienna; 16 (1/2): 73–78. BARRIO-AMORÓS, C. L. & DÍAZ DE PASCUAL, A. (2008): Notes on geographic distribution: Reptilia: Boidae: *Epicrates cenchria cenchria*: distribution extension.— Checklist, Campinas, 4 (3): 243–247. BARRIO-AMORÓS, C. L. & DUELLMAN, W. E. (2009): Herpetofauna de la Sierra de Lema, Estado Bolívar, Venezuela; pp. 137–155. In: LASSO, C. A. & SEÑARIS, J. C. & RIAL, B. A. & FLORES, A. L. (eds.): Evaluación rápida de la biodiversidad de los ecosistemas acuáticos de la Cuenca Alta del Río Cuyuní, Guayana Venezolana. RAP Bulletin of Biological Assessment : Boletín RAP de evaluación Biológica / Conservation International, Washington; 55. BARRIO-AMORÓS, C. L. & MOLINA, C. (2010): Herpetofauna del Ramal de Calderas, Andes de Venezuela; pp: 74–80. In: RIAL, B. A. & SEÑARIS, J. C. & LASSO, C. A. & FLORES, A. (eds.): Evaluación rápida de la biodiversidad y aspectos socioecosistémicos del Ramal de Calderas. Andes de Venezuela.- RAP Bulletin of Biological Assessment : Boletín RAP de evaluación Biológica / Conservation International, Washington; 56. BARRIO, A. C. L. & CHACÓN-ORTIZ, A. & DÍAZ DE PASCUAL, A. (2002): Geographic distribution. Serpentes: *Anilius scytale*.— Herpetological Review, New Heaven; 33 (1): 66–67. BARRIO, C. L. & NAVARRETE, L.F. & Fuentes, O. & Mattei, R. (1998): *Siphlophis cervinus* (Serpentes, Colubridae) en Venezuela.— Acta Biológica Venezolana, Caracas; 18 (1): 49–53. BEEBE, W. (1946): Field notes on the snakes of Kartabo, British Guiana, and Caripito, Venezuela.— Zoologica, New York; 31 (1): 11–52. CALCAÑO, D. & BARRIO-AMORÓS, C. L. (2002): Geographic distribution. Serpentes: *Drymobius rhombifer*.— Herpetological Review, New Heaven; 33 (1): 67. CASTRO-HERRERA, F. & VARGAS-SALINAS, F. (2008): Anfibios y reptiles en el departamento del valle del Cauca, Colombia.— Biota Colombiana, Bogotá; 9: 251–277. EWEL, J. J. & MADRIZ, A. (1968): Zonas de vida de Venezuela. Memoria explicativa sobre el mapa ecológico. Caracas (Editorial Sucre), 265 pp. KAISER, H. & BARRIO-AMORÓS, C. L. & RIVAS, G. & GRISMER, L. (2003): Geographic distribution; Serpentes: *Siphlophis compressus*.— Herpetological Review, New Heaven; 34 (2): 170. LANCINI, A. R. (1986): Serpientes de Venezuela. Caracas (Ernesto Armitano Editor), pp. 262. LANCINI, V. A. R. & KORNACKER, P. M. (1989): Die Schlangen von Venezuela.

Caracas (Armitano Editores), pp. 381. McDIARMID, R. W. & PAOLILLO, A. (1988): Herpetological collections: Cerro de la neblina; pp. 667-670. In: BREWER-CARIAS, C. (Ed.): Cerro de la neblina. Resultados de la expedición 1983-1987. Caracas (FUDEC), pp. 922. RODRÍGUEZ, A. & FUENTES, O. (1995-1996): La fauna ofídica observada en dos regiones de la cuenca amazónica venezolana.- Terra, Caracas; 11-12 (20-21): 77-84. ROZE, J. A. (1966): La taxonomía y zoogeografía de los ofidios de Venezuela.- Universidad Central de Venezuela, Ediciones de la Biblioteca, Caracas; 28: 1-362. SAVAGE, J. M. (2002): The amphibians and reptiles of Costa Rica; a herpetofauna between two continents, between two seas. Chicago (The University of Chicago Press), pp. 934. SÓLÓRZANO, A. (2004): Serpientes de Costa Rica / Snakes of Costa Rica. Santo Domingo de Heredia, Costa Rica (INBIO), pp. 792.

KEY WORDS: Reptilia: Squamata: Serpentes: Colubridae: *Siphlophis compressus*, distribution, new country record, Venezuela

SUBMITTED: May, 5, 2010

AUTHORS: César L. BARRIO-AMORÓS, Fundación Andigena, Apartado Postal 210. 5101-A. Mérida, Venezuela < cesarbla@yahoo.com >; A. CHACÓN-ORTIZ, CEVE, Decanato de Investigación, Universidad Nacional Experimental del Táchira, San Cristóbal. Táchira; J. P. DIASPARRA, Zoológico las Delicias, Vivarium, Maracay, Venezuela; A. M. ORELLANA, CEVE, Decanato de Investigación, Universidad Nacional Experimental del Táchira, San Cristóbal, Táchira; Jaime BAUTISTA, Programa Andes Tropicales, Urb. El Encanto Av. 2 c/c 41, Qta. Irma, PA Mérida, 5101; César MOLINA, Instituto de Zoología y Ecología Tropical, Universidad Central de Venezuela, Apartado 47058, Caracas1041-A, Venezuela

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Herpetozoa](#)

Jahr/Year: 2010

Band/Volume: [23_1_2](#)

Autor(en)/Author(s): Barrio-Amoros Cesar L., Chacon-Ortiz A., Diasparra J.P., Orellana A. M., Bautista Jaime, Molina Cesar

Artikel/Article: [Distribution of Siphlophis compressus \(DAUDIN, 1803\) in Venezuela with a remarkable geographic extension 100-103](#)