### Snakes in the Province of Ha'il, Kingdom of Saudi Arabia, including two new records

Ha'il is a Saudi Arabian principality located in the central north of the country between 25°17'N and 28°52'N, and 39°18'E to 44°21'E. It covers an area of 112,444 km<sup>2</sup>. The zoogeography of the Arabian fauna was subject to studies since long (WALLCE 1876; ANDERSON 1896; SMITH 1983; ARNOLD 1987; JOGER 1987; SINDACO & JEREMČENKO 2008; SINDACO et al. 2013). Although there is comprehensive information available about the snakes of Arabia (among recent publications, e.g., GASPERETTI 1988; AL-SADOON 1989; SCHÄTTI & GASPERETTI 1994; EGAN 2007), little is known about the snakes of the Ha'il region. The first herpetological study in Ha'il was conducted by DEKINESH (1991) who reported 13 species of snakes from the region, which include some doubted species records.

In the past few years, there has been increased interest in the herpetology of Ha'il; SHARAWY & ALSHAMMARI (2009) reported five species of venomous snakes from the Aja Mountains, west of Ha'il city. ALSHAM-MARI et al. (2014) analyzed the gene sequences of Saudi Arabian vipers with results that indicate only two species of the genera Cerastes and Echis to be present in Ha'il. Finally, Alshammari & Ibrahim (2015) reported five species of snakes from the Faid Historical Hema, 90 km northeast of Ha'il. The lizard fauna of the the Ha'il region was surveyed and updated (AL-SHAMMARI 2012; ALSHAMMARI & IBRAHIM 2015), however, the number of snake species in Ha'il remained uncertain.

To compensate for this deficit, the authors surveyed a total of 31 localities covering all types of habitats in the Ha'il region (Fig. 1) such as the An-Nufud desert, rocky areas, hard soil, gravel terrain and green fields. The field trips were conducted at least twice per season from 2010 to 2014. Snakes were captured by hand or using tools. To obtain as many snakes as possible, both snakes observed in the wild and others contributed to the lab by volunteers were recorded. Each study site was surveyed at least once every year. Surveys were conducted at each site during day and night for diurnal and nocturnal species, respectively. A total of 56 specimens were examined. The snakes were labeled and preserved in glass jars containing 70 % ethanol or 10 % formalin, and deposited in the Museum of Biology Department at Ha'il University (HUM).

Twelve species of snake belonging to five families (Boidae, Colubridae, Lamprophiidae, Elapidae and Viperidae) were recorded in the Ha'il region (Table 1), two for the first time: *Lytorhynchus diadema* (DU-MÉRIL, BIBRON & DUMÉRIL, 1854), and *Platyceps rhodorachis* (JAN, 1865).

#### *Eryx jayakari* BOULENGER, 1888

Materials: HUM034, 9 June 2010, Al-Fatkha. HUM035, 22 May 2010, Baqa'a. HUM036, 25 September 2011,Qina. HUM037, 2 October 2011, Qina. Observed at An-Nufud desert and Al-Jithiatha.

This burrowing boid species is widespread in the sandy areas of the Arabian Peninsula (GASPERETTI 1988; SINDACO et al. 2013). Within Ha'il, it was previously reported in the areas of Sumayra and Al-Hayit (DEKINESH 1991).

#### Lytorhynchus diadema (DUMÉRIL, BIBRON & DUMÉRIL, 1854)

Materials: HUM038-039, 23 May 2011, Baqa'a. HUM040, 8 June 2012, Al-Fatkha'a. Observed at Qina and Al-Jithiatha.

The species is widespread in the Arabian Peninsula (GASPERETTI 1988; SCHÄTTI & GASPERETTI 1994; SINDACO et al. 2013). This is the first record within the Province of Ha'il. The nocturnal species was found in the middle of the night in sandy desert areas. It inhabits sand deserts, gravel plains and salt flats "sabkhah" (AMR & DISI 2011). LEVITON et al. (1992) suggested treating Lytrorhynchus kennedyi K. P. SCHMIDT, 1939, and L. gaddi NIKOLSKY, 1907, as subspecies of L. diadema to conform the application of the species concept within the L. diadema group. However, SCHÄTTI & GASPERETTI (1994) did not accept the placement of L. gaddi as a subspecies of L. diadema. The examined specimens are characterized by a striking crown-shaped pattern on the head parietal scales. This consists of a dark brown circle on a tan ground color



Fig. 1: Map of the Province of Ha'il (Saudi Arabia), showing the position of the localities studied.
1 - Ha'il city, 2 - Naqbeen, 3 - Haqrusin, 4 - Al-Qaa'id, 5 - Al-Khuttah, 6 - Qina, 7 - Aja mountains,
8 - Al-Fatkha, 9 - Mawqaq, 10 - Al-Mahafar, 11 - Jubbah, 12 - Ash-Shamli, 13 - Darghat, 14 - Al-Hayit,
15 - Al-Hulaifa As-Sufla, 16 - Al-Ghazalah, 17 - Ar-Rawdha, 18 - As-Sib'aan, 19 - Salma mountains,
20 - An-Nai, 21 - Sumayra, 22 - Faid, 23 - Baq'aa, 24 - Al-Jathiatha, 25 - Turubah, 26 - An-Nufud,
27 - As-Sulaymi, 28 - Sahay, 29 - Al-Jafr, 30 - Al-Mubaydi, 31 - Jofayfa.

with a tan center adjoined frontally with a dark bar across the posterior of the frontal shield and across the supraoculars to the eyes, and distally by a dark brown longitudinal bar that runs to the back of the nape. The dorsal part of the body is yellowish brown or brown, with a series of approximately 30 ovoid or rhomboidal splotches; each splotch is edged with some white scales. The ventral part is uniformly creamy in color.

#### Platyceps elegantissimus (GÜNTHER, 1878)

Materials: HUM027, 27 October 2012, Al-Qa'id. HUM053, 5 May 2014, Al-Qa'id. HUM064, 1 May 2014, north of Ha'il.

This species is present in the northwest, southwest and center of the Arabian Peninsula (GASPERETTI 1988; EL-ORAN et al. 1994; SINDACO et al. 2013). Specimens were collected from sandy areas near green fields, earlier records were reported from Al-Khuttah and Qina by DEKINESH (1991). All specimens collected in the present survey were found at more than 1,100 m a.s.l. In a study criticized by ALOUFI & AMR (2015), MASOOD (2012) reported this species at Jazan, southwest Saudi Arabia, from hard sandstone and rocky habitats at heights ranging from 17 to 472 m a.s.l.

#### Platyceps rhodorachis (JAN, 1865)

Materials: HUM051, 20 April 2013, As-Sulaymi. HUM052, 13 March 2014, Naqbeen.

This xerophilic species is found in in the central and southern parts of the Arabian Peninsula (GASPERETTI 1988; SINDACO et al. 2013). This is the first record of *P. rhodorachis* for the Province of Ha'il. Table 1: A list of snake species located in the Saudi Arabian Province of Ha'il, including the corresponding locality numbers as indicated in Figure 1.

| Species                  | Record locality number               |
|--------------------------|--------------------------------------|
| Eryx javakari            | 6, 8, 14, 21, 23, 24, 26             |
| Lytorhynchus diadema     | 6, 8, 23, 24                         |
| Platyceps elegantissimus | 1, 4, 5, 6, 29                       |
| Platyceps rhodorachis    | 2, 27                                |
| Spalerosophis diadema    | 3, 4, 12, 16, 21, 22, 23, 24, 27, 29 |
| Telescopus dhara dhara   | 1, 7, 13, 14                         |
| Atractaspis engaddensis  | 6, 10, 11, 14, 15, 23, 27            |
| Psammophis schokari      | 1, 5, 6, 9, 11, 14, 15, 21, 23       |
| Rhagerhis moilensis      | 6, 10, 11, 14, 15, 23, 24            |
| Walterinnesia aegyptia   | 18, 22, 23                           |
| Cerastes gasperettii     | 1, 5, 9, 8, 13, 14, 19, 22, 23, 24   |
| Echis coloratus          | 2, 9, 8, 20, 22, 27, 28, 31          |
|                          |                                      |

#### Spalerosophis diadema cliffordii (SCHLEGEL, 1837)

Materials: HUM045, 17 May 2011, As-Sib'aan. HUM046, 22 May 2010, As-Sib'aan. HUM047, 16 May 2014, Haqrusin. HUM054-055, 11 May 2014, Al-Jithiatha. Observed at Al-Qaa'id.

This snake is widespread in Arabia, where it occurs from the sea level to high altitudes (GASPERETTI 1988; SCHÄTTI & GASPERETTI 1994). In the Province of Ha'il it was previously collected from Faid, Al-Jafr, Sumayra, Al-Ghazalah, Ash-Shamli and As-Sulaymi (DEKINESH 1991; ALSHAM-MARI & IBRAHIM 2015).

*Spalerosophis diadema cliffordii* is common in the Ha'il region. Snakes were collected from beds of wadis with scant vegetation, as well as in green fields close to the sandy areas of the An-Nufud desert.

#### Telescopus dhara dhara (FORSSKÅL, 1775)

Materials: HUM048, 15 June 2010, Aja Mountains. HUM056, 21 May 2014, north of Ha'il.

This species is found in most parts of the Arabian Peninsula, in particular along the coasts of the Red and Arabian Seas in Yemen to the north of Oman (WERNER 1988; SCHÄTTI & GASPERETTI 1994; AMR & DISI 2011). In the Province of Ha'il, it was previously recorded from Darghat and Al-Hayit (DEKINESH 1991). In this study, *T. dhara* was found in rocky areas close to farms and vegetated land.

#### Atractaspis engaddensis HAAS, 1950

Materials: HUM011, 26 May 2011, Naqbeen. HUM012, 7 December 2011, As-Sulaymi. HUM013, 27 June 2010, Al-Mubaydi. Observed at the foot of a mountain at Naqbeen, about one hour after dusk.

This venomous snake is known from the western and central parts of Saudi Arabia (SCHÄTTI & GASPERETTI 1994; AMR & DISI 2011). Previous records from Ha'il were from Faid, Ar-Rawdha and Darghat (DEKINESH 1991; ALSHAMMARI & IBRAHIM 2015). *Atractaspis engaddensis* is fossorial and a strictly nocturnal species, rarely found on the surface of the ground.

> Psammophis schokari (FORSSKÅL, 1775)

Materials: HUM042-043, 13 June 2010, Baqa'a. HUM044, 9 June 2010, Al-Fatkha. HUM057, 9 July 2013, Qina. Observed north of Ha'il city (roadkill).

This snake is common in Arabia and ubiquitous from sea level to the highest mountains. It was recorded even from some islands in the Red Sea such as Farasan Al-Kebir and Kamaran (CORKILL & COCHRANE 1965; GASPERETTI 1988; SCHÄTTI & GASPE-RETTI 1994). Previous records in the Province of Ha'il were from the settlements Al-Khuttah, Jubbah, Sumayra, Mawqaq, Al-Hulayfah As-Sufla and Al-Hayit. *Psammophis schokari* was found in the sand desert near green fields and shrubs but also in rocky locations.

#### *Rhagerhis moilensis* (REUSS, 1834)

Materials: HUM041, 15 May 2014, Baqa'a. HUM058-059, 11 March 2012, Jubbah. HUM066, 6 June 2014, Al-Jithiatha. Observed at An-Nufud Desert and Al-Mahafar.

This snake is common to most of Arabia (GASPERETTI 1988; SCHÄTTI & GAS-PERETTI 1994; SINDACO et al. 2013). In the Province of Ha'il it was previously recorded from Al-Hayit, Qina and Al-Hulayfah As-Sufla (DEKINESH 1991). *Rhagerhis moilensis* is well adapted to sandy deserts and arid environments in rocky areas. The otherwise diurnal snake becomes crepuscular during the summer.

#### Walterinnesia aegyptia LATASTE, 1887

Materials: HUM049, 25 June 2010, Baqa'a. HUM050, 22 May 2011, Baqa'a. HUM063, 2 May 2013, As-Sib'aan.

The Desert Cobra is found in northern and central Arabia (AMR & DISI 2011; SIN-DACO et al. 2013). In the Province of Ha'il ALSHAMMARI & IBRAHIM (2015) reported this species from Faid. NILSON & RASTE-GAR-POUYANI (2007) separated the eastern populations of *Walterinnesia* in Iran, Iraq, Turkey, Syria and eastern Saudi Arabia from *W. aegyptia* under the name *W. morgani* (MOCQUARD, 1905), while they considered *W. aegyptia* as the valid name of the snake in Egypt, Israel, Palestine, western Arabia, and Jordan.

This nocturnal and fossorial ophidian is mostly seen on agricultural land. The specimen from As-Sib'aan was 161 cm long, which is the longest size ever reported for this species (comp. GASPERETTI 1988 who reported a specimen of 131.9 cm from north of Riyadh).

## Cerastes gasperettii Leviton & Anderson, 1967

Materials: HUM015-020, May 2010, Baqa'a. HUM021, 10 March 2014, Al-Jithiatha. HUM022-024, 3 June 2010, Al-Fatkha'a. HUM025-026, 5 November 2011, Salma Mountains. HUM014, 6 September 2010, Faid. HUM065, 9 August 2013, Ha'il.

*Cerastes gasperettii* is distributed in the entire Arabian Peninsula, including Jordan, Iraq and western Iran (BUSAIS & AL- JUMAILY 2005; AMR & DISI 2011; SINDACO et al. 2013). In the Province of Ha'il it was previously recorded from Darghat, Al-Hayit, Faid, Mawqaq, Turubah and Al-Khuttah (DEKINESH 1991; ALSHAMMARI & IBRAHIM 2015). This viper is most common in sandy habitats.

## *Echis coloratus* GÜNTHER, 1878

Materials: HUM028, 15 April 2011 Jofayfa. HUM029-031, June 2010, Al-Fatkha. HUM032-033, 6 September 2010, Sahay. HUM060-061, 14 May 2014, An-Nai. HUM062, 3 April 2014, Naqbeen. Observed in an oasis-like habitat in rocky areas at Naqbeen just before dusk and in Al-Modayyeh (Salma Mountains) near a small swamp in which the toad, *Sclerophrys arabica* (HEYDEN, 1827), and its tadpoles were found.

*Echis coloratus* is known to occur in eastern Egypt, Sinai, Palestine, Israel, Jordan, and the Arabian Peninsula (BUSAIS & AL-JUMAILY 2005; AMR & DISI 2011; SIN-DACO et al. 2013). In the Province of Ha'il this snake was previously recorded from Faid, Mawqaq and As-Sulaymi (DEKINESH 1991; ALSHAMMARI & IBRAHIM 2015).

DEKINESH (1991) reported the 59 specimens of snakes from the Ha'il area to represent 13 species. However, four records must be considered doubtful: Cerastes vipera (LINNAEUS, 1758), Dyspeltis scabra (LIN-NAEUS, 1758), Hemorrhois ravergieri (Méné-TRIES, 1832), and Naja arabica (SCORTECCI, 1932) (by this author referred to as Naja *haje*). There are no more records of *Cerastes* vipera (North African, eastwards as far as the Sinai Peninsula) and H. ravergieri (Irano-Turanian, isolated populations in Lebanon, Syria, and N Jordan) from the Arabian Peninsula, whereas the other two species are found in its south and southwest only (GASPERETTI 1988; EGAN 2007; SINDAco et al. 2013). The authors of the present note were not able to trace, at the University of Ha'il, any of the 726 reptilian specimens collected by DEKINESH (1991), thus leaving his identifications unverified.

Many snake species of the Arabian Peninsula are widespread and utilize a variety of habitat types, whereas others are considered endangered because they occur in limited areas and small populations. One of these is *Platyceps elegantissimus*, which is frequently killed by native people due its

#### conspicuous color which is suggestive of a venomous snake.

ACKNOWLEDGMENTS: The authors are indebted to Zuhair Amr (Biology Department, Jordan University of Science and Technology, Irbid) for providing advice and suggestions in the manuscript. Thanks is also due to Mohammad Hereher (Ha'il University) for his assistance in drawing the location map, Ann Paterson (Department of Natural Sciences, William Baptist College, Arkansas)who reviewed the final draft and Muna Muhaureq (Leicester University) for improving the English.

REFERENCES: ALOUFI, A. A. & AMR, Z. S. (2015): On the herpetofauna of the province of Tabuk, Northwest Saudi Arabia.- Herpetozoa, Wien; 27 (3/4): 147-158. AL-SADOON, M. K. (1989): Survey of the reptilian fauna of the Kingdom of Saudi Arabia. I. The snake fauna of the central region.- Journal of the College of Science, King Saud University, Riyadh; 1 (1-2): 53-59. ALSHAMMARI, A. M. (2011): Molecular phylogeny of Viperidae family from different provinces in Saudi Arabia.- International Journal of Biology [electronic resource], Toronto; 3: 56-63. ALSHAMMARI, A. M. (2012): Additional records of lizards in Ha'il Province, Saudi Arabia.- Russian Journal of Herpetology, Moskva; 19 (4): 287-291. ALSHAMMARI, A. & IBRAHIM, A. (2015): Lizards and snakes in the Historical Faid Protected Area (Faid Hema), Ha'il region, Saudi Arabia.- Herpetological Conservation and Biology [electronic resource]; 10 (3): 1021-1029. Alshammari, A. M. & El-Abd, E. & Ciccozzi, M. & Lo Presti, A. & Giovanetti, M. & CELLA, E. (2014): Single-gene versus double-gene tree analyses in molecular classification of Saudi venomous snakes.- Arabian Journal for Science and Engineering, Dhahran; 40 (1): 37-49. AMR, Z. & DISI, A. (2011): Systematics, distribution and ecology of the snakes of Jordan.- Vertebrate Zoology, Frankfurt a. M.; 61 (2): 179-266. ANDERSON, J. (1896): A contribution to the herpetology of Arabia. With a preliminary list of the reptiles and batrachians of Egypt. London (R. H. Porter), pp. 122. ARNOLD, E. (1987): Zoogeography of the reptiles and amphibians of Arabia; pp. 245-256. In: KRUPP, F. & SCHNEIDER, W. & KINZELBACH, R. (Eds.): Proceedings of the Symposium on the Fauna and Zoogeography of the Middle East. Mainz [= Beihefte zum Tübinger Atlas des Vorderen Orients, Wiesbaden; (A) 28: 245-256.]. BUSAIS, S. & AL-JUMAILY, M. (2005): Contribution to the reptiles of Yemen: Distribution study of dangerous snakes in Yemen.- University of Aden Journal of Applied & Natural Sciences; 9 (3): 515-524. CORKILL, N. & COCHRANE, J. (1965): The snakes of the Arabian Peninsula and Socotra - Journal of the Bombay Natural History Society, Bombay; 62 (3): 475-506. DEKINESH, S. (1991): The reptilian fauna of Ha'il district, north of Saudi Arabia.- Journal of the Egyptian-German Society of Zoology, Orman; 5: 177-196. EGAN, D. (2007): Snakes of Arabia; Dubai (Motivate Publishing), pp. 208. EL-ORAN, R. & AL-MEL-HEM, W. & AMR, Z. (1994): Snakes of southern Jordan.-Italian Journal of Zoology, London; 61: 359-367. GASPERETTI, J. (1988): Snakes of Arabia.- Fauna of Saudi Arabia, Basel, Riyad; 9: 169-450. JOGER U. (1987): An interpretation of reptile zoogeography in

Arabia, with special references to Arabian herpetofaunal relations with Africa; pp. 257-271. In: KRUPP, F. & SCHNEIDER, W. & KINZELBACH, R. (Eds.): Proceedings of the Symposium on the Fauna and Zoogeography of the Middle East. Mainz [= Beihefte zum Tübinger Atlas des Vorderen Orients, Wiesbaden; (A) 28: 257-271.]. LEVITON, A. E. & ANDERSON, S. C. & ADLER, K. & MINTON, S. A. (1992): Handbook to Middle East amphibians and reptiles. Oxford (Society for the Study of Amphibians and Reptiles - SSAR) [Contributions to herpetology No. 8]; pp. vii, 252. MASOOD, M. F. (2012): Ecological distribution of snakes' fauna of Jazan region of Saudi Arabia.- Egyptian Academic Journal of Biological Science [electronic resource], Abbasia; 4 (1): 183-197. NILSON, G & RASTEGAR-POUYANI, N. (2007): Walterinnesia aegyptia LATASTE, 1887 (Ophidia: Elapidae) and the status of Naja morgani MOCQUARD, 1905.- Russian Journal of Herpetology, Moskva; 14: 7-14. SCHÄTTI, B. & GASPERETTI, J. (1994): A contribution to the herpetology of southwest Arabia.- Fauna of Saudi Arabia, Basel, Riyad; 14: 348-423. Sharawy, S. M. & Alshammari, A. M. (2009): Checklist of poisonous plants and animals in Aja Mountain, Ha'il region, Saudi Arabia.- Australian Journal of Basic and Applied Sciences [electronic resource], Faisalabad; 3 (3): 2217-2225. SINDACO, R. & JEREMČENKO, V. (2008): The reptiles of the western Palearctic, volume 1. Annotated checklist and distributional atlas of the turtles, crocodiles, amphisbaenians and lizards of Europe, North Africa, Middle East and Central Asia; Latina (Edizioni Belvedere), pp. 543. SINDACO, R. & VENCHÌ, A. & GRIECO, C. (2013): The reptiles of the western Palearctic, volume 2: Annotated checklist and distributional atlas of the snakes of Europe, North Africa, Middle East and Central Asia, with an update to Volume 1; Latina (Edizioni Bel-vedere), pp. 579. SMITH, C. H. (1983): A system of world mammal faunal regions. I. Logical and statistical derivation of the regions.- Journal of Biogeography, Oxford; 10: 455-466. WALLACE, A. R. (1876): The geographical distribution of animals, with a study of the relations of living and extinct faunas as elucidating the past changes of the earth's surface; Vols. 1, 2. London (MacMillan & Co), pp. 576, 640. WERNER, Y. (1988): Herpetofauna survey of Israel (1950-85), with comments on Sinai and Jordan and on zoogeographical heterogeneity; pp. 355-388. In: YOM-TOV, Y. & TCHERNOV, E. (Eds.): Zoogeography of Israel: the distribution and abundance at a zoogeographical cross-road. Dordrecht (Junk), 600 pp. [= Monographiae Biologicae 62].

KEYWORDS: Reptilia: Squamata: Serpentes: Lytorhynchus diadema, Platyceps rhodorachis, new record, distribution, Province of Ha'il, Saudi Arabia

#### SUBMITTED: June 6, 2016

AUTHORS: Ahmed Mohajja ALSHAMMARI<sup>1)</sup>, Salem M. BUSAIS (Corresponding author < s.busais@) gmail.com >) 1, 2) & Adel A. IBRAHIM 1, 3)

1) Department of Biology, Faculty of Science, Ha'il University, 81451 Ha'il, Saudi Arabia;

<sup>2)</sup> Department of Biology, Faculty of Education, Aden University, Khormaksar, 6235 Aden, Yemen; <sup>3)</sup> Department of Zoology, Faculty of Science,

Suez University, 43527 Suez, Egypt.

# **ZOBODAT - www.zobodat.at**

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Herpetozoa

Jahr/Year: 2017

Band/Volume: 30\_1\_2

Autor(en)/Author(s): Alshammari Ahmed Mohajja, Busais Salem M., Ibrahim Abdel A.

Artikel/Article: <u>Snakes in the Province of HaÂ'il, Kingdom of Saudi Arabia,</u> including two new records 59-63