

## First record of *Tarentola mauritanica* (LINNAEUS, 1758), from Athens, Greece

The Moorish Gecko, *Tarentola mauritanica* (LINNAEUS, 1758), has a wide distribution comprising North Africa and South Europe (MARTÍNEZ RICA 1997) and has been introduced to numerous overseas localities (MAHRDT 1998; ACHAVAL & GUDYNAS 1983). In Greece, the species has mainly an insular distribution inhabiting four Ionian Islands (Cephalonia, Ithaca, Zakynthos and Strofades) and Crete, whereas the only mainland populations described are located in the northwestern Peloponnese coastal region (WERNER 1927; ŠTĚPÁNEK 1934, 1940; CHONDROPOULOS 1986; VALAKOS & MYLONAS 1992; VALAKOS et al. 2008). The known Greek range of *T. mauritanica* was recently updated with two new island records from Corfu (MACÁT et al. 2014) and Lesvos (MIZERAKIS & STRACHINIS 2017).

On November 6, 2015, a resident from the municipality of Aigaleo, Athens, sent to the first author three photos of a Moorish Gecko found near her residence ( $38.0022^{\circ}$  N,  $23.6789^{\circ}$  E) (F. Retinioti in litt.). Almost one year later, on October 24, 2015, a resi-

dent of Sepolia, a district in the northwest of Athens, found a fatally injured adult male Moorish Gecko there ( $38.0078^{\circ}$  N,  $23.7193^{\circ}$  E), approximately 4 km west of the first observation's location in Aigaleo (A. Rigkos in litt.). He contacted the authors for identification of the lizard and the specimen was deposited at the Zoological Museum of the National and Kapodistrian University of Athens (ZMUA 4185; snout-vent-length: 61 mm, tail broken, body mass: 4.39 g; Fig. 1). Eight months later, on June 28, 2017, during a field survey, the authors observed about 30 individuals of the Moorish Gecko in Sepolia ( $38.0090^{\circ}$  N,  $23.7188^{\circ}$  E), which were active at noon on a rather warm day ( $36^{\circ}$  C). Most of them were seen on old garden walls surrounded by vegetation, in abandoned buildings and parks, and fled when the authors tried to approach them. One individual was captured and examined while several others were photographed.

This is the first time the species is reported in mainland Greece, outside its known distribution in the western Peloponnese. In the last years, new lizard species, either native to Greece such as *Podarcis peloponnesiacus* (BIBRON & BORY, 1833) (HEDMAN et al. 2017) and *Podarcis muralis* (LAURENTI, 1768) (KARAMETA & PAFILES 2017) or alien, such as *Podarcis siculus*



Fig. 1: Male *Tarentola mauritanica* (LINNAEUS, 1758), from northwest Athens, Greece (ZMUA 4185).

(RAFINESQUE-SCHMALTZ, 1810) (ADAMOPOULOU 2015), were reported from Athens metropolitan area. Like the previous species, the Moorish Gecko must have arrived in Athens through human activities. Being a strongly synanthropic species, *T. mauritanica* has only comparatively recently colonized Europe (HARRIS et al. 2004; RATO et al. 2012) and still expands its range (e.g., ARREDONDO et al. 2014; JESUS et al. 2008). Monitoring the extent and rate of dispersal of the highly adaptive *T. mauritanica* to the wider Athens area and assessing its interactions with the widely distributed local gecko *Hemidactylus turcicus* (LINNAEUS, 1758), would provide additional information on the population and expansion dynamics of the species.

**ACKNOWLEDGMENTS:** The authors thank Mrs. Faye Retinioti and Mr. Argiris Rigkos for contacting and providing us with their photos, information and help.

**REFERENCES:** ACHAVAL, F. & GUDYNAS, E. (1983): Hallazgo de *Tarentola mauritanica* (L., 1758) (Lacertilia, Gekkonidae), en el Uruguay.- Boletín de la Sociedad Zoológica del Uruguay, Montevideo; 1: 7-11. ADAMOPOULOU, C. (2015): First record of *Podarcis siculus* (RAFINESQUE-SCHMALTZ, 1810) from Greece.- Herpetozoa, Wien; 27: 187-188. ARREDONDO, C. & NÚÑEZ, H. (2014): *Tarentola mauritanica* (LINNAEUS, 1758), a new species of lizard for Chile (Reptilia, Phyllodactylidae).- Boletín del Museo Nacional de Historia Natural, Santiago; 63: 73-76. CHONDROPOULOS, B. P. (1986): A checklist of Greek reptiles. 2. The lizards.- Amphibia-Reptilia, Leiden; 7: 217 - 235. HARRIS, D. J. & BATISTA, V. & LYMBERAKIS, P. & CARRETERO, M. A. (2004): Complex estimates of evolutionary relationships in *Tarentola mauritanica* (Reptilia: Gekkonidae) derived from mitochondrial DNA sequences.- Molecular Phylogenetics and Evolution, Cambridge; 30: 855-859. HEDMAN, H. & KAPSALAS, G. & KARAMETA, E. & PSONIS, N. & POULAKAKIS, N. & FOUFOPOULOS, J. & PAFILIS, P. (2017): A new locality record and range extension for *Podarcis peloponnesiacus* (BIBRON & BORY, 1833) (Sauria: Lacertidae) from Athens, Greece.- Herpetozoa, Wien; 29: 190-193. JESUS, J. & LEMOS, R. & GONCALVES, R. & BREHM, A. (2008): First record of *Tarentola mauritanica* (LINNAEUS, 1758) on Porto Santo Island.- Herpetozoa, Wien; 20:175-177. KARAMETA, E. & PAFILIS, P. (2017): The first record of *Podarcis muralis* (LAURENTI, 1768) from Athens, Greece.- Herpetozoa, Wien; 30 (1/2): 87-88. MAČÁT, Z. & STARCOVÁ, M. & ČERVENKA, J. & JABLONSKI, D. & ŠANDERA, M. (2014): A molecular assessment and first record of *Tarentola mauritanica* (Squamata: Phyllodactylidae) on Corfu, Greece.- Salamandra, Mannheim; 50 (3): 172-176. MAHRDT, C. R. (1998): Geographic distribution. *Tarentola mauritanica*.- Herpetological Review, New York; 29 (1): 52. MARTÍNEZ RICA, J. P. (1997): *Tarentola mauritanica* (LINNAEUS, 1758); pp. 214-215. In: GASC, J.-P. & CABEZA, A. & CRNOBRNJA-ÍSAILOVIC, J. & DOLMEN, D. &

GROSSENBACHER, K. & HAFFNER, P. & LESCURE, J. & MARTENS, H. & MARTINEZ RICA, J. P. & MAURIN, H. & OLIVEIRA, M. E. & SOFIANDOU, T. S. & VEITH, M. & ZUIDERWIJK, A. (Eds.): Atlas of amphibians and reptiles in Europe; Paris (Societas Europaea Herpetologica & Museum National d'Histoire Naturelle). MIZERAKIS, V. & STRACHINIS, I. (2017): New record of *Tarentola mauritanica* (Squamata: Phyllodactylidae) from Lesvos island, Greece.- Herpetology Notes, Leiden; 10: 157-159. RATO, C. & CARRANZA, S. & HARRIS, D.J. (2012): Evolutionary history of the genus *Tarentola* (Gekkota: Phyllodactylidae) from the Mediterranean Basin, estimated using multilocus sequence data.- BMC Evolutionary Biology, London; 12: 1-12. ŠTĚPÁNEK, O. (1934): Poznámky k herpetologii Kréty.- Sborník Národního Muzea v Praze, Praha; 1: 8-9. ŠTĚPÁNEK, O. (1940): Herpetologie ostrovů Gaudos a Dia.- Sborník Národního Muzea v Praze, Praha; 2: 107-110. VALAKOS, E. D. & MYLONAS, M. (1992): Distribution and ecological aspects of the herpetofauna of Strofades Islands (Ioanian Archipelago, Greece).- Herpetozoa, Wien; 5: 33-39. VALAKOS, E. D. & PAFILIS, P. & SOTIROPOULOS, P. & LYMBERAKIS, P. & MARANGOU, P & FOUFOPOULOS, J. (2008): Reptiles and amphibians of Greece. Frankfurt am Main. (Edition Chimaira), pp. 463. [Frankfurt contributions to natural history No. 32]. WERNER, F. (1927): Beiträge zur Kenntnis der Fauna Griechenlands.- Zoologischer Anzeiger, Jena; 70: 135-143.

**KEY WORDS:** Reptilia: Squamata: Sauria: Phyllodactylidae; *Tarentola mauritanica*, synanthropic species, distribution, Athens, Greece

SUBMITTED: June 11, 2017

AUTHORS: Ilias STRACHINIS (corresponding author < info@herpetofauna.gr ><sup>1)</sup> & Panayiotis PAFILIS < ppafl@biol.uoa.gr ><sup>2)</sup>.

<sup>1)</sup> Section of Genetics, Development and Molecular Biology, Department of Biology, Aristotle University of Thessaloniki, 541 24 Thessaloniki, Greece.

<sup>2)</sup> Section of Zoology and Marine Biology, Department of Biology, National and Kapodistrian University of Athens, 157 84 Panepistimioupolis, Ilisia, Athens, Greece.

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Herpetozoa](#)

Jahr/Year: 2018

Band/Volume: [31\\_1\\_2](#)

Autor(en)/Author(s): Strachinis Ilias, Pafilis Panayiotis

Artikel/Article: [First record of \*Tarentola mauritanica\* \(LINNAEUS, 1758\), from Athens, Greece 98-99](#)