

A case of unilateral anophthalmia in an adult *Pelophylax* kl. *esculentus* (LINNAEUS, 1758)

Various types of morphological abnormalities are found in amphibians all over the world (OUELLET 2000). Among non-skeletal abnormalities that are not related to coloration, anophthalmia (the absence of the eye) and localized or generalized edema are most frequently detected (HENLE et al. 2017).

Out of a large number of documented instances of anuran and urodelan morphological malformations (HENLE et al. 2012; LANNOO 2009), anophthalmia was reported in 159 cases, involving 47 species of Anura and eight of Urodela. The rarity of anophthalmia in natural populations of urodeles was explained by their high capability of regeneration (HENLE et al. 2017).

Probably because they are easily detected, most teratological findings in natural populations of frogs of the hybridogenic *Pelophylax* kl. *esculentus* (LINNAEUS, 1785) complex apply to malformations of the fingers, toes and limbs such as polydactyly (BORKIN & PIKULIK 1986), ectrodactyly, syndactyly (RAKHIMZHANOVA & KHROMOV 1998), polymely (DUMÉRIL 1865), symmely (BRUCH 1864), clinomely (PUKY 2006), phocomely (KURTYAK 2010) and polyphalangy (ZAMALETDINOV 2014) but also microcephaly (FLAX & BORKIN 2004), melanism, blue-colored individuals (MURPHY 1980), translucent skin (DUBOIS 1968) and anophthalmia. As to the genus *Pelophylax*, anophthalmia was described, e.g., in natural populations of *Pelophylax ridibundus* (PALLAS, 1771) (MARUSCHAK & MURAVYNETS 2013) and reproductive communities formed by individuals of *P. lessonae* (CAMERANO, 1882) and *P. kl. esculentus* (LINNAEUS, 1758) (BERGER & UZZEL 1977).

On September 30, 2017, during the study of water frogs, the authors found an adult male *Pelophylax* kl. *esculentus*, who had only one (right) eye. The frog was captured in a pond of about 4 m², located in Pacov, in the Region of Vysocina, Czech Republic (GPS coordinates: 49.474 N, 14.999 E, 580 m a.s.l.). This and two more

ponds lie next to a narrow stream in a grassy area between the dams of large ponds 1-2 ha in size. Fields surrounding Pacov are used mainly for agricultural purposes providing a potential source of contamination by pesticides. No more malformed frogs were observed at this locality. The skin in the area of the missing eye did not show any visible signs of injury, suggesting that the left eye of this specimen had not developed at all or had been heavily injured during an early larval developmental stage. Nonetheless, the frog was in a good nutritional condition and thus, must have been able to hunt prey successfully despite its reduced spatial orientation. After closer examination, the specimen was photographed and released back to the pond. The causes of the observed monophthalmia remain unknown.

ACKNOWLEDGMENTS: This work was supported by the Internal Grant Agency of the Czech University of Life Sciences Prague (CIGA) through Project No. 20172002.

REFERENCES: BERGER, L. & UZZEL T. (1977): Viability and growth of progeny from different egg size classes of *Rana esculenta* L. (Amphibia, Salientia).- Zoologica Poloniae, Warsaw; 26: 292-317. BORKIN, L. J. & PIKULIK, M. M. (1986): The occurrence of polymely and polydactyly in natural populations of anurans of the USSR.- Amphibia-Reptilia, Leiden; 7: 205-216. BRUCH, C. (1864): Über Missbildungen der Chorda dorsalis (Dichordus), nebst Bemerkungen über Doppelbildungen.- Würzburger Medicinische Zeitschrift, Würzburg; 5: 1-35. DUBOIS, A. (1968): Sur deux anomalies de la grenouille verte (*Rana esculenta*).- Bulletin Mensuel de la Société Linnaéenne de Lyon, Lyon; 37: 316-320. DUBOIS, A. (2017): Rostand's anomaly P in Palaearctic green frogs (*Pelophylax*) and similar anomalies in amphibians.- Mertensiella, Rheinbach; 25: 49-56. DUMÉRIL, A. (1865): Observations sur la monstruosité dite polymélies ou augmentation du nombre des membres chez des batraciens anoures.- Nouvelles Archives du Muséum d'Histoire Naturelle, Paris; 1: 309-319. FLAX, N. L. & BORKIN L. J. (2004): Morphological abnormalities and heavy metal concentrations in anurans of contaminated areas, eastern Ukraine.- Applied Herpetology, Leiden; 1: 229-264. HENLE, K. & MESTER, B. & LENGYEL, S. & PUKY, M. (2012): A review of a rare type of anomaly in amphibians, tail duplication and bifurcation, with description of three new cases in European Species (*Triturus dobrogicus*, *Triturus carnifex*, and *Hyla arborea*).- Journal of Herpetology, Houston, etc.; 46: 451-455. HENLE, K. & DUBOIS, A. & VERSHININ V. (2017): A review of anomalies in natural populations of amphibians and their potential causes.- Mertensiella, Rheinbach; 25: 57-164. KURTYAK, F. F. [Куртjak Ф. Ф.] (2010): Аномалії розвитку *Pelophylax* klepton *esculenta* (LINNAEUS, 1758) (Amphibia, Anura, Ranidae) з теренів Закарпаття [Abnormal develop-



Fig. 1: The monophthalmic specimen of *Pelophylax* kl. *esculentus* (LINNAEUS, 1758) from Pacov, Czech Republic. Observe the complete absence of the left eye.
Photo: Tomáš Holer.

ment in *Pelophylax klepton esculenta* (LINNAEUS, 1758) (Amphibia, Anura, Ranidae) from Zakarpattia (=Transcarpathia). - Науковий вісник Ужгородського університету [Scientific Messenger of the Uzhgorod University (= Transcarpathian State University], Series Biology] Uzhgorod; 28: 132-134. [in Ukrainian]. LANNOO, M. J. (2009): Amphibian malformations: pp. 3089-3111. In: HEATWOLE, H. & WILKINSON J. W. (Eds.): Amphibian biology. Volume 8: Amphibian decline: diseases, parasites, maladies and pollution. Chipping Norton (Surrey Beatty and Sons). MARUSCHAK, A. & MURAVYNETS A. (2013): Morphological abnormalities in natural populations of anurans (Amphibia, Anura) in Ukraine.- Praci Ukrainskogo Gerpetologichnogo Tovaristva [Proceedings of the Ukrainian Herpetological Society], Kyiv; 4: 87-94. MURPHY, J. C. (1980): Green, blue, and yellow frogs.- Bulletin of the Chicago Herpetological Society, Chicago; 15: 103-106. OUELLET, M. (2000): Amphibian deformities: current state of knowledge; pp. 617-661. In: SPARLING, D. W. & LINDER, G. & BISHOP C. A. (Eds.): Ecotoxicology of amphibians and reptiles. Pensacola (SETAC Press). PUKY, M. (2006): Amphibian deformity frequency and monitoring methodology in Hungary.- Froglog, Arlington; 74: 3-4. RAKHIMZHANOVA, A. K. & KHRAMOV, V. A. (1998): Morphological study of the green frog (*Rana ridibunda* PALLAS, 1771) from eastern Kazakhstan province.- Abstracts, page 31. Almaty

(Konzhik Publishing House). Third Asian Herpetological Meeting, 1-5 September 1998 in Almaty, Kazakhstan. ZAMALETDINOV, R. I. (2014): Materials on the occurrence of morphological abnormalities in natural populations of anurans in the Tatarstan Republic. pp 105-111. In: VERSHININ, V. L. & DJUBUYA, A. & HENLE K. & PUKY M. (Eds.): Anomalii i patologii amfibii i reptiliii [Аномалии и патологии амфибий и рептилий: методология, эволюционное значение, возможность оценки здоровья среды]. Ekaterinburg (Uralskii Federalnii Universitet).

KEY WORDS: Amphibia; Anura; Ranidae; *Pelophylax* kl. *esculentus*, malformation, teratology, monophthalmia, anophthalmia; veterinary medicine, Czech Republic

SUBMITTED: January 31, 2018

AUTHORS: Tomáš HOLER (Corresponding author < holer.tomas@gmail.com >) ¹⁾ & Daniel KOLESKA ²⁾

¹⁾ Faculty of Environmental Sciences, Czech University of Life Sciences Prague, Kamýcká 129, Praha 6 - Suchdol, 165 00, Czech Republic.

²⁾ Department of Zoology and Fisheries, Faculty of Agrobiology, Food and Natural Resources, Czech University of Life Sciences Prague, Kamýcká 129, Praha 6 – Suchdol, 165 00, Czech Republic.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Herpetozoa](#)

Jahr/Year: 2018

Band/Volume: [31_1_2](#)

Autor(en)/Author(s): Holer Tomas, Koleska Daniel

Artikel/Article: [A case of unilateral anophthalmia in an adult Pelophylax kl. esculentus \(LINNAEUS, 1758\) 103-104](#)