

New records of the anomaly P syndrome in two water frog species (*Pelophylax ridibundus* and *P. lessonae*) in Russia

Anton O. Svinin¹, Ivan V. Bashinskiy², Vitaly V. Osipov⁴, Leonid A. Neymark², Alexander Yu. Ivanov⁴, Oleg A. Ermakov⁴, Spartak N. Litvinchuk^{5,6}

¹ Mari State University, 424000, Mari El, Yoshkar-Ola, Lenin sq., 1, Russia

² A.N. Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences, 119071, Moscow, Leninskiy pr., 33, Russia

³ Privolzhskaya Lesostep State Nature Reserve, 440031, Penza, Okružnaya str., 12a, Russia

⁴ Saratov branch of VNIRO, 410002, Saratov, Chernyshevskogo str., 152, Russia

⁵ Institute of Cytology, Russian Academy of Sciences, 194064, St. Petersburg, Tikhoretsky pr., 4, Russia

⁶ Dagestan State University, 3367000, Dagestan, Makhachkala, Gadzhiev str., 43-a, Russia

<http://zoobank.org/B608B240-390E-4CA8-80F0-4148F04DC000>

Corresponding author: Anton O. Svinin (ranaesc@gmail.com)

Academic editor: Günter Gollmann ♦ Received 10 October 2019 ♦ Accepted 21 November 2019 ♦ Published 6 December 2019

Abstract

The “anomaly P” was described in Palearctic water frogs of the genus *Pelophylax* by Jean Rostand as complex morphological anomalies of water frogs, including polydactyly, brachymely, hind limb oedema, bone outgrowths, spikes, flexions and additional limbs in the inguinal region. In 2016, the anomaly P syndrome was rediscovered in central Russia, confirming the hypothesis concerning its wider distribution. Here, three new records of this syndrome in two species of western Palearctic water frog from Russia are described.

Key Words

morphological anomalies, anomaly P, polydactyly

In the 1950s, the famous French writer and biologist Jean Rostand discovered morphological anomalies of an unknown etiology in water frogs of the genus *Pelophylax*, which he named “the anomaly P” (Rostand 1971). Recently, it is also known as “Rostand’s anomaly” (Dubois 2017). The anomaly P is one of the longest studied cases of amphibian anomalies in Europe (Dubois 1989; 2017; Ouellet 2000). The anomaly P syndrome refers to morphological anomalies of water frogs that have light and severe forms of manifestation. The light form of the anomaly is symmetrical polydactyly. Specimens suffering from severe forms of the anomaly have brachymely, polydactyly, hind limb oedema, bone outgrowths, spikes, flexions and additional limbs in the inguinal region (Dubois 2017). The anomaly P, in comparison with deformi-

ties caused by trematode *Ribeiroia ondatrae* (Johnson et al. 2001; 2002), had no additional full limbs and can have small autopod elements in inguinal regions only. The most important difference is symmetry of bilateral traits: number of structures and their shapes are usually the same on both sides of the body. Rostand (1971) showed that this is not inherited (after laboratory crosses of abnormal specimens), but caused by a certain factor in the environment. He grew larvae from clutches taken from water bodies with deformed frogs in laboratory conditions and did not encounter any anomalies. He tried to influence the tadpoles with various chemicals, increased water salinity and ultraviolet light exposure. However, none of these studies led to similar anomalies (Rostand 1971; Dubois 2017).

Over a long period of research, Rostand (1971) found several additional records of frogs with the anomaly P in France. Severe forms of the anomaly P were revealed in Trévignon and Penloc'h (Concarneau, Finistère), Lingé (Indre), “Marais Vendéen” and Saint-Philbert-de-Grand-Lieu (Loire-Atlantique) (Fig. 1). One locality with severe forms was also found in Morocco and one case was registered in Amsterdam (Hillenius 1959 in Rostand 1971). Polydactyly was much more dispersed and found in different places across France, including Champdieu (Loire), Soustons, Léon, Aureilhan (Landes), Bordeaux (Gironde), Luxeuil-les-Bains (Haute-Saône), Faverois (Territoire-de-Belfort), Fénétrange (Moselle) and Tassenières (Jura) (Dubois 1984). Rostand (1971) suggested the hypothesis that the infectious agent causing the anomaly P was more widespread than it was initially expected when he discovered his first anomalies from the ponds in Concarneau.

The anomaly P syndrome was studied for 20 years until its disappearance in the 1970s. This happened with localities in Trévignon and Saint-Philbert-de-Grand-Lieu. Hence, it was no longer possible to study anomalies in detail with the aim of finding its causative agent (Dubois 2017). The anomaly was encountered under field experiments when tadpoles were reared together with fish (Rostand and Darré 1968). These authors (Rostand and Darré 1969) also observed the same effect in a laboratory when feeding tadpoles with the contents of fish intestines. The experiments were not repeated owing to the disappearance of the anomaly in the “ponds with monsters”. Finally, Rostand (1971) suggested a hypothesis for the induction of the anomalies by a teratogenic virus transmitted by fish or some components of their diet.

For half a century, the anomaly could not be found despite the searches undertaken (Dubois 1979; 1984). However, in 2016, we rediscovered Rostand's anomaly P (Svinin et al. 2019), confirming the hypothesis concerning its wider distribution (Rostand 1971). Water frogs with the anomaly were found in Ostrovtskovskaya Lesosteppe, Privolzhskaya Lesosteppe Nature Reserve, Penza Region, Russia.

Here, we describe three additional new localities from Russia where the anomaly P syndrome was observed in two species of western Palearctic water frog (Fig. 1). All these new localities have very different conditions. The anomaly P was found in both natural and anthropogenic ponds, small lakes and huge reservoirs, in canopy and open landscapes, nature reserves and within settlements.

Locality 1. Edrovo village (57.916640N, 33.625937E, 196 m alt.), Novgorod Region, 2 August 1999. Here, the pool frog (*P. lessonae*) inhabits an artificial pond. Twenty metamorphosed individuals were collected. Two metamorphosed individuals had severe forms of the anomaly P. Four adults were without anomalies. Samples are stored in the herpetological collections of the Zoological Institute of the Russian Academy of Sciences. A re-investigation of the same pond on 17 August 2019, showed the presence of the carnivorous invasive fish *Perccottus glenii* and the complete absence of water frog tadpoles.




Figure 1. Distribution of severe cases of Rostand's anomaly P: 1 – Port Lyautey near Kenitra, Morocco; 2, 3 – Trévignon and Penloc'h, Finistère, France; 4 – Lingé, France; 5 – Saint-Philbert-de-Grand-Lieu, France; 6 – Amsterdam, Netherlands; 7 – Edrovo village, Russia; 8 – Poperechenskaya Steppe, Russia; 9 – Ostrovtskovskaya Lesosteppe, Russia; 10 – Surskoe Reservoir, Russia (photo: *Pelophylax ridibundus* individual with severe cases from Ostrovtskovskaya Lesosteppe).

Locality 2. Poperechenskaya steppe (53.049986N, 44.294268E), part of Privolzhskaya Lesostep' Nature Reserve, Penza Region, Russia. It is located in the upper part of the Khoper River drainage basin (the same as Ostrovtskovskaya Lesosteppe, where the anomaly P was previously found). Despite the fact that these two parts of the reserve are situated approximately 20 km from each other, they were very distant regarding their watercourses. The pond in Poperechenskaya steppe was made by beavers (*Castor fiber*), is located in a steppe landscape and is inhabited by the marsh frog, *P. ridibundus*. In June 2019, a total of 12 tadpoles were caught. All tadpoles had severe forms of the anomaly P (Fig. 2). One of five sub-adult frogs had symmetrical polydactyly.

Locality 3. Surskoye Reservoir (53.032927N, 45.290373E), Penza Region, Russia. The water area of the reservoir is 110 km². The investigated habitat is a small backwater of the reservoir. It was inhabited by the marsh frog, *P. ridibundus*. In June 2019, we caught 110 tadpoles. Of these tadpoles, 104 were normal, two had polydactyly and four had severe forms of the anomaly P. Six adult frogs were caught showing no abnormalities.

The polydactyly of water frogs has attracted scientists for a long time. One of the first cases of polydactyly was discovered by French zoologist Isidore Geoffroy Saint-Hilaire in 1832 (Geoffroy Saint-Hilaire 1832; Rostand 1971). In the recent area of the Russian Federation, polydactyly was described in 1896 by N.A. Kholodkovsky, who observed specimens of water frogs with polydactyly, both on fore and hind limbs from “Khrenovskoy Bor” (Kholodkovsky 1896). Recently, the bibli-



Figure 2. Specimens with the anomaly P found in Surskoe reservoir: A, B – polydactyly, C, D – severe cases of the anomaly P syndrome.

ography provided by Rothschild et al. (2012) includes approximately 40 publications (together with articles written by Jean Rostand) in which polydactyly in water frogs has been described.

Following Rostand (1971), we consider the anomaly P phenomenon more widespread. Indeed, many cases of symmetric polydactyly, especially those presented in both the fore and hind limbs, probably relate to the anomaly P. Polydactyly, however, may be caused by completely different factors. For example, polydactyly and other morphological limb abnormalities can be a result of the influence of radiation and chemicals (Rostand 1971; Vershinin 1989; Ouellet 2000; Henle et al. 2017). At present, it is difficult to separate these cases, especially in field. However, as a rule, polydactyly caused by the infectious agent of the anomaly P had the following characteristics: symmetry (1), a gradient from 6 to more digits (2), presence on both the hind and fore limbs (3) and mass occurrence (4).

Mass and symmetrical polydactyly in water frogs was repeatedly described in Europe (Table 1). In six locations, frogs had polydactyly both on fore and hind limbs; thus, they were very similar to the anomaly P manifestation. Machado et al. (2010) found several cases that undoubtedly belong to the anomaly P. In the marshy areas of Upper Swabia in southern Germany, during 1981, anomalies were found in 13 of 192 specimens (7%) and labelled as “Seelenhofer Ried”. Such a variant of polydactyly was observed in a marsh frog from a pond near Mordovo village and from a pond near the city of Penza in Russia (Zaks 2008; Fayzulin 2012). Three specimens of a total twenty-three (13%), from a pond in the village of Bol’shaya Lipovitsa in Russia, had polydactyly on the hind and forelimbs (Kozhevnikova and Lada 2016).

Symmetric polydactyly on the hind limbs was noted in the village of Fedorovka (2 of 15 specimens, i.e. 13%) and the Botanical Garden of Samara (1 individual from 19, i.e. 5%) in the Samara Region of Russia (Fayzulin et al. 2018).

It was observed in a pond near the village of Alba in Belarus: 46 of 161 individuals (29%) captured between 1977 and 1979 had polydactyly on their hind limbs (Borkin and Pikulik 1986). In 1983, a specimen with symmetrical polydactyly (eight digits) on the hind limbs was found in the village of Staritsa within the Tambov region in Russia (Lada 1999). Mass cases of polydactyly were found in Minai village and Stanytsia Luhanska suburbs of Ukraine (Kurtyak 2005; Marushchak and Muravynets 2018). A case of symmetrical polydactyly (6 digits on the hind limbs) was also found in Romania at Gherța Mică (Sas and Kovacs 2006).

Additional cases of asymmetric polydactyly were revealed in the following localities in the Samara Region of Russia (Fayzulin 2012; Fayzulin et al. 2018): the village of Klimovka (1 from 16, i.e. 6%), the village of Upper Suskan (1 from 17, i.e. 6%), the Gorniy settlement (1 from 10, i.e. 10%) and the city of Ufa (1 from 26, i.e. 4%).

Our records extend the knowledge of the distribution of the infectious agent and most probably, within these ranges, new regions of “hotspots” with a high occurrence of the anomaly P will be discovered. The hind-limb anomalies in anurans impair locomotor performance (Zamora-Camacho and Aragón 2019), which is likely related with reduced survival (Heinen and Hammond 1997). We also never observed severe cases of Rostand’s anomaly P in adult individuals in wild populations, because individuals showing strong deformities may have been eaten by predators or died after metamorphosis due to injuries and open wounds (our and Rostand’s laboratory observations; Rostand 1971). High frequency of anomalies can decrease survival of individuals and lead to reduction of local populations size (Dubois 2017). The monitoring of water bodies with high abnormality rates will detect variation in frequency of abnormal individuals and determine the most significant environmental factors causing such differences. It is expected that the detailed study of such localities will throw light on the nature of the teratogenic agent of the anomaly P.

Table 1. Distribution of polydactyly and heavy cases of the Rostand's anomaly in European water frogs.

| # | Species | Country | Region | Locality | N / E | Literature |
|---|---|-------------|-----------------------|--|------------------------|--|
| Heavy cases of the anomaly P | | | | | | |
| 1 | " <i>P. esculentus</i> " | France | Finistère | Trévignon | 47.798869, -3.845854 | Rostand 1971; Dubois 1979; 1984; 2017 |
| 2 | " <i>P. esculentus</i> " | France | Finistère | Penloc'h | 47.800546, -3.849952 | Rostand 1971; |
| 3 | " <i>P. esculentus</i> " | France | Indre | Lingé | 46.755949, 1.083394 | Rostand 1971; Dubois 1979; 1984; 2017 |
| 4 | " <i>P. esculentus</i> " | France | Loire-Atlantique | Saint-Philbert-de-Grand-Lieu | 47.077709, -1.679565 | Rostand 1971; Dubois 1979; 1984; 2017 |
| 5 | " <i>P. esculentus</i> " | Morocco | Rabat-Salé-Kénitra | Port Lyautey near Kenitra | 34.306902, -6.578357 | R. Lautie, pers. com. in Rostand 1971 |
| 6 | " <i>P. esculentus</i> " | Netherlands | | Amsterdam | 52.358068, 4.950365 | Hillenius 1959 in Rostand 1971 |
| 7 | <i>P. ridibundus</i> | Russia | Penza Region | Ostrovtskovskaya Lesosteppe | 52.816111, 44.461111 | Svinin et al. 2019 |
| 8 | <i>P. lessonae</i> | Russia | Penza Region | Surskoe Reservoir | 53.032927, 45.290373 | This publication |
| 9 | <i>P. ridibundus</i> | Russia | Penza Region | Poperechenskaya steppe | 53.049986, 44.294268 | This publication |
| 10 | <i>P. ridibundus</i> | Russia | Novgorod Region | Edrovo settlement | 57.916640, 33.625937 | This publication |
| Polydactyly | | | | | | |
| Symmetrical polydactyly on both hind and forelimbs | | | | | | |
| 11 | " <i>P. esculentus</i> " | Russia | Voronezh Region | Chrenovskoy Bor | 51.102176, 40.131986 | Kholodkovsky 1896 |
| 12 | " <i>P. esculentus</i> " | France | Ain | Villars-les-Dombes | 45.997765, 5.024091 | Bonnet and Rey 1937; Rostand 1971; Dubois 1979; 1984; 2017 |
| 13 | " <i>P. esculentus</i> " | Germany | Upper Swabia | Seelenhofer Ried | 48.083333, 9.633333 | Machado et al. 2010 |
| 14 | <i>P. ridibundus</i> | Russia | Penza Region | Sosnovka | 53.1688889, 45.0961111 | Zaks 2008 |
| 15 | <i>P. ridibundus</i> | Russia | Samara | Mordovo | 53.176002, 49.438077 | Fayzulin 2012 |
| 16 | <i>P. ridibundus</i> | Russia | Tambov Region | B. Lipovitsa | 52.531394, 41.350586 | Kozhevnikova and Lada 2016 |
| Symmetrical polydactyly on hindlimbs | | | | | | |
| 17 | " <i>P. esculentus</i> " | France | Loire | Champdieu | 45.646192, 4.065011 | Rostand 1971; Dubois 1979; 1984; 2017 |
| 18 | " <i>P. esculentus</i> " | France | Gironde | Bordeaux | 44.890266, -0.575800 | Rostand 1971; Dubois 1979; 1984; 2017 |
| 19 | " <i>P. esculentus</i> " | France | Landes | Soustons | 43.764041, -1.337326 | Rostand 1971; Dubois 1979; 1984 |
| 20 | " <i>P. esculentus</i> " | France | Landes | Léon | 43.897340, -1.320508 | Rostand 1971; Dubois 1979; 1984 |
| 21 | " <i>P. esculentus</i> " | France | Landes | Aureilhan | 44.227901, -1.215550 | Rostand 1971; Dubois 1979; 1984 |
| 22 | " <i>P. esculentus</i> " | Turkey | près d'Istanbul | Tekirdag | 40.926399, 27.400486 | Dubois 1979; 1984 |
| 23 | " <i>P. esculentus</i> " | France | Haute-Saône | Luxeuil-les-Bains | 47.818387, 6.386234 | Dubois 1984 |
| 24 | " <i>P. esculentus</i> " | France | Territoire-de-Belfort | Faverois | 47.518561, 7.033751 | Dubois 1984 |
| 25 | " <i>P. esculentus</i> " | France | Moselle | Fénétrange | 48.841808, 7.003869 | Dubois 1984 |
| 26 | " <i>P. esculentus</i> " | France | Jura | Tassenières | 46.918473, 5.506221 | Dubois 1984 |
| 27 | <i>P. esculentus</i> , <i>P. lessonae</i> | Belarus | Minsk | Alba fishery near Nesvizh | 53.201240, 26.637838 | Borkin and Pikulik 1986 |
| 28 | <i>P. ridibundus</i> | Russia | Tambov Region | Staritsa village | 52.606942, 42.797774 | Lada 1999 |
| 29 | <i>P. esculentus</i> complex | Russia | Tatarstan | Kazan | 55.802554, 49.139874 | Zamaletdinov 2003 |
| 30 | <i>P. ridibundus</i> , <i>P. esculentus</i> | Moldova | | Bugornya | 47.958231, 28.823739 | Bezman-Moseiko et al. 2014 |
| 31 | <i>P. ridibundus</i> , <i>P. esculentus</i> | Moldova | | Plot' | 47.974755, 29.160899 | Bezman-Moseiko et al. 2014 |
| 32 | <i>P. ridibundus</i> | Russia | Samara Region | Fedorovka | 53.466, 49.665 | Fayzulin et al. 2018 |
| 33 | <i>P. ridibundus</i> | Russia | Samara Region | Samara, Botanical Garden | 53.215, 50.179 | Fayzulin et al. 2018 |
| 34 | <i>P. esculentus</i> | Romania | Satu-Mare County | Gherța Mică locality | 47.933333, 23.233333 | Sas and Kovacs 2006 |
| 35 | <i>P. ridibundus</i> | Ukraine | Luhansk region | Stanytsia Luhanska suburbs, Siverskyi Donets river | 48.665029, 39.470719 | Marushchak and Muravynets 2018 |
| 36 | <i>P. esculentus</i> | Ukraine | Zakarpattia | Mynai village | 48.590665, 22.282290 | Kurtyak 2005 |
| Asymmetrical polydactyly on hindlimbs | | | | | | |
| 37 | <i>P. ridibundus</i> | Russia | Samara Region | Klimovka | 53.487, 49.018 | Fayzulin et al. 2018 |
| 38 | <i>P. esculentus</i> | Russia | Samara Region | Verhniy Suskan | 53.818, 49.311 | Fayzulin et al. 2018 |
| 39 | <i>P. lessonae</i> | Russia | Samara Region | Settlement Gornyi | 52.997, 51.061 | Fayzulin et al. 2018 |
| 40 | <i>P. ridibundus</i> | Russia | Samara Region | Ufa, Lokotki | 54.542, 55.931 | Fayzulin et al. 2018 |

Acknowledgments

Authors are thankful to anonymous reviewers for valuable comments. We would like to express our grateful to

Professor Dr. Alain Dubois (Muséum National d'Histoire Naturelle, Paris) for discussions throughout the course of this work. The work was supported by the Russian Foundation of Basic Research No. 18-34-00059.

References

- Bezman-Moseiko OS, Borkin LJ, Rozanov YM, Litvinchuk SN (2014) Mass anomalies of hind limbs in green frogs (*Pelophylax esculentus* complex) in Transnistria: the problem of factors and bioindication. Anomalies and pathologies of amphibians and reptiles: methodology, evolutionary significance, the possibility of assessing the health of the environment. Yekaterinburg: 13–19.
- Bonnet A, Rey M (1937) Sur quelques cas de polydactyly et de schistodactylie observés en série chez la grenouille. Bulletin de la Société zoologique de France 62: 21–25.
- Borkin LJ, Pikulik MM (1986) The Occurrence of Polymely and Polydactyly in Natural Populations of Anurans of the USSR. Amphibia-Reptilia 7: 205–216. <https://doi.org/10.1163/156853886X00019>
- Dubois A (1979) Anomalies and mutations in natural populations of the *Rana esculenta* complex (Amphibia, Anura). Mitteilungen aus dem Zoologischen Museum in Berlin 55: 59–87. <https://doi.org/10.1002/mmnz.4830550108>
- Dubois A (1984) L'anomalie P des Grenouilles vertes (complexe de *Rana* kl. *esculenta* Linné, 1758) et les anomalies voisines chez les Amphibiens. In: Vago C, Matz G (Eds) Comptes rendus du Premier Colloque international de Pathologie des Reptiles et des Amphibiens, Angers, Presses de l'Université d'Angers: 215–221.
- Dubois A (2017) Rostand's anomaly P in Palaearctic green frogs (*Pelophylax*) and similar anomalies in amphibians. Mertensiella (Studies on Anomalies in Natural Populations of Amphibians) 25: 49–56.
- Fayzulin AI (2012) Occurrence and morphological anomalies variety of populations of marsh frog (Anura, Amphibia) of the Middle Volga. Proceedings of the Samara Scientific Center of the Russian Academy of Sciences 14(5): 150–154.
- Fayzulin AI, Chikhlyayev IV, Mineev AK, Kuzovenko AE, Mihaylov RA, Zaripova FF, Popov AI, Ermakov OA (2018) New Data on the Anomalies of Tailless Amphibians of the Volga Basin. Second International Conference on Amphibian and Reptiles Anomalies and Pathology: Methodology, Evolutionary Significance, Monitoring and Environmental Health. KnE Life Sciences: 29–35. <https://doi.org/10.18502/kl.v4i3.2099>
- Geoffroy Saint-Hilaire I (1832) Histoire générale et particulière des anomalies de l'organisation chez l'homme et les animaux. Ouvrage comprenant des recherches sur les caractères, la classification, l'influence physiologique et pathologique, les rapports généraux, les lois et les causes des monstruosités, des variétés et vices de conformation, ou traité de teratology Volume I. Paris: Ballière. Librairie de l'Académie Royale de Médecine, 746 pp. <https://doi.org/10.5962/bhl.title.50400>
- Heinen JT, Hammond G (1997) Antipredator behaviors of newly metamorphosed green frogs (*Rana clamitans*) and leopard frogs (*R. pipiens*) in encounters with eastern garter snakes (*Thamnophis s. sirtalis*). American Midland Naturalist 137: 136–144. <https://doi.org/10.2307/2426762>
- Henle K, Dubois A, Vershinin V (2017) A review of anomalies in natural populations of amphibians and their potential causes. Mertensiella (Studies on Anomalies in Natural Populations of Amphibians) 25: 57–164.
- Johnson PTJ, Lunde KB, Ritchie EG, Reaser JK, Launer AE (2001) Morphological abnormality patterns in a California amphibian community. Herpetologica 57: 336–352.
- Johnson PTJ, Lunde KB, Thurman EM., Ritchie EG, Wray SW, Sutherland DR, Kapfer JM, Frest TJ, Bowerman J, Blaustein AR (2002) Parasite (*Ribeiroia ondatrae*) infection linked to amphibian malformations in the western United States. Ecological Monographs 72: 151–168. [https://doi.org/10.1890/0012-9615\(2002\)072\[0151:PROILT\]2.0.CO;2](https://doi.org/10.1890/0012-9615(2002)072[0151:PROILT]2.0.CO;2)
- Kholodkovsky N (1896) Two examples of the polydactyly. Protocols of meetings of the Imperial St. Petersburg Society of Naturalists 27(1): 74–80, 86–87.
- Kozhevnikova VN, Lada GA (2016) On polydactyly in the marsh frog *Pelophylax ridibundus* (Pallas, 1771) in Tambov Province. Tambov University Bulletin. Series: Natural and Technical Sciences 21(1): 265–268. <https://doi.org/10.20310/1810-0198-2016-21-1-265-268>
- Kurtyak FF (2005) Specificity of the population structure of Western-Palearctic green frogs *Rana esculenta* complex of the flat Transcarpathian. Scientific Bulletin of Uzhgorod University. Series: Biology 16: 172–175.
- Lada GA (1999) Polydactyly in anurans in the Tambov Region (Russia). Russian Journal of Herpetology 6: 104–106.
- Machado C, Kwet A, Schlüter A (2010) Polydactyly and polymely in two populations of *Rana temporaria* and *Pelophylax esculentus* (Anura, Ranidae) in southern Germany. Salamandra 46(4): 239–242.
- Marushchak OY, Muravynets OA (2018) Morphological abnormalities in tailless amphibians (Amphibia, Anura) in Ukraine. Geo & Bio 16: 76–82. <https://doi.org/10.15407/gb.2018.16.076>
- Ouellet M (2000) Amphibian deformities: current state of knowledge. In: Sparling DW, Linder G, Bishop CA (Eds) Ecotoxicology of Amphibians and Reptiles. SETAC Press, Pensacola, Florida, 617–661.
- Rostand J, Darré P (1968) Conditions déterminantes de l'anomalie P chez *Rana esculenta*. Comptes rendus des séances de la Société de biologie et de ses filiales, Paris, 162: 1682–1683.
- Rostand J, Darré P (1969) Action tératogène des déjections de certains poissons sur les larves de *Rana esculenta*. Comptes rendus des séances de la Société de biologie et de ses filiales, Paris, 163: 2033–2034.
- Rostand J (1971) Les étangs à monstres. Histoire d'une recherche (1947–1970). Stock, Paris.
- Rothschild BM, Schultze HP, Pellegrini R (2012) Herpetological osteopathology: annotated bibliography of amphibians and reptiles. Springer, New York, 450 pp. <https://doi.org/10.1007/978-1-4614-0824-6>
- Sas I, Kovacs E-H (2006) Hexadactyly case at a *Rana* kl. *esculenta* sample from the north-western part of Romania. Analele Universității din Oradea. Fascicula Biologie 13: 52–53.
- Svinin AO, Bashinskiy IV, Litvinchuk SN, Neymark LA, Osipov VV, Katsman EA, Ermakov OA, Ivanov AY, Vedernikov AA, Drobot GP, Dubois A (2019) First record of the Jean Rostand's "anomaly P" in the marsh frog, *Pelophylax ridibundus*, in central Russia. Alytes 37: 31–45.
- Vershinin VL (1989) Morphological anomalies of amphibians in the urban area. Russian Journal of Ecology 3: 58–66.
- Zaks MM (2008) On the morphological anomalies of green frogs (*Rana ridibunda*, *R. lessonae*) of the city of Penza. Izvestia V. G. Belinsky PSPU 14 (10): 63–65.
- Zamaletdinov RI (2003) Morphological anomalies in urban populations of tailless amphibians (on the example of Kazan). Current Studies in Herpetology 2: 148–153.
- Zamora-Camacho FJ, Aragón P (2019) Hindlimb abnormality reduces locomotor performance in *Pelobates cultripes* metamorphs but is not predicted by larval morphometrics. Herpetozoa 32: 125–131. <https://doi.org/10.3897/herpetozoa.32.e35654>

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Herpetozoa](#)

Jahr/Year: 2019

Band/Volume: [32](#)

Autor(en)/Author(s): Svinin Anton O., Bashinskiy Ivan V., Osipov Vitaly V., Neymark Leonid A., Ivanov Alexander Yu., Ermakov Oleg A., Litvinchuk Spartak N.

Artikel/Article: [New records of the anomaly P syndrome in two water frog species \(*Pelophylax ridibundus* and *P. lessonae*\) in Russia 277-281](#)