

First record of the Diadem Snake, *Spalerosophis diadema* (Squamata, Colubridae) for Lebanon

Rami Khashab¹, Daniel Jablonski²

¹ *Lebanese Wildlife, Building 3393, ground floor, Beit Meri, Matn, Lebanon*

² *Department of Zoology, Comenius University in Bratislava, Ilkovičova 6, Mlynská dolina, 842 15 Bratislava, Slovakia*

<http://zoobank.org/909C5D45-FB8E-48BC-A883-335C924AE766>

Corresponding author: Daniel Jablonski (daniel.jablonski@uniba.sk)

Academic editor: Yurii Kornilev ♦ Received 30 November 2021 ♦ Accepted 28 February 2022 ♦ Published 5 April 2022

Abstract

We report the first record of the Diadem Snake, *Spalerosophis diadema cliffordii* (Schlegel, 1837) for Lebanon, based on an adult male found in the semi-desert area of the Beqaa Valley. This record increases the number of species of reptiles for Lebanon to 52, with the family Colubridae now comprising 17. The record also highlights the Beqaa Valley as a promising, but currently insufficiently investigated area for further herpetological research due to its biogeographic connection to the large semi-desert and desert areas of Syria.

Key Words

Middle East, Levant, distribution, faunistics, diversity

Lebanon is a small country with a high species and habitat diversity, but herpetologically still insufficiently investigated. Thus, further efforts are necessary for studying species composition, distribution and genetic diversity of local amphibians and reptiles to better understand their evolutionary history and taxonomy in a wider, Middle Eastern context. Lebanon is located in the Levant Region that is characterised by the highest reptile richness and endemism in the Western Palearctic (Ficetola et al. 2018). However, our herpetological knowledge is still limited in Lebanon, including species mapping and genetic diversity (cf. Tamar et al. 2015; Jablonski and Sadek 2019a, b). Despite the long herpetological research tradition in the wider region (e.g. Böttger 1880; Zinner 1967), it was only a genetic approach that allowed, for example, the discovery of the second species of the tree frog *Hyla felixarabica* Gvoždík, Kotlík, Moravec, 2010 in Lebanon (Dufresnes et al. 2019). In addition, several morphologically well-recognised species of herpetofauna [(e.g. *Myriopholis macrorhyncha* (Jan, 1860), *Platyceps rogersi* (Anderson, 1893)] are expected to be present, but have not yet been reported in Lebanon despite their

confirmed presence in the surrounding areas (cf. Sindaco et al. 2013; Bar et al. 2021).

Although the family Colubridae in Lebanon is represented by 16 species and eight genera, the distribution of some of them within the country is still not clear (genus *Eirenis*; Sindaco et al. 2013). The presence of the Diadem snake, *Spalerosophis diadema* (Schlegel, 1837), a widely distributed species in the Western Palearctic, is also unclear in Lebanon. Although the species is listed as part of the snake fauna of Lebanon in some sources (e.g. Disi et al. 2001), it has not been confirmed by field observations, nor museum voucher specimens. It is the only member of the genus in Western Asia with a very unstable subspecific taxonomy comprising of three taxa: *S. d. diadema* (India, Pakistan), *S. d. cliffordii* (Schlegel, 1837) from Mauritania to Turkey and Iran and *S. d. schirasianus* (Jan, 1863) mostly from central and south Asia (Marx 1959; Baig and Masroor 2008). The subspecies *S. d. cliffordii* is known from both neighbouring countries of Lebanon (Sindaco et al. 2013), yet it has never been recorded inside Lebanon despite the availability of

potentially suitable habitats. Here, we report on the first record of *S. diadema* for Lebanon.

An adult male *S. diadema cliffordii* (Fig. 1A–F) was killed by local people and then collected by Wihad Dandich on 2 June 2021, near the village of Ras El Assi (ريصاعلا سار), Hermel area, Beqaa Valley, north-eastern Lebanon (34.3342°N, 36.4017°E, datum WGS84; elevation 736 m; Fig. 2A, B). The specimen was found in an open dry semi-desert area with condensed sandy soil and rocks and with small agricultural fields scattered around (Fig. 1H). Small rocky valleys with caverns and invaginations in the walls are present in the wider vicinity that represents a suitable habitat for the species (e.g. Disi et al. 2001).

The specimen was immediately frozen for further examination under the voucher number LW-R-001/2021 and is stored in 70% ethanol in the collection of the Lebanese Wildlife NGO, Beit Meri, Matn, Lebanon. The snout-vent length (SVL) of the specimen is 1,100 mm with 204 ventral scales, tail length (TL) is 210 mm with

73 subcaudal scales. The weight of the specimen is 460 g. The length and weight were measured by tape and digital scales after the snake was frozen. The colouration and the pattern are typical for *S. diadema cliffordii* (Marx 1959; Schätti et al. 2009; Fig. 1A), comprising brown blotches on a light, sandy-orange background. Ventral scales are light with reddish places. The colouration of the body and eyes, however, changed partly due to preservation in ethanol.

Starting in the spring of 2018, we conducted 13, one or two-days long field trips to the area. Day, as well as night surveys, were conducted. Reptile species recorded in the Hermel area of the Beqaa Valley, where the specimen of *S. diadema* was found, were *Trapelus ruderatus* (Olivier, 1804) (Agamidae), *Mesalina microlepis* (Angel, 1936), *Ophisops elegans* Ménétries, 1832 (both Lacertidae), *Eumeces schneiderii pavimentatus* (Geoffroy De St. Hilaire, 1827) (Scincidae), *Telescopus nigriceps* (Ahl, 1924), *Eirenis* aff. *coronella* (Schlegel, 1837), *Dolichophis jugularis*




Figure 1. The specimen of *Spalerosophis diadema cliffordii* from Lebanon. **A, B.** The dorsal and ventral view of the body; **C–F.** Detail views of the head.




Figure 2. A. The map of Lebanon indicating the position of the record (red star); B. The habitat where the specimen of *Spalerosophis diadema cliffordii* was found.

(Linnaeus, 1758) (all Colubridae), *Malpolon insignitus* (Geoffroy Saint Hilaire, 1827) (Psammophiidae), and *Testudo graeca* Linnaeus, 1758 (Testudinidae) (Bosch et al. 1998; Hraoui-Bloquet et al. 2002; Jablonski and Khashab, pers. obs.). According to our observations, dense rodent holes present in the locality provide good shelter for some reptiles, including *S. diadema*. In these shelters, we also found snake sheds from *D. jugularis*, *M. insignitus*, as well as other, unidentified snake species, possibly *Platycephalus* sp. Our record of *S. diadema* thus increases the number of species of reptiles for Lebanon to 52, with the family Colubridae now comprising 17 (cf. Mahlow et al. 2013; Sindaco et al. 2013; Jablonski and Sadek 2019a).

In the Levant Region that lies between the Nur Mountains and the Sinai Peninsula, *S. diadema* is found in semi-desert to desert habitats, rocky slopes or border rocky areas near sandy dunes (Disi et al. 2001; Werner 2016; Bar et al. 2021). The semi-desert areas of Lebanon (continuing from dry areas of the Homs Desert in Syria) are limited only to the northern part of the Beqaa Valley (Fig. 2A) that coincides with the edge area of the range

of several other species of reptiles with affiliation to such environment, for example, *Trapelus ruderatus*, *Eumeces schneiderii pavimentatus* or *Mesalina microlepis* (see Hraoui-Bloquet et al. 2002). Thus, the reason why *S. diadema* was recorded for Lebanon only now is most probably due to the small-size of suitable habitats limited to the northern part of the Beqaa Valley and limited research in the area due to the overall region's poor accessibility connected with the security situation. However, the Beqaa Valley represents probably an important biogeographical corridor for reptile species inhabiting arid environments connecting to Syrian deserts and thus provides the potential for newly-recorded species in Lebanon.

Acknowledgements

We thank Wihad Dandich for his help during the fieldwork. We thank two anonymous reviewers and the editor for their comments. This study was supported by the grant of the Scientific Grant Agency of the Slovak Republic VEGA 1/0242/21.

References

- Baig KL, Masroor R (2008) The snakes of the genus *Spalerosophis* Jan, 1865 in Indo-Pakistan and Iran (Squamata: Serpentes: Colubridae). *Herpetozoa* 20: 109–115.
- Bar A, Haimovitch G, Meiri S (2021) Field Guide to the Amphibians and Reptiles of Israel. Edition Chimaira, Frankfurt am Main, 511 pp.
- Bosch HAJ, Bischoff W, Schmidler JF (1998) Bemerkenswerte Reptilienfunde im Libanon. *Herpetofauna* 20: 19–32.
- Böttger O (1880) Die Reptilien und Amphibien von Syrien, Palaestina und Cypern. Senckenbergische naturforschende Gesellschaft 1880: 132–219.
- Disi AM, Modrý D, Nečas P, Rifai L (2001) Amphibians and reptiles of the Hashemite Kingdom of Jordan: An Atlas and Field Guide. Chimaira, Frankfurt am Main, 408 pp.
- Dufresnes C, Mazepa G, Jablonski D, Sadek RA, Litvinchuk SN (2019) A river runs through it: tree frog genomics supports the Dead Sea Rift as a rare phylogeographical break. *Biological Journal of Linnean Society* 128: 130–137. <https://doi.org/10.1093/biolinnean/blz076>
- Ficetola GF, Falaschi M, Bonardi A, Padoa-Schioppa E, Sindaco R (2018) Biogeographical structure and endemism pattern in reptiles of the Western Palearctic. *Progress in Physical Geography* 42: 220–236. <https://doi.org/10.1177/0309133318765084>
- Hraoui-Bloquet S, Sadek RA, Sindaco R, Venchi A (2002) The herpetofauna of Lebanon: new data on distribution. *Zoology in the Middle East* 27: 35–46.
- Jablonski D, Sadek RA (2019a) The species identity and biogeography of *Blanus* (Amphisbaenia: Blanidae) in Lebanon. *Zoology in the Middle East* 65: 208–214. <https://doi.org/10.1080/09397140.2019.1604471>
- Jablonski D, Sadek RA (2019b) The Caucasian Toad, *Bufo verrucosissimus* (Pallas, 1814) in the Levant: evidence from mitochondrial DNA. *Herpetozoa* 32: 255–258. <https://doi.org/10.3897/herpetozoa.32.e37560>

- Mahlow K, Tillack F, Schmidtler JF, Müller J (2013) An annotated checklist, description and key to the dwarf snakes of the genus *Eirenis* Jan, 1863 (Reptilia: Squamata: Colubridae), with special emphasis on the dentition. *Vertebrate Zoology* 63: 41–85.
- Marx H (1959) Review of the Colubrid snake genus *Spalerosophis*. *Fieldiana. Zoology* 39: 347–361. <https://doi.org/10.5962/bhl.title.7187>
- Schätti B, Tillack F, Helfenberger N (2009) A contribution to *Spalerosophis microlepis* Jan 1865, with a short review of the genus and a key to the species (Squamata: Serpentes: Colubridae). *Herpetozoa* 22: 115–135.
- Sindaco R, Venchi A, Grieco C (2013) The Reptiles of the Western Palearctic 2. Annotated checklist and distributional atlas of the snakes of Europe, North Africa, the Middle East and Central Asia, with an update to the vol. 1. *Societas Herpetologica Italica, Via Adige, Latina*, 543 pp.
- Tamar K, Carranza S, Bosch H, Sindaco R, Moravec J, Meiri S (2015) Hidden relationships and genetic diversity: Molecular phylogeny and phylogeography of the Levantine lizards of the genus *Phoenicolacerta* (Squamata: Lacertidae). *Molecular Phylogenetics and Evolution* 91: 86–97. <https://doi.org/10.1016/j.ympev.2015.05.002>
- Werner YL (2016) *Reptile life in the Land of Israel*. Frankfurt am Main, Edition Chimaira, 494 pp.
- Zinner H (1967) Herpetological collection trips to the Lebanon 1965 and 1966. *Israel Journal of Zoology* 16: 49–58.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Herpetozoa](#)

Jahr/Year: 2022

Band/Volume: [35](#)

Autor(en)/Author(s): Khashab Rami, Jablonski Daniel

Artikel/Article: [First record of the Diadem Snake, *Spalerosophis diadema* \(Squamata, Colubridae\) for Lebanon 77-80](#)