

# Gigantism in tadpoles and color anomalies in European green toad *Bufo viridis* from Poland

Michał Szkudlarek<sup>1</sup>, Jan Marek Kaczmarek<sup>2</sup>, Marta Piasecka<sup>3</sup>, Wiktor Grenas<sup>3</sup>, Mikołaj Kaczmarski<sup>2</sup>

<sup>1</sup> Department of Zoology, Institute of Biological Sciences, University of Zielona Góra, Prof. Z. Szafrana 1, 65-516 Zielona Góra, Poland

<sup>2</sup> Department of Zoology, Poznań University of Life Sciences, Wojska Polskiego 71c, 60-625 Poznań, Poland

<sup>3</sup> Poznań, Poland

<http://zoobank.org/34BB0BCF-8B39-4C1D-BD29-830617470621>

Corresponding author: Michał Szkudlarek ([michalszkudlarek@protonmail.com](mailto:michalszkudlarek@protonmail.com))

Academic editor: Günter Gollmann ♦ Received 23 March 2022 ♦ Accepted 9 May 2022 ♦ Published 7 June 2022

## Abstract

Some new cases of anomalies in wild *Bufo viridis* from Western Poland are presented together with a discussion of their potential causes and a comparison with similar cases from the literature. The anomalies in question are color aberrations (pre- and post-metamorphic) and developmental pathologies (pre-metamorphic). The color anomalies include an ochre patch in a juvenile, leucism and albinism in both tadpoles and juvenile individuals. Developmental pathologies described in this paper include gigantism, edema, curvature of tail, and asymmetric bodies.

## Key Words


aberration, Bufonidae, leucism, pathologies

Aberrant amphibian individuals can be found both in large (due to mere populations' size) and small populations. The latter is partly due to genetic drift (Hitchings and Beebee 1997) and, especially in the case of populations that are genetic isolates (like many urban amphibian populations), inbreeding (Hitchings and Beebee 1997; Vershinin 2004; Weyrauch and Grubb 2006). Weakened predation pressures in urban habitats might hypothetically also contribute to a higher prevalence of aberrant individuals in such places due to them usually being of lower fitness, hence benefiting from the scarcity of predators (Strachinis and Tsarouhas 2021). Atypically colored amphibians, usually having other co-occurring anomalies and looking less cryptic, show a lower survival rate than normally colored ones (Henle et al. 2017a); however, it seems that they are now quite frequently reported and attract general interest (Jablonski et al. 2014; Henle et al. 2017a; Kolenda et al. 2017). To date, several color anomalies have been reported in the European green toads *Bufo viridis* (Laurenti, 1768), including

albinism, axanthism, erythrism, leucism, abnormal ochre patches and retinal depigmentation (Flindt 1985; Andrä 2011; Jablonski et al. 2014; Henle et al. 2017c; Lunghi et al. 2017; Strachinis and Tsarouhas 2021). Herein, we present several rare or previously unreported color anomalies in *B. viridis* from Poland.

On 4 August 2017 at about 20:00 we found and photographed a juvenile individual of *B. viridis* with an ochre patch extending from the middle of the parotid glands onto the head (Fig. 1A, B). The observation was made in Tarnowo Podgórze, near man-made water reservoirs [52°27'53"N, 16°38'38"E], where dozens of freshly metamorphosed conspecifics with typical coloration were present. Unfortunately, the individual was in a closed and fenced area, so it was impossible to catch it for a more detailed inspection. As far as we observed, it did not behave differently from typically colored conspecifics.

Divergent patterns and colorations other than melanism and albinism can be caused by parasites,



**Figure 1.** Aberrant juveniles of *B. viridis* from Poland. **A, B.** Juvenile with an ochre patch; **C, D.** Juvenile leucistic individual.

diseases, chemicals, inbreeding, radioactivity, or other mutagenic factors (Henle et al. 2017c). To date, *B. viridis* individuals with abnormal yellow/ochre patches were found in 1980 at a quarry in Roßwag, Germany (Henle et al. 2017c). Those individuals had yellow patches of various sizes either on the flanks or on the dorsum (see fig. 13 in Henle et al. (2017c) and fig. 32 in Henle et al. (2017a)). Henle et al. (2017c) crossed one of them with a normally colored individual, and 25% of the offspring developed ochre patch/patches, which indicates that they were heritable. The authors also speculated that the mutation was most likely caused by irradiation from radioactive contaminants. In our case, a sewage treatment plant located ca 200 m away and/or a waste pond for thermal spring water located ca 50 m away could have been the sources of mutagenic factors.

Atypical yellow dorsal spots dubbed as partial flavism were observed in the yellow-bellied toad *Bombina variegata* (Linnaeus, 1758) and in the spadefoot toad *Pelobates fuscus* (Laurenti, 1768) in two localities in Poland (Kolenda et al. 2017). In *B. variegata* yellow is part of the normal ventral coloration so only a different location of that spot can be considered abnormal. *Pelobates fuscus*, on the other hand, had many yellow spots of various sizes. Unfortunately, we do not have any histological skin sections that would allow a better diagnosis of the described cases.

In 2019, in an artificial pond in the city park Rosarium, Poznań [52°25'26"N, 16°55'56"E], we found

free-swimming whitish tadpoles (Gosner stage 25 (Gosner 1960)) of *B. viridis* among several thousands of normally pigmented ones (see the description of this site in Kaczmarski et al. 2019). We caught 10 aberrant and 10 typically colored tadpoles for further rearing. In most atypical individuals, the background color was dirty white or light grey, but the eyes were typically colored (Fig. 2A). During development 6 whitish individuals were affected by edema, curvature of tail and asymmetric bodies. Nine out of 10 reared whitish individuals had trouble freeing their front limbs and finally died before the end of metamorphosis. The color of the individual who survived metamorphosis can be described as leucistic, that is: white, with slightly pigmented spots and typically colored eyes (Fig. 1C, D). All individuals with normal coloration did not show any malformations and underwent metamorphosis without problems.

*Bufo viridis* tadpoles typically reach a maximum total length of 52 mm (Günther and Podloucky 1996, according to Henle et al. 2017c), but in our case, three individuals with color aberration reached even 80 mm total length (Fig. 2B), which allows them to be classified as giant (see fig. 17, in Henle et al. (2017c)). As far as we know, gigantism is presumed to be more common among hybrids of *Bufo bufo* (Linnaeus, 1758) and *B. viridis* (Flindt (1984) in Henle et al. (2017c)). However, we confirmed that the here described gigantic tadpoles belonged to *B. viridis* by PCR-amplifying and sequencing



**Figure 2.** Aberrant tadpoles and juvenile of *B. viridis* from Poland. **A.** Whitish tadpole with a normally pigmented one; **B.** Leucistic giant (52 mm–80 mm TL) tadpoles; **C.** Albinistic tadpoles, some of which are undergoing metamorphosis; **D.** Freshly metamorphosed albinistic individual. Eyes are red, which classifies this individual as a complete albino.

one mitochondrial gene (16S rRNA) and one nuclear gene (POMC) and comparing them to reference sequences from *Bufo bufo*, *Epidalea calamita* and *Pseudepidalea viridis/variabilis*. Arms remaining within the branchial cavity can be caused by chemical pollution (Henle et al. 2017c). Curvature of tail or body might be related to chemical pollution, inbreeding, increased natural level of UV-B, radioactivity, or other mutagenic factors (Henle et al. 2017c). Gigantism in tadpoles can be induced by chemical pollution or inbreeding (Henle et al. 2017c). Gigantism and leucism in tadpoles of *B. viridis*, along with 30 other anomalies in that species among tadpoles, toadlets, and adults, were described from a quarry near Roßwag (Henle 1981, 1982; Henle et al. 2017c). In our

case, the population is located in the city center and occupies a large park area. However, the relatively small number of tadpoles with unusual coloration indicates that the frequency of this mutation does not differ from the norm. Similarly, one leucistic tadpole was recorded in an artificial pond in Milan along with hundreds of normally pigmented conspecifics (Lunghi et al. 2017).

In May 2021, in a temporary pool in an industrial/rural area of Poznań [52°26'24"N, 16°52'53"E], we found 30 free-swimming completely albinistic tadpoles (Gosner stage 25 (Gosner 1960)) of *B. viridis* among several thousands of normally pigmented ones. We caught 10 aberrant and 10 normally colored tadpoles for further rearing. Their development took place without any

problems (Fig. 2C), and the young individuals underwent metamorphosis faultlessly (Fig. 2D). Although albinism in *B. viridis* has already been described several times in the literature (Flindt 1985; Andrä 2011; Lunghi et al. 2017; Strachinis and Tsarouhas 2021), as far as we know, our case is the second one recorded in Poland (Życzyński 2002). Albinism has been associated with spontaneous tyrosinase mutations in three Japanese frog species (Miura et al. 2018). In *B. viridis* color aberrations can be heritable, caused by recessive genes (Boschwitz 1963). They can also be transient; in some cases, the eggs are white, and the hatched tadpoles acquire pigmentation over time (Christaller 1983; Flindt 1985; Henle et al. 2017b). With intensifying anthropogenic pressure, and cameras becoming more accessible and omnipresent, we are likely to see more reports of anomalies in amphibians. However, their causes can be unravelled only with the use of controlled conditions that would allow the measurement of as many potential causal factors as possible.

The authors are thankful to Maciej Pabijan (Jagiellonian University) for genetic testing of the gigantic tadpoles and to three anonymous reviewers for their valuable comments and remarks. The authors have declared that no competing interests exist.

## References

- Andrä E (2011) Aspekte der Biologie der Wechselkröte. In: Wechselkröten-Symposium des LBV, Munich (Germany), May 2011, 1–14. [http://www.lars-ev.de/pdf\\_pub/Andrae\\_Aspekte%20der%20Biologie%20der%20Wechselkroete\\_Web.pdf](http://www.lars-ev.de/pdf_pub/Andrae_Aspekte%20der%20Biologie%20der%20Wechselkroete_Web.pdf)
- Boschwitz D (1963) Histological investigation of inherited incomplete albinism. *Zoologischer Anzeiger* 170: 19–23.
- Christaller J (1983) Vorkommen, Phänologie und Ökologie der Amphibien des Enzkreises. *Jahreshefte der Gesellschaft für Naturkunde in Württemberg* 138: 153–182.
- Flindt R (1985) Latenter Albinismus und Mißbildungen bei Kaulquappen von Wechselkröten *Bufo viridis* LAURENTI, 1768. *Salamandra* 21: 298–303. <https://www.salamandra-journal.com/index.php/home/contents/1985-vol-21/1116-flindt-r/file>
- Gosner KL (1960) A simplified table for staging anuran embryos and larvae with notes on identification. *Herpetologica* 16(3): 183–190. <https://www.jstor.org/stable/3890061>
- Henle K (1981) A unique case of malformations in a natural population of the green toad (*Bufo viridis*) and its meaning for environmental politics. *British Herpetological Society Bulletin* 4: 48–49. <https://www.thebhs.org/publications/the-herpetological-bulletin/issue-number-4-winter-1981/2105-hb004-07/file>
- Henle K (1982) Ein außergewöhnlicher Fall von Mißbildungen bei der Wechselkröte (*Bufo viridis*). *Unicornis* 2: 40–42.
- Henle K, Dubois A, Vershinin V (2017a) A review of anomalies in natural populations of amphibians and their potential causes. *Mertensiella* 25: 57–164.
- Henle K, Dubois A, Vershinin V (2017b) Commented glossary, terminology and synonyms of anomalies in natural population of amphibians. *Mertensiella* 25: 9–48.
- Henle K, Dubois A, Rimpp K, Vershinin V (2017c) Mass anomalies in green toads (*Bufo viridis*) at a quarry in Roßwag, Germany: inbred hybrids, radioactivity or an unresolved case?. *Mertensiella* 25: 185–242.
- Hitchings SP, Beebe TJC (1997) Genetic substructuring as a result of barriers to gene flow in urban *Rana temporaria* (common frog) populations: implications for biodiversity conservation. *Heredity* 79: 117–127. <https://doi.org/10.1038/hdy.1997.134>
- Jablonski D, Alena A, Vlcek P, Jandzik D (2014) Axanthism in amphibians: A review and the first record in the widespread toad of the *Bufo viridis* complex (Anura: Bufonidae). *Belgian Journal of Zoology* 144: 93–101. <https://doi.org/10.26496/bjz.2014.69>
- Kaczmarek M, Szala K, Kloskowski J (2019) Early onset of breeding season in the green toad *Bufo viridis*. *Herpetozoa* 32: 109–112. <https://doi.org/10.3897/herpetozoa.32.e35825>
- Kolenda K, Najbar B, Najbar A, Kaczmarek P, Kaczmarek M, Skawiński T (2017) Rare colour aberrations and anomalies of amphibians and reptiles recorded in Poland. *Herpetology Notes* 10: 103–109. <https://www.biotaxa.org/hn/article/view/22366>
- Lunghi E, Monti A, Binda A, Piazzini I, Salvadori M, Cogoni R, Riefolo LA, Biancardi C, Mezzadri S, Avitabile D, Ficetola GF, Mulargia M, Manca S, Blaimont P, Di Cerbo AR, Manenti R (2017) Cases of albinism and leucism in amphibians in Italy: new reports. *Natural History Sciences* 4: 73–80. <https://doi.org/10.4081/nhs.2017.311>
- Miura I, Tagami M, Fujitani T, Ogata M (2018) Spontaneous tyrosinase mutations identified in albinos of three wild frog species. *Genes & Genetic Systems* 92(4): 189–196. <https://doi.org/10.1266/ggs.16-00061>
- Strachinis I, Tsarouhas N (2021) First report of albino green toads *Bufo viridis* in Greece. *Herpetological Bulletin* 155: 44–45. <https://doi.org/10.33256/hb155.4445>
- Vershinin VL (2004) Frequency of iris depigmentation in urban populations of *Rana arvalis* Frogs. *Russian Journal of Ecology* 35: 58–62. <https://doi.org/10.1023/B:RUSE.0000011112.17428.f5>
- Weyrauch SL, Grubb TC (2006) Effects of the interaction between genetic diversity and UV-B radiation on wood frog fitness. *Conservation Biology* 20: 802–810. <https://doi.org/10.1111/j.1523-1739.2006.00334.x>
- Życzyński A (2002) Ochrona herpetofauny w warszawskim kompleksie parkowym Pola Mokotowskie. *Przegląd Przyrodniczy* 3: 137–140.

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Herpetozoa](#)

Jahr/Year: 2022

Band/Volume: [35](#)

Autor(en)/Author(s): Szkudlarek Micha, Kaczmarek Jan Marek, Piasecka Marta, Grenas Wiktor, Kaczmarski Mikolaj

Artikel/Article: [Gigantism in tadpoles and color anomalies in European green toad \*Bufo viridis\* from Poland 95-98](#)