

First record of *Anatololacerta pelasgiana* (Mertens, 1959) in mainland Greece: another new species in Athens

Apostolos Christopoulos¹, Charikleia-Foteini Pantagaki^{2,3}, Nikos Poulakakis^{2,3,4}, Panayiotis Pafilis^{1,5}

¹ Section of Zoology and Marine Biology, Department of Biology, National and Kapodistrian University of Athens, Panepistimioupolis, 15784 Athens, Greece

² Department of Biology, School of Sciences and Engineering, University of Crete, Heraklion, Greece

³ Natural History Museum of Crete, School of Sciences and Engineering, University of Crete, Heraklion, Greece

⁴ Institute of Molecular Biology and Biotechnology (IMBB), Foundation for Research and Technology – Hellas (FORTH), Heraklion, Greece

⁵ Zoological Museum, National and Kapodistrian University of Athens, Greece, Panepistimioupolis, 15784 Athens, Greece

<https://zoobank.org/DC996BE2-22C2-43D6-8050-26E9C281BDF2>

Corresponding author: Apostolos Christopoulos (laniusapo@yahoo.gr)

Academic editor: Peter Mikulíček ♦ Received 16 November 2022 ♦ Accepted 2 December 2022 ♦ Published 14 December 2022

Abstract

Urban habitats receive an increasing number of species due to anthropogenic activities, mainly transportations. Here, we report a new addition to the herpetofauna of Athens (Greece): a small population of the Pelasgian wall lizard (*Anatololacerta pelasgiana*) was found in a suburb of the Athenian metropolitan area. The species normally occurs in southwestern Anatolia and southeastern Aegean islands and this is the first record in the Greek mainland. Allochthonous species that successfully colonize cities raise new challenges to urban ecology.

Key Words

introduction, lizard, Mediterranean, phylogenetic analysis, urban ecology

The lacertid genus *Anatololacerta* comprises five species according to the latest phylogenetic review of the taxon (Karakasi et al. 2021). Three of them occur both in Turkey and in some southeastern Greek islands: *A. anatolica* (Werner, 1900) in Samos and Ikaria, *A. finikensis* (Eiselt & Schmidtler, 1986) in Psomi islet and *A. pelasgiana* (Mertens, 1959) in Kastelorizo, Rhodes, Symi and surrounding islets (Fig. 1; Karakasi et al. 2021; Uetz et al. 2021). The Pelasgian rock lizard is a small bodied (SVL up to 65 mm) diurnal species that feeds on invertebrates, mainly insects (Valakos et al. 2008). The species prefers rocky areas, cultivated lands with dry-stone walls, and light deciduous forests, while it also frequents small human settlements and ruins where it can be seen climbing on the walls (Lymberakis et al. 2018).

On 27 November 2020, we visited the western outskirts of Athens for a herpetological survey. The broader area, known collectively as Elaionas (*Ελαιώνας*, literally olive grove), the historical part of Athens where olive trees were cultivated since the antiquity is nowadays a degraded district hosting small industries and logistics companies together with abandoned buildings and uncultivated lands (Metaxas et al. 2007). At approximately 14:15 (air temperature around 18 °C), we observed a single lacertid lizard running and hiding under some garbage at the side of a stream (37.974649°N, 23.688932°E; 18 m a.s.l.). After a while, the lizard emerged from its hiding place and was photographed. We did not encounter any other individuals and thus we returned to the spot a week later (6 December 2020). This time we found eight lizards and captured one




Figure 1. Known distribution of *Anatololacerta pelasgiana* (species range in gray). An arrow points to the new record in western Athens.

adult female (Fig. 2; SVL: 62 mm, tail length: 149 mm, body weight: 4.9 g). One month later (6 January 2021) we visited the spot again and counted 23 individuals (10 males, eight females, five juveniles) within a distance of about 300 meters along the stream. The presence of juveniles indicates that lizards do reproduce in this locality.

Based on the morphological characters and the coloration pattern of the captured individual and the ones we observed in the spot, we concluded that the lizards belonged to the genus *Anatololacerta*. To identify the species, we removed a tail tip (10 mm) from the captured female and sent it to the Molecular Systematics Lab of the Natural History Museum of Crete (NHMC) of the University of Crete. The captured specimen was deposited in the Herpetological Collection of the Zoological Museum of the National and Kapodistrian University of Athens (ZMUA 8624).

Total genomic DNA was extracted from the above specimen using a standard ammonium acetate protocol (Bruford et al. 1998). A fragment (~ 430 bp) of the mitochondrial gene (mtDNA) encoding for cytochrome *b* (*cyt b*), which is one of the most commonly amplified gene markers in the Lacertidae, was amplified through PCR using the primers GLUDG and CB2 (Palumbi 1996), following the conditions described in Karakasi et al. (2021). Double stranded sequencing was carried out using the Big-Dye Terminator v.3.1 Cycle Sequencing kit

on an ABI3730 automated sequencer (CEMIA, Larissa, Greece), following the manufacturer's protocol and using the same primers as in PCR. Sequences were edited using CodonCode Aligner v.9.0.1 (CodonCode Corporation). The identity and authenticity of the produced sequence was evaluated with a BLAST search in the NCBI genetic database (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>). This search revealed that the produced *cyt b* sequence had high similarity with other available *cyt b* sequences of *Anatololacerta pelasgiana* in GenBank (E-value = 0).

To confirm the above results, we carried out a phylogenetic analysis using the data from Karakasi et al. (2021), which is the most recent and complete dataset including all species of *Anatololacerta* from the eastern Mediterranean region (for more details see Table 1). The alignment of the sequences was performed using the ClustalW implemented in MEGAX v.10.2.2 (Kumar et al. 2018), whereas the best model of nucleotide substitution was selected using the PartitionFinder (PF) v.2.1 (Guindon et al. 2010; Lanfear et al. 2012; Lanfear et al. 2016). The pairwise distances (p-distances) were estimated in MEGAX. Bayesian Inference (BI) was performed in MrBayes v.3.2.7 (Ronquist et al. 2012), with four runs and eight chains for each run. Each chain ran for 10^7 generations sampling every 10^3 generations. Several MCMC convergence diagnostics were used to check for convergence and stationarity following the manual's instructions. The



Figure 2. The captured individual, in situ photography (A) and in hand (B).

first 25% trees were discarded as burn-in, as a measure to sample from the stationary distribution and avoid the possibility of including random, sub-optimal trees. A majority rule consensus tree was then produced from the posterior distribution of trees, and the posterior probabilities were calculated as the percentage of samples recovering any particular clade. Posterior probabilities ≥ 0.95 indicate statistically significant support (Huelsenbeck and Ronquist 2001).

In total, 408 base pairs (bp) of *cyt b* sequence were obtained from the examined specimen collected in Athens (Accession number in GenBank: [OP831897](https://www.ncbi.nlm.nih.gov/nuclseq/OP831897)). Pairwise genetic distances (p-distance) between the specimen from Athens and all the others varied from 0 (from *A. pelasgiana* from the island of Rhodes) to 9.8% (from *A. ibrahimi* from Turkey), but were above 14.6% when compared with the outgroup taxa (*Phoenicolacerta*, *Iberolacerta*, *Lacerta*, *Parvilacerta*, and *Hellenolacerta*). The best-fit nucleotide substitution model selected by PF was HKY+I+G. In BI (harmonic mean $-\ln L = -4844.13$), the MCMC convergence diagnostics did not provide any clues of non-convergence and indicated stationarity. Considering the *Anatololacerta* specimen is from Athens, it forms a highly supported clade [posterior probability (p.p.) = 1.00] with *A. pelasgiana* from Dodekanisa (Rhodes, Symi) and Turkey, showing closer proximity with the specimens of *A. pelasgiana* from Rhodes (Fig. 3).

The new Athenian home suits the Pelasgian lizard well. The stream along the banks of which we found the small population is a degraded torrent stream's bed bounded between two stone-built walls, approximately 2.5 meters high, comprising plenty of shelters and basking sites (Fig. 4). The stream is flowing most of the year, attracting insects and thus providing invertebrate food to the lizards.

The vegetation is relatively sparse, with dominant species the Jerusalem thorn (*Parkinsonia aculeata*), black poplar (*Populus nigra*), castor bean (*Ricinus communis*), tree of heaven (*Ailanthus altissima*) and fig tree (*Ficus carica*). Though *A. pelasgiana* has been reported twice in the past to expand its range on other islands such as Kasos (Kornilios and Thanou 2016) and Kastelorizo (Kalaentzis et al. 2018), this is the first time it has settled in a mainland site within a large city.

Though we don't know the exact origin of the new population (based on the Bayesian Inference tree we hypothesize that it comes from Rhodes), we presume that the Pelasgian lizards arrived in Athens through one of the many transport and logistics companies that are located in the area, receiving and shipping goods to and from destinations all over the country. Reptilian unintentional human-mediated transportation is a well-known avenue of new introductions (Krysko and MacKenzie-Krysko 2016; Santos et al. 2018; Medina et al. 2019; Bisbal-Chinesta et al. 2020; Oskyrko et al. 2020), a case particularly common in large metropolitan centers (Tiatragul et al. 2020; Vaughn et al. 2021). Settling in Athens makes *A. pelasgiana* the eighth lizard species reported to have established population in the Greek capital city during the last seven years (*Podarcis siculus* – Adamopoulou 2015; *Chamaeleo chamaeleon* – Dimaki et al. 2015; *Podarcis muralis* – Karameta and Pafilis 2017; *Podarcis peloponnesiacus* – Hedman et al. 2017; *Podarcis vaucheri* – Spilani et al. 2018; *Tarentola mauritanica* – Strachinis and Pafilis 2018; *Algyroides nigropunctatus* – Deimezis-Tsikoutas et al. 2020). The adaptations in urban life that these new populations will adopt and their possible future interactions is a fascinating topic inviting further research (Sol et al. 2013; Johnson and Munshi-South 2017).

Table 1. List of specimens examined in the present study with their corresponding taxon names, voucher numbers, country/region/locality names (detailed only where available), reference of the study in which they were previously used (if any), coordinates given in decimal degrees and accession numbers in GenBank. Clade assignment based on Karakasi et al. (2021).

Code	Clade	Species	Locality	Coordinates (Longitude, Latitude)	cyt b	Reference
1	Clade B	<i>A. pelasgiana</i>	Athens, Greece	23.68893, 37.97464	OP831897	Present study
2	Clade A	<i>A. anatolica</i>	Turkey: Kaz Dag: Gure	26.88096, 39.62310	LN611203	Bellati et al. (2015)
3	Clade A	<i>A. anatolica</i>	Turkey: Kaz Dag: Gure	26.88096, 39.62310	LN611204	Bellati et al. (2015)
4	Clade A	<i>A. anatolica</i>	Turkey: Izmir: Karagol	27.21653, 38.55811	MW087597	Karakasi et al. (2021)
5	Clade A	<i>A. anatolica</i>	Turkey: Aydin: Pasayaylasi	27.88985, 37.93787	LN611207	Bellati et al. (2015)
6	Clade A	<i>A. anatolica</i>	Turkey: Aydin: Pasayaylasi	27.89219, 37.94394	MW087630	Karakasi et al. (2021)
7	Clade A	<i>A. anatolica</i>	Turkey: Aydin: Pasayaylasi	27.89219, 37.94394	MW087631	Karakasi et al. (2021)
8	Clade A	<i>A. anatolica</i>	Greece: Samos	26.89663, 37.73897	LN611212	Bellati et al. (2015)
9	Clade A	<i>A. anatolica</i>	Greece: Samos	26.66754, 37.78866	LN611210	Bellati et al. (2015)
10	Clade A	<i>A. anatolica</i>	Greece: Ikaria isl.	26.06290, 37.61707	LN611214	Bellati et al. (2015)
11	Clade A	<i>A. anatolica</i>	Greece: Ikaria isl.	26.05140, 37.56600	MW087496	Karakasi et al. (2021)
12	Clade B	<i>A. pelasgiana</i>	Greece: Symi isl.	27.84900, 36.60460	MW087492	Karakasi et al. (2021)
13	Clade B	<i>A. pelasgiana</i>	Greece: Symi isl.	27.84900, 36.60460	MW087493	Karakasi et al. (2021)
14	Clade B	<i>A. pelasgiana</i>	Greece: Symi isl.	27.84900, 36.60460	MW087494	Karakasi et al. (2021)
15	Clade B	<i>A. pelasgiana</i>	Greece: Symi isl.	27.83000, 36.62000	LN611224	Bellati et al. (2015)
16	Clade B	<i>A. pelasgiana</i>	Greece: Rodos isl.	27.94270, 36.28960	MW087487	Karakasi et al. (2021)
17	Clade B	<i>A. pelasgiana</i>	Greece: Rodos isl.	28.22115, 36.44388	MW087489	Karakasi et al. (2021)
18	Clade B	<i>A. pelasgiana</i>	Greece: Rodos isl.	28.22115, 36.44388	MW087490	Karakasi et al. (2021)
19	Clade B	<i>A. pelasgiana</i>	Greece: Rodos isl.	28.21767, 36.43549	GQ142137	Pavlicev and Mayer (2009)
20	Clade B	<i>A. pelasgiana</i>	Greece: Rodos isl.	28.21000, 36.33000	LN611220	Bellati et al. (2015)
21	Clade B	<i>A. pelasgiana</i>	Greece: Rodos isl.	28.21000, 36.33000	LN611221	Bellati et al. (2015)
22	Clade B	<i>A. pelasgiana</i>	Greece: Rodos isl.	28.21000, 36.33000	LN611222	Bellati et al. (2015)
23	Clade B	<i>A. pelasgiana</i>	Turkey: Antalya: Elmali	29.80464, 36.52431	MW087533	Karakasi et al. (2021)
24	Clade B	<i>A. pelasgiana</i>	Turkey: Antalya: Korkuteli	30.02850, 37.13719	MW087609	Karakasi et al. (2021)
25	Clade B	<i>A. pelasgiana</i>	Turkey: Denizli: Tavas	29.11686, 37.59461	MW087648	Karakasi et al. (2021)
26	Clade B	<i>A. pelasgiana</i>	Turkey: Denizli: Tavas	29.11686, 37.59461	MW087649	Karakasi et al. (2021)
27	Clade C	<i>A. finikensis</i>	Greece: Kastellorizo: Psomi	29.63720, 36.11530	MW087500	Karakasi et al. (2021)
28	Clade C	<i>A. finikensis</i>	Turkey: Antalya: near Sarilar	29.76800, 36.22376	LN611230	Bellati et al. (2015)
29	Clade C	<i>A. finikensis</i>	Turkey: Antalya: Altinyaka	30.40508, 36.68172	MW087646	Karakasi et al. (2021)
30	Clade C	<i>A. finikensis</i>	Turkey: Antalya: Karaman	30.16038, 36.94415	LN611228	Bellati et al. (2015)
31	Clade D	<i>A. ibrahimi</i>	Turkey: Burdur: Yakakoy	30.34522, 37.69956	MW087524	Karakasi et al. (2021)
32	Clade D	<i>A. ibrahimi</i>	Turkey: Antalya: Manavgat	31.54650, 36.88653	MW087586	Karakasi et al. (2021)
33	Clade D	<i>A. ibrahimi</i>	Turkey: Antalya: Manavgat	31.55906, 36.86689	MW087624	Karakasi et al. (2021)
34	Clade D	<i>A. ibrahimi</i>	Turkey: Antalya: Gazipasa	32.45828, 36.49711	MW087572	Karakasi et al. (2021)
35	Clade D	<i>A. ibrahimi</i>	Turkey: Mersin: Anamur	32.83011, 36.21247	MW087481	Karakasi et al. (2021)
36	Clade D	<i>A. ibrahimi</i>	Turkey: Icel: Abanoz	32.95107, 36.32493	LN611237	Bellati et al. (2015)
37	Clade D	<i>A. ibrahimi</i>	Turkey: Mersin: Gulnar	33.37964, 36.27850	MW087583	Karakasi et al. (2021)
38	Clade D	<i>A. ibrahimi</i>	Turkey: Mersin: Gulnar	33.37964, 36.27850	MW087585	Karakasi et al. (2021)
39	Clade E	<i>A. danfordi</i>	Turkey: Icel: Kavaklipinar	34.35123, 37.01912	LN611241	Bellati et al. (2015)
40	Clade E	<i>A. danfordi</i>	Turkey: Mersin: Camliyayla	34.57811, 37.14658	MW087638	Karakasi et al. (2021)
41	Clade E	<i>A. danfordi</i>	Turkey: Mersin: Camliyayla	34.57811, 37.14658	MW087640	Karakasi et al. (2021)
42	Clade E	<i>A. danfordi</i>	Turkey: Adana: Kozan	35.86014, 37.50894	MW087613	Karakasi et al. (2021)
43	Clade E	<i>A. danfordi</i>	Turkey: Adana: Feke	35.95111, 37.82644	MW087552	Karakasi et al. (2021)
44	Clade E	<i>A. danfordi</i>	Turkey: Adana: Feke	35.95111, 37.82644	MW087555	Karakasi et al. (2021)
45	Clade E	<i>A. danfordi</i>	Turkey: Adana: Saimbeyli	36.09392, 38.00275	MW087636	Karakasi et al. (2021)
46	Outgroup	<i>Iberolacerta aranica</i>	France: Serre de Ventailou	–	AY151876	Carranza et al. (2004)
47	Outgroup	<i>Iberolacerta horvathi</i>	Croatia: Northwest	–	AY256648	Carranza et al. (2004)
48	Outgroup	<i>Hellenolacerta graeca</i> 1	Greece: Mystras	22.37201, 37.07046	LN611249	Bellati et al. (2014)
49	Outgroup	<i>Hellenolacerta graeca</i> 2	Greece: Serveika	22.31710, 37.15508	LN611250	Bellati et al. (2014)
50	Outgroup	<i>Phoenicolacerta troodica</i> 1	Cyprus: Kakopetria	32.91028, 34.97667	MW087633	Karakasi et al. (2021)
51	Outgroup	<i>Phoenicolacerta troodica</i> 2	Cyprus: Kakopetria	32.91028, 34.97667	MW087634	Karakasi et al. (2021)
52	Outgroup	<i>Lacerta agilis</i>	Greece: Makedonia: Florina	–	KJ940308	Sagonas et al. (2014)
53	Outgroup	<i>Parvilacerta parva</i>	Turkey: Sivas: Osmandede	37.02020, 38.74674	LN611248	Bellati et al. (2014)

References

- Adamopoulou C (2015) First record of *Podarcis siculus* (Rafinesque-Schmaltz, 1810) from Greece. *Herpetozoa* 27(3/4): 187–188.
- Bellati A, Carranza S, Garcia-Porta J, Fasola M, Sindaco R (2015) Cryptic diversity within the *Anatololacerta* species complex (Squamata: Lacertidae) in the Anatolian Peninsula: evidence from a multi-locus approach. *Molecular Phylogenetics and Evolution* 82: 219–233. <https://doi.org/10.1016/j.ympev.2014.10.003>
- Bisbal-Chinesta JF, Tamar K, Gálvez Á, Alberó L, Vicent-Castelló P, Martín-Burgos L, Alonso M, Sánchez R, Ortega C, Gómez A, Candel D, Cervera M, Carranza S, Blain HA (2020) Trade and stowaways: molecular evidence for human-mediated translocation of eastern skinks into the western Mediterranean. *Amphibia-Reptilia* 41(1): 49–62. <https://doi.org/10.1163/15685381-20191249>
- Bruford MW, Hanotte O, Burke T (1998) Multi and single locus DNA fingerprinting. In: Hoelzel AR (Ed.) *Molecular genetic analysis of populations: a practical approach*. IRL Press, 225–269.
- Carranza S, Arnold EN, Amat F (2004) DNA phylogeny of *Lacerta* (*Iberolacerta*) and other lacertine lizards (Reptilia: Lacertidae): did

- competition cause long-term mountain restriction? *Systematics and Biodiversity* 2: 57–77. <https://doi.org/10.1017/S1477200004001355>
- Deimezis-Tsikoutas A, Kapsalas G, Antonopoulos A, Strachinis I, Pafilis P (2020) *Algyroides nigropunctatus* (Squamata: Lacertidae) in the city of Athens: An unexpected finding. *Russian Journal of Herpetology* 27(3): 172–174. <https://doi.org/10.30906/1026-2296-2020-27-3-172-174>
- Dimaki M, Chondropoulos B, Legakis A, Valakos E, Vergetopoulos M (2015) New data on the distribution and population density of the African Chameleon, *Chamaeleo africanus* and the Common Chameleon, *Chamaeleo chamaeleon* in Greece. *Hyla* 2015(1): 36–43.
- Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O (2010) New Algorithms and Methods to Estimate Maximum-Likelihood Phylogenies: Assessing the Performance of PhyML 3.0. *Systematic biology* 59: 307–321. <https://doi.org/10.1093/sysbio/syq010>
- Hedman H, Kapsalas G, Karameta E, Psonis N, Poulakakis N, Foufopoulos J, Pafilis P (2017) First record of *Podarcis peloponnesiacus* (Bibron et Bory, 1833) from outside the Peloponnese. *Herpetozoa* 29(3/4): 190–193.
- Huelsenbeck JP, Ronquist F (2001) MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics* 17: 754–755. <https://doi.org/10.1093/bioinformatics/17.8.754>
- Johnson MTJ, Munshi-South J (2017) Evolution of life in urban environments. *Science* 358: eaam8327. <https://doi.org/10.1126/science.aam8327>
- Kalaentzis K, Strachinis I, Katsiyiannis P, Oefinger P, Kazilas C (2018) New records and an updated list of the herpetofauna of Kastellorizo and the adjacent islet Psomi (Dodecanese, Greece). *Herpetology Notes* 11: 1009–1019.
- Karakasi D, Ilgaz Ç, Kumlutaş Y, Candan K, Güçlü Ö, Kankılıç T, Beşer N, Sindaco R, Lymberakis P, Poulakakis N (2021) More evidence of cryptic diversity in *Anatololacerta* species complex Arnold, Arribas & Carranza, 2007 (Squamata: Lacertidae) and re-evaluation of its current taxonomy. *Amphibia-Reptilia* 42(2): 201–216. <https://doi.org/10.1163/15685381-bja10045>
- Karameta E, Pafilis P (2017) First record of *Podarcis muralis* (Laurenti, 1768) from Athens, Greece. *Herpetozoa* 30: 87–88.
- Kornilios P, Thanou E (2016) Two additions to the herpetofauna of Kasos (Aegean Sea, Greece) and the role of human-mediated dispersals. *Herpetological Review* 47(4): 633–635.
- Krysko KL, MacKenzie-Krysko C (2016) First report of the mourning gecko, *Lepidodactylus lugubris* (Duméril & Bibron, 1836), from The Bahamas. *Caribbean Herpetology* 54: 1–2. <https://doi.org/10.31611/CH.54>
- Kumar S, Stecher G, Li M, Knyaz C, Tamura K (2018) MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. *Molecular Biology and Evolution* 35(6): 1547–1549. <https://doi.org/10.1093/molbev/msy096>
- Lanfear R, Calcott B, Ho SYW, Guindon S (2012) PartitionFinder: Combined selection of partitioning schemes and substitution models for phylogenetic analyses. *Molecular Biology and Evolution* 29: 1695–1701. <https://doi.org/10.1093/molbev/mss020>
- Lanfear R, Frandsen PB, Wright AM, Senfeld T, Calcott B (2016) PartitionFinder 2: New Methods for Selecting Partitioned Models of Evolution for Molecular and Morphological Phylogenetic Analyses. *Molecular biology and evolution* 34(3): 772–773. <https://doi.org/10.1093/molbev/msw260>
- Lymberakis P, Pafilis P, Poulakakis N, Sotiropoulos K, Valakos ED (2018) Amphibians and Reptiles of the Aegean Sea. In: Sfenthourakis S, Pafilis P, Parmakelis A, Poulakakis N, Triantis KA (Eds) *Biogeography and Biodiversity of the Aegean*. Broken Hill Publications, 169–190.
- Medina JAO, Cen DIC, Noh MMC, Vázquez JRC (2019) First record of the Moorish Gecko, *Tarentola mauritanica* (Linnaeus, 1758) (Squamata: Phyllodactylidae), in Mexico. *Herpetology Notes* 12: 971–974.
- Metaxas G, Sfakianaki E, Spyropoulos N (2007) Using Geographical Information Systems in the environmental assessment of urban environments: the Elaionas area in Attica, Greece. *World Transactions on Engineering and Technology Education* 7: 127–133.
- Oskyrko O, Laakkonen H, Silva-Rocha I, Jablonski D, Marushchak O, Uller T, Carretero MA (2020) The possible origin of the common wall lizard, *Podarcis muralis* (Laurenti, 1768) in Ukraine. *Herpetozoa* 33: 87–93. <https://doi.org/10.3897/herpetozoa.33.e49683>
- Palumbi SR (1996) *Nucleic Acids II: The Polymerase Chain Reaction*. In: Hillis DM, Moritz C, Mable BK (Eds) *Molecular Systematics*. Sinauer, Sunderland, MA, 205–248.
- Pavlicev M, Mayer W (2009) Fast radiation of the subfamily Lacertinae (Reptilia: Lacertidae): History or methodical artefact? *Molecular Phylogenetics and Evolution* 52: 727–734. <https://doi.org/10.1016/j.ympev.2009.04.020>
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61: 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Sagonas K, Poulakakis N, Lymberakis P, Parmakelis A, Pafilis P, Valakos ED (2014) Molecular systematics and historical biogeography of the green lizards (*Lacerta*) in Greece: Insights from mitochondrial and nuclear DNA. *Molecular Phylogenetics and Evolution* 76: 144–154. <https://doi.org/10.1016/j.ympev.2014.03.013>
- Santos JL, Žagar A, Drašler K, Rato C, Ayres C, Harris DJ, Carretero MA, Salvi D (2018) Phylogeographic evidence for multiple long-distance introductions of the common wall lizard associated with human trade and transport. *Amphibia-Reptilia* 40(1): 121–127. <https://doi.org/10.1163/15685381-20181040>
- Sol D, Lapedra O, González-Lagos C (2013) Behavioural adjustments for a life in the city. *Animal Behaviour* 85: 1101–1112. <https://doi.org/10.1016/j.anbehav.2013.01.023>
- Spilani L, Strachinis I, Lampropoulos A, Tsigas P, Poulakakis N, Pafilis P (2018) *Podarcis vaucheri* (Sauria: Lacertidae) far away from home: a new invasive species in Greece. *Amphibia-Reptilia* 39(3): 363–368. <https://doi.org/10.1163/15685381-18000002>
- Strachinis I, Pafilis P (2018) First record of *Tarentola mauritanica* (Linnaeus, 1758) from Athens, Greece. *Herpetozoa* 31(1/2): 98–99.
- Tiatragul S, Hall JM, Warner DA (2020) Nestled in the city heat: urban nesting behavior enhances embryo development of an invasive lizard. *Journal of Urban Ecology* 6: 1. <https://doi.org/10.1093/jue/juaa001>
- Uetz P, Freed P, Aguilar R, Hošek J (Eds) (2021) *The Reptile Database*. <http://www.reptile-database.org> [accessed November 10, 2021]
- Valakos ED, Pafilis P, Lymberakis P, Maragou P, Sotiropoulos K, Foufopoulos J (2008) *The amphibians and reptiles of Greece*. Edition Chimaira, Frankfurt, 463 pp.
- Vaughn PL, McQueen W, Gangloff EJ (2021) Moving to the city: testing the implications of morphological shifts on locomotor performance in introduced urban lizards. *Biological Journal of the Linnean Society* 134: 141–153. <https://doi.org/10.1093/biolinnean/blab076>

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Herpetozoa](#)

Jahr/Year: 2022

Band/Volume: [35](#)

Autor(en)/Author(s): Christopoulos Apostolos, Pantagaki Charikleia-Foteini,
Poulakakis Nikos, Pafilis Panayiotis

Artikel/Article: [First record of *Anatololacerta pelasgiana* \(Mertens, 1959\) in mainland Greece: another new species in Athens 239-244](#)