

Melanism in the grass snake *Natrix natrix* (Linnaeus, 1758) from the Danube Delta Biosphere Reserve, Romania

Geanina Fănar^{1,2,3*}, Alexandra E. Telea^{1,2,3}, Iulian Gherghel^{1,4}, Raluca Melenciuc^{5*}

¹ Faculty of Natural and Agricultural Sciences, Ovidius University of Constanța, Constanța, Romania

² Association Chelonia Romania, Bucharest, Romania

³ Emil Racoviță Institute of Speleology of Romanian Academy of Sciences, Bucharest, Romania

⁴ Department of Exact Sciences and Natural Sciences, Institute of Interdisciplinary Research, Alexandru Ioan Cuza University of Iași, Iași, Romania

⁵ Faculty of Biology, Alexandru Ioan Cuza University, Iași, Romania

<https://zoobank.org/CD5A060B-94D3-49D1-81CA-C67D51C07B04>

Corresponding author: Alexandra E. Telea (alexandra.telea@gmail.com)

Academic editor: Silke Schweiger ♦ Received 19 April 2022 ♦ Accepted 7 June 2022 ♦ Published 21 December 2022

Abstract

Animal colouration has a significant ecological role in defence, reproduction, and thermoregulation. In the case of melanism, it is a complex topic. Besides potential disadvantages such as higher risk of predation, melanistic ectotherms may have certain physiological advantages such as more efficient thermoregulation in colder climates and thus, reduced basking time. The common grass snake (*Natrix natrix*) is a widespread species throughout Europe and Asia. It exhibits a wide range of colour polymorphisms, from olive to dark grey, even albinistic and melanistic. Between 2016 and 2021, we conducted fieldwork in the Danube Delta Biosphere Reserve (DDBR) with the aim to document the geographic range of melanistic grass snakes. We categorised the melanistic expressions of *N. natrix* individuals as melanotic, completely melanistic, and partially melanistic. Melanistic snakes were encountered in all six localities visited, suggesting that the occurrence of melanistic grass snakes in the DDBR is geographically widespread. We observed both juveniles (n=2) and adults (n=11) with melanism, suggesting that individuals are born melanistic. However, the proportion of melanistic individuals in the general population of *N. natrix* from the DDBR is unknown. Only at Histria locality we studied the proportion of melanism in the grass snake population and 6.3% of the snakes caught were melanistic. Body size comparisons are not statistically significant because of the low sample size. The *N. natrix* melanistic morph's geographical distribution in the DDBR is most likely due to an interaction of climate and habitats, which offer a thermal advantage in the face of predation pressure.

Key Words

colour morphs, ectotherms, melanotic, polymorphism

Introduction

Animal colouration provides ecological roles to organisms such as the capacity to warn predators, increase crypsis, signal potential mating partners, and is subject to strong selective pressures (Clusella-Trullas et al.

2007). In ectothermic vertebrates the intra- and inter-specific variation in colour provides thermoregulatory opportunities to exploit niches that maximise their thermal performance optimums (Clusella-Trullas et al. 2007; Geen and Johnston 2014). For instance, darker colour is supposedly favoured in cooler climates since

* These authors shared first authorship.

dark individuals (i.e., lower skin reflectance) have an advantage under low temperature conditions, as they heat up faster than lighter individuals (Clusella-Trullas et al. 2007; Hodges 2018). Melanism occurs when the quantity of skin melanin pigment is found in large amounts at the expense of other colours (Majerus 1998; van Grouw 2017; Măranducă et al. 2019). This can lead to animal populations with complete melanistic individuals that display an entirely black colour phenotype or variants such as melanotic (nearly entirely expressed melanism) or pseudo-/partial- melanism (Majerus 1998; Zuffi 2008; Domeneghetti et al. 2016). Various melanin concentrations in animal populations led to varying degrees of darker pigmentation leading to continuous intraspecific polymorphism (True 2003). The drivers of melanistic polymorphism are still subject to debate and research (Majerus 1998; Zuffi 2008; Domeneghetti et al. 2016). Laboratory experiments and field studies on ectotherms showed that melanistic morphs use different thermoregulation strategies to morphs with less melanin, such as different use of habitat and lower basking time. Sampling of *Vipera aspis* in the Swiss Alps had indicated that melanistic individuals prefer cooler microhabitats with higher vegetation cover, compared to non-melanistic ones (Muri et al. 2015). Thus, melanistic morphs preserve their body temperature better in colder weather and through physiological and behavioural adaptations are protected against overheating in warm weather conditions (Gibson and Falls 1979; Clusella-Trullas et al. 2007; Tanaka 2007; Muri et al. 2015). Studies on melanistic snakes showed that they are generally smaller than non-melanistic individuals from the same population and prone to increased predation, which could indicate that melanistic individuals could potentially have minimal thermal benefits compared to non-melanistic individuals (Madsen 1987; Bury et al. 2020).

Melanism is generally not rare in certain snakes, especially in colder environments (Clusella-Trullas et al. 2007; Lorient et al. 2008; Zuffi 2008). Across Europe there have been recorded melanistic individuals in different snake species such as *Zamenis longissimus* (Zadravec and Lauš 2011), *Coronella austriaca* (Nash et al. 2016), *Vipera berus* (Terhivuo 1990; Strugariu et al. 2009; Nash et al. 2016). Melanistic individuals have also been recorded in the semi-aquatic *Natrix* species (Baran 1976; Naumov 2007; Ajtić et al. 2013; Gvozdenović and Schweiger 2014; Jablonski and Kautman 2017).

The grass snake *N. natrix* is a species known for its polymorphism, with a wide range of colours and morphs which have been reported within its range. Dorsal pattern can range from typical olive-green, brown or greyish to individuals with rows of black spots or bars, or double yellow stripes (Fuhn and Vancea 1961; Thorpe 1979; Kabisch 1999; Speybroeck et al. 2016).

We still have a limited understanding of the distribution and frequency of the melanistic morph in *N. natrix* populations throughout its range. Reports of melanistic

individuals of grass snakes have been made from Turkey (Baran 1976; Habiboğlu et al. 2016; Yenmiş et al. 2022), Bulgaria (Naumov and Tomović 2005; Stojanov et al. 2011; Mollov 2012), Serbia (Bjelica and Anđelković 2021), Croatia (Zadravec and Lauš 2011), Montenegro (Gvozdenović and Schweiger 2014), Bosnia and Herzegovina (Bašić and Zimić 2016), Slovakia (Jandzik 2004), Poland (Błażuk 2007; Kolenda et al. 2017; Bury et al. 2020). In Romania, there are only three general literature mentions of melanistic *N. natrix*, all of them from the Danube Delta Biosphere Reserve (DDBR) (Fuhn and Vancea 1961; Halpern et al. 2002; Zamfirescu et al. 2010).

The aim of our paper is to document the geographic range of melanistic individuals of grass snakes (*N. natrix*) from the DDBR. We also discuss possible advantages and disadvantages for the occurrence of melanism in the study region.

Methods

Study area

The Danube Delta is composed of a network of channels, lagoons, marshes, levees, and three main Danube river lobes (Chilia to the north, Sulina in the middle and Sfântul Gheorghe to the south) and the Razelm-Sinoe lagoon complex (Vespremeanu-Stroe et al. 2017). The DDBR has an average altitude of 0.52 m above mean Black Sea level, with the highest point at 12.4 m altitude, and covers an area of 5800 km² and supports rich biodiversity, with numerous populations of bird and reptile species (Găstescu 2009; Zamfirescu et al. 2010; Marinov et al. 2019). According to the Köppen-Geiger classification system, the region is classified as a C climate, warm temperate, fully humid with hot summers (Kottek et al. 2006). The DDBR is influenced by the Black Sea and is characterised by torrid summers, very cold winters and is considered the windiest and sunniest region of Romania (Ciulache and Torică 2003; Găstescu 2009). The habitats are diverse, ranging from anthropic settlements, agricultural fields and pastures to sand dunes with reeds and a few areas with forests, in close proximity to water bodies (Doroftei 2013). 12 reptile species have been recorded here (*Emys orbicularis*, *Testudo graeca*, *Eremias arguta*, *Lacerta agilis*, *Lacerta viridis*, *Podarcis tauricus*, *Coronella austriaca*, *Elaphe sauromates*, *Dolichophis caspius*, *Natrix natrix*, *Natrix tessellata*, *Vipera ursinii*); of these, *N. natrix* is the most common species due to the suitable habitats (Zamfirescu et al. 2010; Cogălniceanu et al. 2013).

Data collection and analysis

We conducted opportunistic and transect surveys of melanistic individuals in the areas accessible by foot, between 8:00 – 14:00h, during the snake's active season

(March–October), from 2016 to 2021 throughout the DDBR. Melanistic grass snakes were visually searched for and caught by hand when possible. Upon capture, we measured snake's body mass (BM) using a spring / digital scale (0.1 g accuracy) and photographed alongside a graph paper, using DSLR cameras with standard lenses (Nikon D7100 and Olympus OM-D E-M1 MarkII). The snakes' snout-vent length (SVL) was then measured in ImageJ (Rasband 1997). The location of each melanistic individual was recorded using Orux Maps and Geo Tracker applications for mobile phone, using geographic coordinates in datum WGS84. At one 32 ha locality, the Histria Archaeological Complex (Fig. 1), the methodology described above was applied to both melanistic and non-melanistic grass snakes.

Depending on the morphological expression of melanistic characteristics, we categorised the melanistic morphs of grass snakes following Zuffi (2008) as:

- completely melanistic: complete melanism expressed, with a few ventral white scales still retained (Fig. 3C);
- melanotic: where melanism is nearly entirely expressed, with a few morphological characteristics which break the melanistic expression, in grass snakes – a few dorsal scales coloured with dark brown pigment (Figs 2, 3B);
- partially melanistic: where the normal colour pattern has an increased black pigmentation (Fig. 3A).

For the statistical analysis between groups of melanistic and non-melanistic grass snakes from Histria locality we have applied a One-Way Anova test.




Figure 1. Danube Delta Biosphere Reserve and the locations of the melanistic grass snakes *Natrix natrix* found during our study and in literature (map source: [OpenStreetMap.org](https://www.openstreetmap.org) and [geo-spatial.org](https://www.geo-spatial.org)).

Results

We found 13 melanistic individuals in close proximity to six localities: Letea, Maliuc, Sulina, Sfantu Gheorghe, Periboina, and Histria (Fig. 1). All melanistic snakes were found in habitats characterised by sand dunes with reed vegetation, surrounded by water bodies such as lakes, canals or the coastal area. Four were adult females (SVL between 580 – 870 mm), six were adult males (SVL between 500 – 882 mm) and two were juveniles (SVL 185–198 mm) (Table 1). One adult male from Periboina locality lacks morphometrics data because it was found opportunistically by colleagues. An adult found in Sulina locality, with unknown sex was found dead on the road in a state that precluded measuring.

Table 1. Characteristics of the melanistic individuals of grass snakes *Natrix natrix* found in DDBR between 2016 and 2021. Abbreviations as follows: Localities L-Letea, M-Maliuc, S-Sulina, SG-Sfantu Gheorghe, H-Histria, P-Periboina (Fig. 1); Melanism CM-completely melanistic, PM-partially melanistic, MI-melanotic; Age A-Adult, J-Juvenile; *Dead on road.

Locality	Date	Sex	Age	SVL (mm)	Weight (g)	Coordinates	Melanism	Figure
L	14.05.16	M	A	500	42.9	45°17.16'N, 29°31.44'E	CM	
L	04.06.17	–	J	198	30	45°17.16'N, 29°31.44'E	CM	
M	16.08.17	F	J	185	18.8	45°12.00'N, 29°6.66'E	PM	
S	06.06.17	–	A*	–	–	45°9.36'N, 29°39.24'E	CM	
SG	17.06.17	F	A	870	284.8	44°53.76'N, 29°35.58'E	CM	
SG	22.07.17	M	A	882	191.4	44°53.76'N, 29°35.58'E	CM	
SG	22.07.17	M	A	590	95.9	44°53.76'N, 29°35.58'E	MI	
SG	22.07.17	M	A	641	81.6	44°53.76'N, 29°35.58'E	PM	3A
P	24.04.21	M	A	–	–	44°36.78'N, 28°55.80'E	CM	3C
P	01.05.21	F	A	680	–	44°36.84'N, 28°55.86'E	CM	
H	13.10.18	F	A	625	76	44°32.82'N, 28°46.38'E	MI	
H	17.10.20	M	A	630	82	44°32.82'N, 28°46.50'E	MI	2
H	19.10.21	F	A	580	70	44°32.94'N, 28°45.72'E	MI	3B

We encountered all three categories of melanism expression: 53.8% completely melanistic individuals (n=7), 15.4% partially melanistic (n=2), and 30.8% melanotic (n=4) (Table 1, Figs 2, 3).

At Histria we caught a total of 48 adult grass snakes, of which only three snakes were melanistic (6.3%). The three melanistic grass snakes from Histria had an average SVL of 611.66 mm and BM of 76.00 g. The non-melanistic grass snakes had an average SVL of 609.71 mm, and BM of 76.26 g (Table 2). The difference in body size between the two categories is statistically non-significant (df=1, F=0.00, p>0.05) for both SVL and BM.



Figure 2. Polymorphism observed in *Natrix natrix* populations from DDBR. Top common morph; middle double yellow stripes pattern; bottom melanotic morph (photo credit: E. A. Telea & G. Fănară).

Discussion

Melanistic grass snakes have been found in all six localities surveyed, four of these not mentioned before in literature (Fig. 1). Although *N. natrix* is a widespread species in Romania (Cogălniceanu et al. 2013), previous recorded melanistic were all observed in the DDBR, in four localities (Fuhn and Vancea 1961; Halpern et al. 2002; Zamfirescu et al. 2010). Literature data and our results suggest that melanism in *N. natrix* is geographically widespread in the DDBR.

We have encountered all three melanistic morph categories, as defined after Zuffi (2008), and as much as we could identify them, they have been previously reported in *N. natrix* across its range (Fuhn and Vancea 1961; Naumov and Tomović 2005; Zamfirescu et al. 2010; Zadravec and Lauš 2011; Mollov 2012; Gvozdenović and Schweiger 2014; Habiboğlu et al. 2016; Kolenda et al. 2017; Bjelica and Anđelković 2021; Yenmiş et al. 2022). Both the present study and recorded cases in literature of melanistic grass snakes rely on visual encounter surveys. Fuhn and Vancea (1961) mentioned that the most common melanistic morph of grass snakes in the Romanian DDBR was the melanotic phenotype. However, in our study, the completely melanistic one was more abundant and widespread, found in the majority of localities

visited (Figs 1, 3C, Table 1). In literature, this morph seems to be the most commonly reported melanistic phenotype in grass snakes and has been reported before in Poland (Kolenda et al. 2017), Bulgaria (Naumov and Tomović 2005; Mollov 2012), Turkey (Habiboğlu et al. 2016), Serbia (Bjelica and Anđelković 2021), and in the Romanian DDBR (Zamfirescu et al. 2010). The melanotic phenotype, on the other hand, was the second most often encountered in our study, numerically and geographically (Figs 1, 3B, Table 1). This morph was also recorded in Turkey in a record published by Yenmiş et al. (2022). The third morph, the partially melanistic morph, was the least encountered numerically and in only two localities visited (Figs 1, 3A, Table 1). The partially melanistic grass snakes were previously recorded in grass snakes in Croatia and Montenegro (Zadravec and Lauš 2011; Gvozdenović and Schweiger 2014).

In the small sample size from Histria locality (Fig. 1, Table 2), melanistic and non-melanistic grass snakes did not differ in body length or body size. In a previous study, Bury et al. (2020) showed that the melanistic grass snakes are smaller than the typically coloured individuals of both sexes. The small size of melanistic snakes is counterintuitive, seeing as, generally, a thermoregulatory advantage leads to higher growth rates, larger body size, and greater reproductive success (Bittner and King 2003).




Figure 3. Examples of encountered melanistic morphs of *Natrix natrix*. **A.** dorsal view of partially melanistic morph with double yellow stripes pattern, Sfântu Gheorghe (photo credit: I. Gherghel); **B.** dorsal view of melanotic morph, Histria (photo credit: E.A. Telea); **C.** partial lateral and ventral views of completely melanistic morph, Periboina (photo credit: F. Stănescu).

Forsman (1995) identified no consistent difference in the proportion of time spent basking and maintaining higher temperatures between melanistic and zigzag individuals of *V. berus*. Such a situation could explain our population case: low altitude combined with the DDBR's climate and high-density vegetation habitats could offer some advantages for thermoregulatory behaviour for melanistic grass snakes against predation pressure, but not enough to make a considerable difference in higher growth rate opportunities between melanistics and non-melanistics. But because of the small sample size of grass snakes from this locality, we stress that further studies are needed for a better sample size in order to test differences in size and growth rate between the colour morphs.

Table 2. Body sizes of grass snakes *Natrix natrix* from Histria locality.

	SVL (mm)	BM (g)
	min–max mean±s.e.	min–max mean±s.e.
Melanistic (n = 3)	580–630 611.66±15.89	70–82 76.00±3.46
Non-melanistic (n = 45)	445–785 609.71±11.26	25–169 76.26±4.73

All surveys were done during the day, from morning to early afternoon. All melanistic grass snakes we found were either hidden in reed vegetation or in close proximity to an abundant reed vegetation habitat. This habitat choice might have affected the success of our detection and capture of melanistic grass snakes. However, such behaviour and habitat choice might not have been accidental. Hypothetically, it could have been advantageous to the melanistic grass snakes as they could maintain a thermal benefit in microhabitats with less exposure to the sun (e.g., high-density vegetation), while also avoiding predator (and researcher) detection. From an evolutionary point of view, in snakes, melanism is established when the thermoregulatory advantage outweighs the risks (Clusella-Trullas et al. 2007; Broennimann et al. 2014). Studies have shown that at a population level melanistic snakes are more predated upon; because of their distinctiveness, predators have an easier task of singling out the dark individual (Andrén and Nilson 1981; Madsen 1987). At continental level, darker individuals of animals (i.e., individuals with higher concentration of melanin) tend to live in warm and humid climates (Gloger, 1833 in Millien et al. 2006). However, melanism is thought to be more advantageous in colder environments such as high-altitude habitats,

or even peninsular and coastal habitats characterized by cooler and more humid weather (Millien et al. 2006; Clusella-Trullas et al. 2007; Geen and Johnston 2014). In the case of the DDBR, the region is characterized by a warm, sunny and humid climate, with strong winds from the coastal regions. The DDBR is also known for a very high abundance and diversity of bird species. Although it is a suitable region for many birds, including species which prey upon snakes, such as *Circaetus gallicus*, *Circus* sp. and *Ciconia* sp. (Prejbeanu and Rada 2007; Gâstescu 2009; Marinov et al. 2019), the melanistic grass snake individuals have plenty of dense habitats suitable for hiding from potential predators. The abundant reed vegetation offers protection both against predators and overheating in the DDBR's warm and humid climate, but also protection against the characteristic cool, strong winds for this coastal region (Ciulache and Torică 2003; Kottek et al. 2006). The presence and abundance of *N. natrix* melanistic morph's geographical distribution in Romania in the DDBR is likely due to the interaction of climate and habitat, which offers a thermal advantage in the face of predation pressure.

Acknowledgements

We are grateful to Dan Cogălniceanu for his helpful comments on versions of the manuscript and to our collaborators Florina Stănescu and Alexandra Șoimu for one of the observations presented here. We also thank the editor and the reviewers for their effort and time in helping us improve the manuscript; their commitment was of great value. This work was partially supported by a Romanian Ministry of Education and Research grant CNCS – UEFISCDI, project number PN-III-P1-1.1-TE-2019-1233, and project number PN-III-P1-1.1-PD-2021-0591 within PNCDI III. Research permits were provided by the Danube Delta Biosphere Reserve Administration (05/09.03.2016, 05/16.03.2017, 02/22.02.2018, 13/23.05.2019, 100/24.06.2020 and 13/28.01.2021). Access to the Histria Fortress Archaeological Complex was given by the National History and Archaeology Museum Constanța.

References

- Ajtić R, Tomović L, Sterijovski B, Crnobrnja-Isailović J, Djordjević S, Djurakić M, Golubović A, Simović A, Arsovski D, Andjelković M, Krstić M, Šukalo G, Gvozdenović S, Aïdam A, Michel CL, Ballouard J-M, Bonnet X (2013) Unexpected life history traits in a very dense population of dice snakes. *Zoologischer Anzeiger-A Journal of Comparative Zoology* 252: 350–358. <https://doi.org/10.1016/j.jcz.2012.10.001>
- Andrén C, Nilson G (1981) Reproductive success and risk of predation in normal and melanistic colour morphs of the adder, *Vipera berus*. *Biological Journal of the Linnean Society* 15: 235–246. <https://doi.org/10.1111/j.1095-8312.1981.tb00761.x>
- Baran I (1976) Türkiye yılanlarının taksonomik revizyonu ve coğrafi dağılımları. Tübitak Yayınları, 177 pp.
- Bašić M, Zimić A (2016) Findings of melanistic individuals of Grass Snake (*Natrix natrix* Linnaeus, 1758) in Bosnia and Herzegovina. *Prilozi fauni Bosne i Hercegovine*: 25–29.
- Bittner TD, King RB (2003) Gene flow and melanism in garter snakes revisited: a comparison of molecular markers and island vs. coalescent models. *Biological Journal of the Linnean Society* 79: 389–399. <https://doi.org/10.1046/j.1095-8312.2003.00199.x>
- Bjelica V, Anđelković M (2021) Different shades of snake: Peculiar coloration in an urban population of the Grass Snake, *Natrix natrix* (Linnaeus, 1758). *Ecologica Montenegrina* 39: 81–85. <https://doi.org/10.37828/em.2021.39.9>
- Błażuk J (2007) Herpetofauna of the San River valley along the Otryt mountain ridge and of the adjacent areas (Western Bieszczady Mts.). *Reptiles. Roczniki Bieszczadzkie* 15: 181–229.
- Broennimann O, Ursenbacher S, Meyer A, Golay P, Monney J-C, Schmocker H, Guisan A, Dubey S (2014) Influence of climate on the presence of colour polymorphism in two montane reptile species. *Biology Letters* 10: 20140638. <https://doi.org/10.1098/rsbl.2014.0638>
- Bury S, Mazgajski TD, Najbar B, Zajac B, Kurek K (2020) Melanism, body size, and sex ratio in snakes—new data on the grass snake (*Natrix natrix*) and synthesis. *The Science of Nature* 107: 22. <https://doi.org/10.1007/s00114-020-01678-x>
- Ciulache S, Torică V (2003) Clima Dobrogei. In: Dobrogea I. *Analele Universității București. Geografie. Universitatea din București, Facultatea de Geografie*, 83–106.
- Clusella-Trullas S, van Wyk JH, Spotila JR (2007) Thermal melanism in ectotherms. *Journal of Thermal Biology* 32: 235–245. <https://doi.org/10.1016/j.jtherbio.2007.01.013>
- Cogălniceanu D, Rozyłowicz L, Székely P, Samoilă C, Stănescu F, Tudor M, Székely D, Iosif R (2013) Diversity and distribution of reptiles in Romania. *ZooKeys*: 49–76. <https://doi.org/10.3897/zookeys.341.5502>
- Domeneghetti D, Mondini S, Bruni G (2016) Melanism and pseudo-melanism in the Common Wall Lizard, *Podarcis muralis* Laurenti, 1768 (Reptilia: Lacertidae) in central Italy. *Herpetology Notes* 9: 307–309.
- Doroftei M (2013) Climate change adapted management plan for Danube Delta Biosphere Reserve. Danube Delta National Institute. Climate-change adapted management plan, 135 pp.
- Forsman A (1995) Heating rates and body temperature variation in melanistic and zigzag *Vipera berus*: does colour make a difference? *Annales Zoologici Fennici* 32: 365–374.
- Fuhn IE, Vancea Ș (1961) XIV Fauna Republicii Populare Române. Reptilia. Academiei Republicii Populare Române, Bucharest, Romania, 349 pp.
- Gâstescu P (2009) The Danube Delta biosphere reserve. Geography, biodiversity, protection, management. *Romanian Journal of Geography* 53: 139–152.
- Geen MRS, Johnston GR (2014) Coloration affects heating and cooling in three color morphs of the Australian bluetongue lizard, *Tiliqua scincoides*. *Journal of Thermal Biology* 43: 54–60. <https://doi.org/10.1016/j.jtherbio.2014.04.004>
- Gibson AR, Falls JB (1979) Thermal biology of the common garter snake *Thamnophis sirtalis* (L.): II. The effects of melanism. *Oecologia* 43: 99–109. <https://doi.org/10.1007/BF00346675>

- van Grouw H (2017) The dark side of birds: melanism—facts and fiction. *Bulletin of the British Ornithologists' Club* 137: 12–36. <https://doi.org/10.25226/bboc.v137i1.2017.a9>
- Gvozdenović S, Schweiger M (2014) Melanism in *Natrix natrix* and *Natrix tessellata* (Serpentes: Colubridae) from Montenegro. *Ecologica Montenegrina* 1: 231–233. <https://doi.org/10.37828/em.2014.1.30>
- Habiboğlu T, Çiçek K, Tok CV (2016) A case of melanism in the Grass snake, *Natrix natrix* (L., 1758) (Reptilia: Colubridae) from Gökçeada (Imbros), Turkey. *Biharean Biologist* 10: 60–61.
- Halpern B, Péchy T, Kiss JB (2002) Scientific research on two isolated *Vipera ursinii moldavica* populations of the Romanian Black Sea coast: preliminary results. *Cercetări Marine – Recherches Marines* 34: 301–311.
- Hodges RJ (2018) Seeing the heat with inexpensive thermography: natural history observations on the northern viper (*Vipera berus*) and grass snake (*Natrix helvetica*). *The Herpetological Bulletin*: 5–13.
- Jablonski D, Kautman J (2017) Melanism in *Natrix tessellata* (Serpentes: Colubridae) from Slovakia. *Herpetology Notes* 10: 173–175.
- Jandzik D (2004) Partial melanism in the grass snake *Natrix natrix* (Reptilia: Colubridae) from northeastern Slovakia. *Acta Zoologica Universitatis Comenianae* 46: 75–77.
- Kabisch K (1999) *Natrix natrix* (Linnaeus, 1758) – Ringelnatter. In: Böhme W (Ed.) *Handbuch der Reptilien und Amphibien Europas*, Bd. 3/IIA, Schlangen (Serpentes) II. Schlangen. Aula-Verlag Wiesbaden, 513–580.
- Kolenda K, Najbar B, Najbar A, Kaczmarek P, Kaczmarek M, Skawiński T (2017) Rare colour aberrations and anomalies of amphibians and reptiles recorded in Poland. *Herpetology Notes* 10: 103–109.
- Kottek M, Grieser J, Beck C, Rudolf B, Rubel F (2006) World Map of the Köppen-Geiger climate classification updated. *Meteorologische Zeitschrift* 15: 259–263. <https://doi.org/10.1127/0941-2948/2006/0130>
- Lorion S, Bonnet X, Brischoux F, De Crignis M (2008) Is melanism adaptive in sea kraits? *Amphibia-Reptilia* 29: 1–5. <https://doi.org/10.1163/156853808783431523>
- Madsen T (1987) Are juvenile grass snakes, *Natrix natrix*, aposematically coloured? *Oikos* 48: 265–267. <https://doi.org/10.2307/3565512>
- Majerus MEN (1998) *Melanism: evolution in action*. 2nd edn. Oxford University Press, 338 pp.
- Mărănducă M, Brănișteanu D, Șerban DN, Brănișteanu DC, Stoleriu G, Manolache NG, Șerban IL (2019) Synthesis and physiological implications of melanic pigments (Review). *Oncology Letters*. <https://doi.org/10.3892/ol.2019.10071>
- Marinov M, Doroșencu AC, Vasile A, Bolboacă LE, Kiss JB, Nanu C, Tošić K, Tudor M (2019) Recent data regarding colonial waterbirds in Danube Delta Biosphere Reserve (Romania). *Journal of Environmental Protection and Ecology* 20: 276–284.
- Millien V, Lyons SK, Olson L, Smith FA, Wilson AB, Yom-Tov Y (2006) Ecotypic variation in the context of global climate change: revisiting the rules. *Ecology Letters* 9: 853–869. <https://doi.org/10.1111/j.1461-0248.2006.00928.x>
- Mollov I (2012) Another case of melanism in the Grass snake *Natrix natrix* (Linnaeus, 1758) (Reptilia: Colubridae) from Bulgaria. *ZooNotes* 28: 1–3.
- Muri D, Schuerch J, Trim N, Golay J, Baillifard A, El Taher A, Dubey S (2015) Thermoregulation and microhabitat choice in the polymorphic asp viper (*Vipera aspis*). *Journal of Thermal Biology* 53: 107–112. <https://doi.org/10.1016/j.jtherbio.2015.06.009>
- Nash DJ, Robinson S, Lewis TR (2016) Observations on the distribution of melanistic snakes in Britain. *The Herpetological Bulletin* 136: 19–22.
- Naumov B (2007) Subclass snake. In: Biserkov V (Ed.) *A Field Guide to Amphibians and Reptiles of Bulgaria*. Green Balkans, Sofia, Bulgaria, 196 pp.
- Naumov B, Tomović L (2005) A case of melanism in *Natrix natrix* (Linnaeus, 1758) (Reptilia: Colubridae) in Bulgaria. *Acta Zoologica Bulgarica* 57: 253–254.
- Prejbeanu I, Rada C (2007) The Danube Delta—an avian paradise. *Interdisciplinary Environmental Review* 9: 210–220. <https://doi.org/10.1504/IER.2007.053989>
- Rasband WS (1997) ImageJ. U. S. National Institutes of Health, Bethesda, Maryland, USA. <https://imagej.nih.gov/ij/>
- Speybroeck J, Beukema W, Bok B, van der Voort J (2016) *Field guide to the amphibians and reptiles of Britain and Europe*. Bloomsbury Publishing, 432 pp.
- Stojanov A, Tzankov N, Naumov B (2011) 35 Die amphibien und reptilien Bulgariens. Edition Chimaira, 582 pp.
- Strugariu A, Zamfirescu ȘR, Gherghel I (2009) First record of the adder (*Vipera berus berus*) in Argeș County (Southern Romania). *Biharean Biologist* 3: 163–166.
- Tanaka K (2007) Thermal biology of a colour-dimorphic snake, *Elaphe quadrivirgata*, in a montane forest: do melanistic snakes enjoy thermal advantages? *Biological Journal of the Linnean Society* 92: 309–322. <https://doi.org/10.1111/j.1095-8312.2007.00849.x>
- Terhivuo J (1990) Relative regional abundance and colour morphs of the adder (*Vipera berus* L.), grass snake (*Natrix natrix* L.), slow worm (*Anguis fragilis* L.) and common toad (*Bufo bufo* L.) in Finland. *Annales Zoologici Fennici* 27: 11–20.
- Thorpe RS (1979) Multivariate analysis of the population systematics of the ringed snake, *Natrix natrix* (L.). *Proceedings of the Royal Society of Edinburgh, Section B, Biological Sciences* 78: 1–62. <https://doi.org/10.1017/S026972700001294X>
- True JR (2003) Insect melanism: the molecules matter. *Trends in Ecology & Evolution* 18: 640–647. <https://doi.org/10.1016/j.tree.2003.09.006>
- Vespremeanu-Stroe A, Zăinescu F, Preoteasa L, Tăuți F, Rotaru S, Morhange C, Stoica M, Hanganu J, Timar-Gabor A, Cărdan I, Piotrowska N (2017) Holocene evolution of the Danube delta: An integral reconstruction and a revised chronology. *Marine Geology* 388: 38–61. <https://doi.org/10.1016/j.margeo.2017.04.002>
- Yenmiş M, Bayrakçı Y, Ayaz D (2022) Skin structure, coloration, and habitat utilization in typical and melanistic morphs of the grass snake (*Natrix natrix*). *The Science of Nature* 109: 12. <https://doi.org/10.1007/s00114-022-01794-w>
- Zadavec M, Lauš B (2011) Melanism variations in *Natrix natrix* (Linnaeus, 1758) and *Zamenis longissimus* (Laurenti, 1768) in Croatia. *Hyla: Herpetological bulletin* 2011: 39–42.
- Zamfirescu ȘR, Strugariu A, Gherghel I, Zamfirescu O (2010) Sfântu Gheorghe (Tulcea, Romania): an important herpetological area. *Analele Științifice ale Universității “Al. I. Cuza”, Iași, Biologie animală* LVI, 119–128.
- Zuffi MAL (2008) Colour pattern variation in populations of the European Whip snake, *Hierophis viridiflavus*: does geography explain everything? *Amphibia-Reptilia* 29: 229–233. <https://doi.org/10.1163/156853808784124929>

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Herpetozoa](#)

Jahr/Year: 2022

Band/Volume: [35](#)

Autor(en)/Author(s): Fanaru Geanina, Telea Alexandra E., Gherghel Iulian, Melenciuc Raluca

Artikel/Article: [Melanism in the grass snake *Natrix natrix* \(Linnaeus, 1758\) from the Danube Delta Biosphere Reserve, Romania 257-263](#)