

# The advertisement calls of *Pristimantis galdi* Jiménez de la Espada, 1870 and *Pristimantis katoptroides* (Flores, 1988) (Anura, Strabomantidae)

Diego Batallas<sup>1,2</sup>, Jorge Brito<sup>3</sup>

<sup>1</sup> Departamento de Biodiversidad, Ecología y Evolución, Facultad de Ciencias Biológicas, Programa de Doctorado en Biología, Universidad Complutense de Madrid, Calle J.A. Nováis 12, Madrid 28040, Spain

<sup>2</sup> Laboratorio de Biología Evolutiva, Colegio de Ciencias Biológicas y Ambientales COCIBA, Universidad San Francisco de Quito USFQ, vía Interoceánica y Diego de Robles 17-1200-841, Quito, Ecuador

<sup>3</sup> Instituto Nacional de Biodiversidad (INABIO). Calle Rumipamba 341 y Av. de Los Shyris. Casilla Postal 17-07-8976, Quito, Ecuador

<https://zoobank.org/17257AAD-00F7-49D0-803B-1D33F705F48F>

Corresponding author: Diego Batallas ([bioandino\\_kakaram@hotmail.com](mailto:bioandino_kakaram@hotmail.com))

Academic editor: Eva Ringler ♦ Received 18 November 2022 ♦ Accepted 13 March 2023 ♦ Published 21 March 2023

## Abstract

In this study we describe for the first time the calls of *Pristimantis galdi* and *Pristimantis katoptroides*. Recordings were obtained in Sangay National Park, Ecuador. We highlight the importance of recording *P. galdi* since its call has been recorded after 153 years of having been described as a species. The call of *P. galdi* consists of 7 to 9 short notes, the sounds of which are similar to a hammer hitting a nail, with a mean dominant frequency of 2.39 kHz. In turn, the call of *P. katoptroides* consists of a single note, the sound of which is similar to a metallic “tic”, with a mean dominant frequency of 1.74 kHz. We compared the advertisement calls of *P. galdi*, *P. katoptroides* and *P. roni* as these species share similar morphological characteristics and are grouped in the spiny green frog’s ecotype. Despite these morphological similarities, their advertisement calls are different. Obtaining calls of *Pristimantis* species in Ecuador might prove difficult with short-term studies due to the great sampling efforts that may be needed to get these recordings. Therefore, implementing active and passive monitoring could help improve our knowledge of acoustic signals in Ecuador’s rainfrogs.

## Key Words

acoustic communication, calls, Sangay National Park, spiny green frogs

Acoustic communication is one of the Anurans’ most important and conspicuous ethological traits, through which they transmit messages between conspecifics (Duellman and Trueb 1994; Angulo 2006; Wells 2007). Acoustic emissions respond to a specific social context and function (Toledo et al. 2015), where the advertisement call is the most commonly heard one in nature. This type of call announces sexual receptivity, position in a territory, and individual size (Gerhardt and Huber 2002; Wells 2007; Toledo et al. 2015; Köhler et al. 2017). In general, the advertisement call consists of a repetitive series of notes

or pulses and, as a result of this stereotypical structure, tends to have a high degree of specificity among species (Gerhardt 2001; Wells and Schwartz 2007). These characteristics have allowed the use of the advertisement call as an identification feature among species (Angulo 2006). Moreover, analyzing and determining call variations has allowed for gaining a better understanding of the diversity of certain groups. Especially in those where morphological characteristics are insufficient to differentiate and diagnose species. (Angulo and Reichle 2008; Rivera-Correa et al. 2022).

*Pristimantis* Jiménez de la Espada, 1870 is a genus of anuran restricted to the New World, known as the most speciose vertebrate genus (595 species; Frost 2023). Nearly 20 species of *Pristimantis* have been described in the last two years in Ecuador, thereby making it the country with the highest description rate per year of this genus (Reyes-Puig and Mancero 2022). In fact, Ecuador counts 254 species, out of which 61.8% (157 species) are endemic (Ron et al. 2022). Despite this considerable rate of discoveries and publications, a large percentage of the vocalizations of the *Pristimantis* individuals of Ecuador remains unknown (Batallas and Brito 2016). In this study, we describe for the first time the advertisement calls of *Pristimantis galdi* Jiménez de la Espada, 1870 and *Pristimantis katoptroides* (Flores, 1988). Furthermore, we compare these results with the call of *Pristimantis roni* Yanez-Muñoz, Bejarano-Muñoz, Brito-M and Batallas-Revelo, 2014, since it is a species with some morphological resemblance.

Fieldwork was conducted in June and September of 2014 in the lower zone of Sangay National Park (administrative jurisdiction comprising the eastern foothills of the park within an altitudinal range of 600 to 1800 m), which corresponds to the Subtropical zoogeographic region (Albuja et al. 2012). The ecosystem comprises montane evergreen forest on the southern slope of the Eastern Andes (Ministerio Del Ambiente del Ecuador 2013). The habitat is characterized by trees whose gnarled trunks and branches support several species of epiphytes such as orchids and bromeliads. The emergent vegetation reaches up to 30 m in height, and there is a visual predominance of romerillo (*Prumnopitys montana*) and palma real (*Dictyocaryum lamarckianum*) trees. In the understory, there is visual predominance of herbaceous plants of the Araceae and Marantaceae families (Brito et al. 2017).

Two calls from a *Pristimantis galdi* individual collected at Sardinayacu lake complex (2°04'20.5"S, 78°12'52.4"W, 1,800 m) were recorded on June 24, 2014, between 14:00–16:00h. Meanwhile, 30 calls from three *Pristimantis katoptroides* individuals collected at Danu (2°04'45.3"S, 78°09'37.3"W, 1,360 m) were recorded on September 21, 2014, between 19:00–20:00h. The calls were obtained using a digital recorder (Olympus WS-802), coupled to a directional microphone (Sennheiser ME 66-K6), placed between 1–3m from the active individual. All recordings were made at a sampling rate of 44.1 kHz and 16 “bits” resolution, saving the audio files in the uncompressed WAV format. It is also worth noting that no fixed hours were set for recording during fieldwork, and it was rather the vocal activity of species that influenced the sampling effort. Collected specimens were sacrificed according to Chen and Combs (1999) protocols and preserved following Simmons (2002). They are deposited at the División de Herpetología del Instituto Nacional de Biodiversidad (DHMECN), Quito, Ecuador. The recordings are deposited at the Fonoteca Zoológica ([www.fonozoo.com](http://www.fonozoo.com)) del Museo Nacional de Ciencias Naturales (CSIC), Madrid, Spain (Appendix 1). Identification of specimens was based on morphological characteristics described in Lynch and Duellman, (1980), Flores (1988), and Duellman and Lehr (2009).

The spectral and temporal properties of the advertisement calls were analyzed with the software Raven 1.6 (K. Lisa Yang Center for Conservation Bioacoustics at the Cornell Lab of Ornithology 2022). The spectrograms were obtained using the Hann window at 90% overlap and 512 samples of DFT size. The parameters analyzed were: (1) Dominant frequency of the entire call and each of the elements emitted in it; (2) Harmonics; (3) Duration of calls; (4) Intervals between calls; (5) Calls/minute; (6) Notes/call; (7) Duration of notes; (8) Intervals between notes; (9) Notes/second. Definitions, terminology and measurements of acoustic parameters follow the terms of Köhler et al. (2017), and Sueur (2018). The figures were processed in R software (R Core Team 2022). For this, the audio files were imported with the tuneR package version 1.4.1 (Ligges et al. 2018). Subsequently, the oscillogram and spectrogram were created using the Seewave package version 2.2.0 (Sueur et al. 2008). With the values of the analyzed parameters, the measures of central tendency (means) and dispersion (maximum, minimum, and standard deviation) were calculated.


#### *Pristimantis galdi* Jiménez de la Espada, 1870 (Fig. 1A)

The recorded male was calling from shrub vegetation approximately 150 cm above the ground. The advertisement call (Fig. 2A, Table 1) consists of 7 to 9 notes. Call duration ranges from 1730 to 2017 ms. The mean note duration is  $44.62 \pm 12.50$  ms, with a mean rate of  $4 \pm 0.96$  notes/second. The mean interval between notes duration is  $217.57 \pm 54.16$  ms. The mean dominant frequency is  $2.39 \pm 0.16$  kHz, showing a partial harmonic whose mean frequency is  $4.76 \pm 0.36$  kHz. Its calls are a constant metallic sound, similar to a hammer hitting a nail.

#### *Pristimantis katoptroides* (Flores 1988) (Fig. 1B)

Recorded males were calling perched on tree branches, about 2–4 m above the ground. This species vocalizes in choruses of about seven to ten males, often observed in smaller groups of up to four individuals in the same tree. The advertisement call consists of a single note with a mean duration of  $54.73 \pm 5.69$  ms. The mean interval between notes is  $44.84 \pm 25.14$  ms, and mean rate is  $1.80 \pm 0.99$  calls/minute. The mean dominant frequency is  $1.74 \pm 0.11$  kHz, showing four partial harmonics in the spectrogram (Fig. 2B, Table 1). The calls have a metallic sound, onomatopoeically resembling the “tic” of a bell.


We describe for the first time the spectral and temporal parameters of the advertisement call of *Pristimantis galdi* and *Pristimantis katoptroides*. The former is assigned to the *Pristimantis* (*Pristimantis*) *galdi* species group (Hedg-es et al. 2008; Padial et al. 2014), where other species’ calls remain unknown. Both, *P. galdi* and *P. katoptroides* (not assigned to any group), are placed in the spiny green frog’s ecotype (Yanez-Muñoz et al. 2014). Within this ecotype, *Pristimantis roni* is morphologically similar to *P. katoptroides* (Yanez-Muñoz et al. 2014). However, these species with similar morphological characteristics differ in the properties of their advertisement calls. The



**Figure 1.** Individuals in their natural habitat. **A.** Adult male of *Pristimantis galdi* (DHMECN 12189, SVL 20 mm) from Sardinayacu, Parque Nacional Sangay, Ecuador; **B.** Adult male of *Pristimantis katoptroides* (DHMECN 12169, SVL 28.15 mm) from Danu, Parque Nacional Sangay, Ecuador.

call structure in both species is different, as *P. roni* emits several notes while *P. katoptroides* emits one. The calls of *P. katoptroides* are shorter, with longer intervals between calls and a lower number of emissions per minute. Finally, the dominant frequency in *P. katoptroides* is lower than that of *P. roni*. In the case of *P. galdi*, the call structure is similar (both species emit several notes), and the dominant frequency is lower than that of *P. roni* (see Fig. 3, Table 1 for further details). The advertisement calls from the three compared species share that they resemble a metallic sound (an onomatopoeic sound of a “Tic”).

Among cryptic species, the use of acoustic diagnostic features for identification could be an alternative to morphometric and molecular diagnosis (Padial et al. 2009; Rivera-Correa et al. 2022). Even though morphologically similar, *P. galdi*, *P. katoptroides*, and *P. roni* have calls with different features, which supports the specificity of their acoustic signals at the species level. However, the magnitude of cryptic diversity and their advertisement calls remain largely unknown. (Funk et al. 2012; Reyes-Puig and Mancero 2022). Direct developing frogs, as is the case of




**Figure 2.** Spectrograms and Oscillograms of the advertisement calls described in this study. **A.** *Pristimantis galdi* (DHMECN 12189, SVL 20 mm, 18 °C air temperature); **B.** *Pristimantis katoptroides* (DHMECN 12169, SVL 28.15 mm, 17.1 °C air temperature).

**Table 1.** Spectral and temporal values of *Pristimantis* calls belonging to the spiny green frog’s ecotype (*sensu* Yáñez-Muñoz et al. 2014). The abbreviations used in the parameters correspond to: kHz = kilohertz, ms = milliseconds, s = seconds. In the analyzed sample, n=number of specimens/calls/notes.

	<i>P. galdi</i> n=1/2/16	<i>P. katoptroides</i> n=3/30	<i>P. roni</i> n=1/9/50
Dominant frequency (kHz)	2.19–2.58 (2.39 ± 0.16)	1.60–2.06 (1.74 ± 0.11)	3.14–3.35 (3.32 ± 0.06)
Harmonic 1 (kHz)	4.35–5.34 (4.76 ± 0.36)	3.19–3.99 (3.43 ± 0.24)	–
Harmonic 2 (kHz)	–	4.94–5.09 (5.03 ± 0.05)	–
Harmonic 3 (kHz)	–	6.53–6.84 (6.65 ± 0.10)	–
Harmonic 4 (kHz)	–	8.80–9.91 (9.65 ± 0.42)	–
Call duration (ms)	1730–2017	42–64 (54.73 ± 5.69)	40–1247 (493.09 ± 341.67)
Interval between calls (s)	–	16.05–94.92 (44.84 ± 25.14)	2.09–7.93 (6.13 ± 1.71)
Call rate (calls/min)	–	0.64–3.72 (1.80 ± 0.99)	6.73–12.44 (8.79 ± 1.76)
Notes per call	7–9	1	1–5
Note duration (ms)	30–77 (44.62 ± 12.50)	–	29–45 (36.84 ± 4.68)
Interval between notes (ms)	151–325 (217.57 ± 54.16)	–	175–298 (249.56 ± 35.91)
Note rate (notes/s)	2.54–5.29 (4 ± 0.96)	–	–
Source	This work	This work	Yáñez-Muñoz et al. 2014

*Pristimantis* species, do not conform to chorus aggregations and individuals are dispersed in their habitat (Duellman and Lehr 2009). Therefore, obtaining recordings of their calls could be more difficult due to their life history traits. In fact, the data collected in this study comes from two of the 17 expeditions conducted between 2010 to 2016 in Sangay National Park and buffer zones (Brítez et al. 2017).



**Figure 3.** Comparative oscillograms and spectrograms of advertisement calls of three *Pristimantis* species belonging to the spiny green frog's ecotype. **A.** *P. galdi* (DHMECN 12189, SVL 20 mm, 18 °C air temperature); **B.** *P. katoptroides* (DHMECN 12169, SVL 28.15 mm, 17.1 °C air temperature); **C.** *P. roni* (DHMECN 11322, SVL 16.55 mm, 15.6 °C air temperature).

Bioacoustics, behavioral and natural history information for species of the genus *Pristimantis* is very scarce, which contrasts with their abundant diversity (Hutter et al. 2016; Rivera-Correa et al. 2021). In Ecuador, such a contrast becomes more pronounced when taking into consideration the description rate of new species presented in recent years (Reyes-Puig and Mancero 2022). The fact that the call from a species such as *P. galdi*, which was described 153 years ago, has just been recorded for the first time suggests that recording the calls from *Pristimantis* species in Ecuador is difficult to obtain with short-term studies including direct visual encounters. Even though these methods are complementary and necessary for detecting and recording species in prospective phases (Boullhesen et al. 2021), complementary methods, such as passive acoustic monitoring, could help with recording the acoustic signals in one of Ecuador's most diverse and yet unknown groups of frogs.

## Acknowledgments

We thank the staff at Sangay National Park, especially Victor León, for supporting field logistics. Thanks to the Tenecota family for providing housing and logistic support in the localities of Sardinayacu and Danu. We are grateful to Glenda Pozo, Patricia Bejarano, Jenny Curay and Hernan Orellana for their invaluable field work assistance. The revision of Oliver Thomas, who provided useful comments on the first version of the manuscript, and who also revised the grammar, is gratefully acknowledged. We especially thank Edgar Lehr and Eva Ringler for their valuable and constructive comments that helped to improve the quality of this manuscript. Thanks also

to the Ministerio del Ambiente de Morona Santiago for issuing research permit N°. 05-2014-I-B-DPMS/MAE. This work is dedicated to the memory of Luis Rivadeneira.

## References

- Albuja L, Almendáriz A, Barriga R, Cáceres F, Montalvo L, Román J (2012) Fauna de Vertebrados del Ecuador. Escuela Politécnica Nacional. Editorial Arial 12, Quito, 490 pp.
- Angulo A, Reichle S (2008) Acoustic signals, species diagnosis, and species concepts: the case of a new cryptic species of *Leptodactylus* (Amphibia, Anura, Leptodactylidae) from the Chapare region, Bolivia. *Zoological Journal of the Linnean Society* 152: 59–77. <https://doi.org/10.1111/j.1096-3642.2007.00338.x>
- Angulo A (2006) Fundamentos de bioacústica y aspectos prácticos de grabaciones y análisis de cantos. In: Angulo A, Rueda-Almonacid JV, Rodríguez-Mahecha JV, La Marca E (Eds) *Técnicas de inventario y monitoreo para los anfibios de la Región Tropical Andina*. Conservación Internacional, Serie Manuales para la conservación N°2, Panamericana Formas e Impresos SA, Bogotá, 93–134.
- Batallas D, Brito J (2016) Descripción de las vocalizaciones de *Pristimantis ganonotus* (Anura: Craugastoridae) de Ecuador. *Cuadernos de Herpetología* 30: 31–34.
- Brito J, Batallas D, Yáñez MH (2017) Ranas terrestres *Pristimantis* (Anura: Craugastoridae) de los bosques montanos del Río Upaño, Ecuador: Lista anotada, patrones de diversidad y descripción de cuatro especies nuevas. *Neotropical Biodiversity* 3: 125–156. <https://doi.org/10.1080/23766808.2017.1299529>
- Boullhesen M, Vaira M, Barquez RM, Akmentins MS (2021) Evaluating the efficacy of visual encounter and automated acoustic survey methods in anuran assemblages of the Yungas Andean forests of Argentina. *Ecological Indicators* 127: 107750. <https://doi.org/10.1016/j.ecolind.2021.107750>



- Chen HM, Combs CA (1999) An alternative anesthesia for amphibians: ventral application of benzocaine. *Herpetological Review* 30: 34.
- Duellman WE, Lehr E (2009) Terrestrial-breeding Frogs (Strabomantidae) in Peru. Natur und Tier-Verlag, Münster, Germany.
- Duellman WE, Trueb L (1994) Biology of Amphibians. The John Hopkins University Press, London. <https://doi.org/10.56021/9780801847806>
- Flores G (1988) Two new species of Ecuadorian *Eleutherodactylus* (Leptodactylidae) of the *E. crucifer* Assembly. *Journal of Herpetology* 22: 34–41. <https://doi.org/10.2307/1564354>
- Frost DR (2023) Amphibian Species of the World: an Online Reference. Version 6.1. American Museum of Natural History, New York, USA. <https://amphibiansoftheworld.amnh.org/index.php> [Accessed 9 January 2023]
- Funk WC, Caminer M, Ron SR (2012) High levels of cryptic species diversity uncovered in Amazonian frogs. *Proceedings of the Royal Society B* 279: 1806–1814. <https://doi.org/10.1098/rspb.2011.1653>
- Gerhardt HC, Huber F (2002) Acoustic Communication in Insects and Anurans: Common Problems and Diverse Solutions. Chicago University Press, USA, 542 pp.
- Gerhardt HC (2001). Acoustic communication in two groups of closely related tree frogs. *Advances in the Study of Behavior* 30: 99–167. [https://doi.org/10.1016/S0065-3454\(01\)80006-1](https://doi.org/10.1016/S0065-3454(01)80006-1)
- Hedges SB, Duellman WE, Heinicke MP (2008) New World direct-developing frogs (Anura, Terrarana), molecular phylogeny, classification, biogeography, and conservation. *Zootaxa* 1737:1–182. <https://doi.org/10.11646/zootaxa.1737.1.1>
- Hutter CR, Liu V, Kell T, Lyons JA, Guayasamin, JM (2016). The Natural History, Distribution, and Conservation of Lonely Rainfrogs, *Pristimantis eremitus*. *Herpetologica* 72: 13–22. <https://doi.org/10.1655/Herpetologica-D-14-00020>
- K. Lisa Yang Center for Conservation Bioacoustics at the Cornell Lab of Ornithology (2022) Raven Pro: Interactive Sound Analysis Software (Version 1.6.3) [Computer software]. Ithaca, NY: The Cornell Lab of Ornithology. <https://ravensoundsoftware.com/>
- Köhler J, Jansen M, Rodríguez A, Kok PJR, Toledo LF, Emmrich M, Glaw F, Haddad CFB, Rödel MO, Vences M (2017) The use of bioacoustics in anuran taxonomy: theory, terminology, methods and recommendations for best practice. *Zootaxa* 4251: 1–124. <https://doi.org/10.11646/zootaxa.4251.1.1>
- Ligges U, Krey S, Mersmann O, Schnackenberg S (2018) tuneR: Analysis of music. R package versión 1.3.3. <https://CRAN.R-project.org/package=tuneR>
- Lynch JD, Duellman WE (1980) The *Eleutherodactylus* of the Amazonian slopes of the Ecuadorian Andes (Anura: Leptodactylidae). Miscellaneous Publications, Museum of Natural History, University of Kansas 69: 1–86. <https://doi.org/10.5962/bhl.title.16222>
- Ministerio del Ambiente del Ecuador (2013) Sistema de Clasificación de los Ecosistemas del Ecuador Continental. Subsecretaría de Patrimonio Natural, Quito, 232 pp.
- Padial JM, Grant T, Frost DR (2014) Molecular systematics of Terraranas (Anura: Brachycephaloidea) with an assessment of the effects of the alignment and optimality criteria. *Zootaxa* 3825: 1–132. <https://doi.org/10.11646/zootaxa.3825.1.1>
- Padial JM, De la Riva I (2009) Integrative taxonomy reveals cryptic Amazonian species of *Pristimantis* (Anura: Strabomantidae). *Zoological Journal of the Linnean Society* 155: 97–122. <https://doi.org/10.1111/j.1096-3642.2008.00424.x>
- R Development Core Team (2022) R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing. <http://www.R-project.org>
- Reyes-Puig C, Mancero E (2022) Beyond the species name: an analysis of publication trends and biases in taxonomic descriptions of rainfrogs (Amphibia, Strabomantidae, Pristimantis). *ZooKeys* 1134: 73–100. <https://doi.org/10.3897/zookeys.1134.91348>
- Rivera-Correa M, Correa-Medina H, Venegas-Valencia K, Daza JM (2022) Genetic diversity, acoustic signal and geographic distribution of a colourful rain frog of the genus *Pristimantis* (Anura: Strabomantidae). *Herpetology Notes* 15: 215–227.
- Ron SR, Merino-Viteri A, Ortiz DA (2022) Anfíbios del Ecuador. Version 2022.0. Museo de Zoología, Pontificia Universidad Católica del Ecuador. <https://bioweb.bio/faunaweb/amphibiaweb> [Accessed 11 January 2023]
- Simmons JE (2002) Herpetological collecting and collection management. *Herpetological Circular* 31: 1–153.
- Sueur J (2018) Sound Analysis and Synthesis with R. Cham, Switzerland: Springer, 637 pp. <https://doi.org/10.1007/978-3-319-77647-7>
- Sueur J, Aubin T, Simonis C (2008) Seewave: a free modular tool for sound analysis and synthesis. *Bioacoustics* 18: 213–226. <https://doi.org/10.1080/09524622.2008.9753600>
- Toledo LF, Martins IA, Bruschi DP, Passos MA, Alexandre C, Haddad CF (2015) The anuran calling repertoire in the light of social context. *Acta Ethologica* 18: 87–99. <https://doi.org/10.1007/s10211-014-0194-4>
- Wells KD (2007) The Ecology and Behavior of Amphibians. University of Chicago Press, Chicago, 1148 pp. <https://doi.org/10.7208/chicago/9780226893334.001.0001>
- Wells KD, Schwartz JJ. (2007) The behavioral ecology of anuran communication. In: Narins PM, Feng AS, Fay RR, Popper AN (Eds) Hearing and sound communication in amphibians. Springer Handbook of Auditory Research, Springer, New York, 44–86. [https://doi.org/10.1007/978-0-387-47796-1\\_3](https://doi.org/10.1007/978-0-387-47796-1_3)
- Yáñez-Muñoz MH, Bejarano-Muñoz EP, Brito J, Batallas D (2014) Ranas terrestres de los Andes Surorientales de Ecuador II: Una nueva especie de *Pristimantis* verde espinosa de los bosques montanos del Parque Nacional Sangay (Anura: Craugastoridae). *Avances en Ciencias e Ingenierías* 6: 63–77. <https://doi.org/10.18272/aci.v6i2.180>

## Appendix 1

### Specimens recorded and collected:

*Pristimantis galdi* (Ecuador): Provincia de Morona Santiago, Complejo lacustre de Sardinayacu, Parque Nacional Sangay: DHMECN 12189, SVL 20 mm.46; FZ 11490–91

*Pristimantis katoptroides* (Ecuador): Provincia de Morona Santiago, Danu, Parque Nacional Sangay: DHMECN 12169, SVL 28.15 mm, FZ 11492; DHMECN 12170, SVL 24.05 mm, FZ 11493; DHMECN 12172, SVL 26.24 mm, FZ 11493.

*Pristimantis roni* (Ecuador): Provincia de Morona Santiago,  
Complejo lacustre de Sardinayacu, Parque Nacional  
Sangay: DHMECN 11322 SVL 16.55 mm, FZ 14208

## Supplementary material 1

***Pristimantis galdi* recording 1. Fonozoo library code: FZ 11491**

Authors: Diego Batallas, Jorge Brito

Data type: WAV file

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/herpetozoa.36.e97705.suppl1>

## Supplementary material 2

***Pristimantis galdi* recording 2. Fonozoo library code: FZ 11490**

Authors: Diego Batallas, Jorge Brito

Data type: WAV file

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/herpetozoa.36.e97705.suppl2>

## Supplementary material 3

***Pristimantis katoptroides* recording 1. Fonozoo library code: FZ 11492**

Authors: Diego Batallas, Jorge Brito

Data type: mp3 file

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/herpetozoa.36.e97705.suppl3>

## Supplementary material 4

***Pristimantis katoptroides* recording 2. Fonozoo library code: FZ 11493**

Authors: Diego Batallas, Jorge Brito

Data type: mp3 file

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/herpetozoa.36.e97705.suppl4>

## Supplementary material 5

***Pristimantis katoptroides* recording 3. Fonozoo library code: FZ 11494**

Authors: Diego Batallas, Jorge Brito

Data type: mp3 file

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/herpetozoa.36.e97705.suppl5>

## Supplementary material 6

***Pristimantis roni* recording. Fonozoo library code: FZ 14208**

Authors: Diego Batallas, Jorge Brito

Data type: WAV file

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/herpetozoa.36.e97705.suppl6>

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Herpetozoa](#)

Jahr/Year: 2023

Band/Volume: [36](#)

Autor(en)/Author(s): Batallas Diego, Brito Jorge

Artikel/Article: [The advertisement calls of \*Pristimantis galdi\* Jiménez de la Espada, 1870 and \*Pristimantis katoptroides\* \(Flores, 1988\) \(Anura, Strabomantidae\) 59-64](#)