

The secret life of a rock-dweller: arboreal acrobatics observed in the European leaf-toed gecko *Euleptes europaea*

Daniele Salvi¹, Emanuele Berrilli¹, Giacomo Bruni², Matteo Garzia¹, Veronica Gomes³, Giacomo Radi⁴, Michel-Jean Delaugerre⁵

¹ Department of Health, Life & Environmental Sciences, University of L'Aquila, Via Vetoio snc, 67100 L'Aquila-Coppito, Italy

² Viale Palmiro Togliatti, 50019 Sesto Fiorentino, Italy

³ CIBIO Research Centre in Biodiversity and Genetic Resources, InBIO, BIOPOLIS, Universidade do Porto, Campus de Vairão, Rua Padre Armando Quintas, 7, 4485-661 Vairão, Portugal

⁴ Museo di Storia Naturale della Maremma, Str. Corsini, 5, 58100 Grosseto, Italy

⁵ Conservatoire du littoral, Rés. St Marc, 2, rue Juge Falcone 20200 Bastia, France

<https://zoobank.org/C6B5E0C8-195D-43BF-82E1-E8B0B58466C0>

Corresponding author: Michel-Jean Delaugerre (dmighe@gmail.com)

Academic editor: Günter Gollmann ♦ Received 13 March 2023 ♦ Accepted 9 May 2023 ♦ Published 9 June 2023

Abstract

The European leaf toed Gecko, *Euleptes europaea*, is a strictly nocturnal species endemic to the western Mediterranean and has long been considered a rock-specialist as it is associated with this habitat during its entire daily and life cycle. In this study, we report observations of arboreal behaviour in *E. europaea*, collected during field research over a 40-year period from across the entire species range. We provide a review of the available information on this topic that contributes to a refined view of the habitat uses and arboreal abilities of this species. Arboreal behaviour in *E. europaea* was observed throughout the year, across different macrohabitats, on a wide variety of tree, shrub, and bush species, on various parts of the plant (trunk, branches, fronds, twigs, leaves), and at different height from the ground. Remarkably, *E. europaea* shows an extraordinarily agile arboreal locomotion associated with striking morphological adaptations to an arboreal lifestyle, namely a prehensile tail bearing a terminal adhesive pad that supplements grasping force, an equilibrium asset, and scansor adhesion both in static condition and during escape. We conclude that *E. europaea* is a climbing gecko (opposed to ground dwelling), occupying both rocky and arboreal microhabitats. While the evolutionary origin and ecological drivers of the arboreal behaviour of *E. europaea* remains to be fully investigated, this realization has important implications for designing fieldwork research and management strategies for conservation.

Key Words

arboreal behaviour, Gekkota, microhabitat use, prehensile tail, rock crevices, tail pads, vegetation cover

Introduction

Understanding the habitat use of species is crucial for ecologists, biogeographers, and conservationists (MacArthur and Pianka 1966; Caughley 1994; Hanski and Gyllenberg 1997). Species differ greatly in the range of habitat they

use. Some species are habitat generalists while others are specialists, with a gradient of variation between the two extremes. Habitat selection is particularly important for ectothermic lizards such as geckos, because it influences thermal ecology, physiological performance, and individual behaviour, thus having an impact upon population and

community dynamics (Pulliam and Danielson 1991; Pandit et al. 2009). However, for many species, we still have a very limited knowledge of their habitat use.

Here, we report original observations and literature data on the European leaf toed Gecko *Euleptes europaea* that warrant a reconsideration of its habitat use. This strictly nocturnal gecko, endemic to the western Mediterranean (Fig. 1), belongs to a monotypic genus of the Gondwanan family Sphaerodactylidae, mainly distributed in the Neotropics (Gamble et al. 2008), thus representing an outstanding biogeographical relict within the European fauna.




Figure 1. Distribution range the European leaf toed Gecko *Euleptes europaea* and geographic location of the observations of arboricolous behaviour reported in Table 1.

The European Leaf-toed Gecko is considered a rock-specialist and has several anatomical, behavioural, ecological adaptations to this habitat with which it is associated during its entire daily and life cycle (Salvidio et al. 2010). During the day, narrow rock crevices provide this species protection against predators and the opportunity to thermoregulate. Its small size, SVL 30–40 mm (Salvidio et al. 2010; Delaunier and Corti 2020), its smooth (not keeled) body scales and its consequent ability to flatten itself to an extreme extent are probable morphological adaptations to crevice life. Crevices with openings as narrow as 3–5 mm permit this gecko to stay in both dorsal and ventral contact with the stone, preserving it from light and direct sun exposure and from desiccating cold winds, and providing optimal microclimatic conditions (thermal and hygrometric). Geckos are able to thermoregulate by conduction moving inside the shelter heated by the sun; whereas predators, such as birds, mammals, and other squamates, are too large to enter the crevices. In the roomiest cracks, large aggregations up to several dozen geckos may occur (Delaunier 1981, 1992; Salvidio et al. 2010). Along with suitable climatic conditions, the availability and the quality of retreat sites and egg laying sites are likely the

key factor governing the presence and the abundance of this gecko (Salvidio and Oneto 2008).

The nocturnal life of this species is also linked to the rocks. At night, *E. europaea* forages mostly on rocky surfaces (from horizontal to vertical to overhanging) that it climbs easily thanks to its distal toe pads and claws that allow attachment to dusty rock surfaces where other geckos with more powerful basal pads cannot do so – such as *Tarentola mauritanica*– (Russell and Delaunier 2017). The heat stored by the rocks during the day is slowly released overnight making the rocks warmer than the air, wood, and other substrates, enabling this gecko to efficiently thermoregulate by thigmothermy during the night (Delaunier 1984; Salvidio and Oneto 2008).

However, while the association of *E. europaea* to rocky habitats is well established, little is known about its ability to utilize vegetation near or far from rocks. For example, in the original description of this species it was reported that the species could be found under the bark of trees (Gené 1839), and recent observations suggest that, when rats are present, *E. europaea* shifts its spatial behaviour and feeds beneath vegetations (Delaunier et al. 2019). Delaunier (1992) also hypothesized that the animal may disperse in the vegetation during the hottest days of the year when the thermal supplement of the rocky substratum is no longer essential. Nevertheless, the arboreal aptitude of this gecko is still to be explored in order to have a more accurate view of its ecological niche and microhabitat use. Here we provided a comprehensive description of new field observations and a review of the available information on this topic that contribute to a refined view on the habitat use and arboreal abilities of *Euleptes europaea*.

Materials and methods

We collected observations of arboreal behaviour of *E. europaea* during field research in an opportunistic way. That is, vegetation was not primarily searched for the presence of the species but rather the species was accidentally spotted on vegetation while searching on rocks or for other species. Additional observations were gathered from colleagues and from the literature. We use the term “arboreal behaviour” in a broad meaning, including individuals climbing on trees, bushes, or low shrubs either dwelling on the bark, the branches or the leaves. For this tiny gecko, the main difference is likely dwelling on rock faces or on vegetation of any kind. On the other hand, we did not consider as ‘arboreal behaviour’ many observations we gathered during the years of *E. europaea* escape jumps (from heights of up to 3 m) toward other rock and into the vegetation, usually at the base of rock boulders, where they quickly disappear.

For each observation, we described the presence of the geckos on vegetation and their behaviour, and we reported the type of vegetation, the year and locality of the observations and the observers.

Results and discussion

A total of 32 observations of one or more individuals of *E. europaea* on vegetation were collected during 27 surveys. Observations cover 40 years (1982–2022) and 21 locations scattered across the species range (Fig. 1). Detailed information for each observation is reported in Table 1 and Suppl. material 2.

Eighteen different plant/tree species were used by *E. europaea*, mostly maquis species but also alien species such as *Eucalyptus* and *Asparagus aethiopicus* (Table 2). Although the opportunistic sampling does not allow proper statistical testing, the absence of large differences in observation frequency distribution across plant species (Table 2), suggests that the wide range of trees, bushes, and shrubs used by *E. europaea* reflects their availability at the sampling sites rather than plant selection by the gecko. Likewise, although our field research was mainly biased towards

rocky environments (as this was considered as the elective habitat of *E. europaea*), and thus most observations concern vegetation growing on or nearby rocky surfaces, we also observed *E. europaea* on vegetation relatively far from rocks.

Arboreal behaviour of *E. europaea* was observed throughout the year, across the entire species range, on a wide variety of tree, shrub, bush species, on various parts of the plant (trunk, branches, fronds, twigs, leaves) at different heights above the ground (up to 3 m), strongly supporting that this is not an occasional behaviour of this species. This is further supported by a paper made available at the same time of our study that shows a high occupancy probability of *E. europaea* on *Eucalyptus* trees in an insular site (Deso et al. 2023). On the two *Eucalyptus* stands studied by these authors, *E. europaea* was the only gecko occupying the less-anthropized area. It is not yet known how high *Euleptes* geckos dwell in those trees and if this arboricolous habitat does provide egg-laying sites.

Table 1. Observations of arboricolous behaviour of *Euleptes europaea* (*Ee*); see Fig. 1 for the location of the observations. Island (I); islet (is).

Observation	Locality	Year	Period	Observer
1 <i>Ee</i> found on the nape of MD's neck while passing through bush by night	Scandula ⁵ (W Corsica)	1982	May	M. Delaugerre
Most of the <i>Ee</i> forage by night between the base of the granite and a mattress of plants	Lavezzu I ²³ (S Corsica)	1986–2022	June to October	M. Delaugerre, C. Corti, M. Biaggini and P. Lo Cascio
2 <i>Ee</i> found on a bird nest box on <i>Arbutus unedo</i> (2 m high), not far from rocky boulders	Scandula (W Corsica)	1992	June	JL Martin
1 <i>Ee</i> in a bird nest box on <i>Erica arborea</i> (2 m high), not far from rocky boulders	Scandula (W Corsica)	1993	May	JL Martin
1 <i>Ee</i> climbing on <i>Rosmarinus officinalis</i> close to a schist face (Suppl. material 1: fig. S1a)	Port-Cros I ³ (Provence)	2003	October	M. Delaugerre
1 <i>Ee</i> climbing on <i>Lotus cytisoides</i> growing at the base of the rock (Suppl. material 1: fig. S1b)	Gabinière is ⁴ (Provence)	2003	October	M. Delaugerre
1 <i>Ee</i> climbing on <i>Malva arborea</i> (1.30 m high)	Toro is ²¹ (SE Corsica)	2005	April	M. Delaugerre
4 <i>Ee</i> (out of 23 diurnal sightings) found under the bark of dead <i>Ulmus minor</i> trees	Tino is ¹⁰ (Liguria)	2006	20–29 June	F. Oneto, D. Ottonello, and S. Salvidio ^a
1 <i>Ee</i> found on the vest of MD while passing by bushes of <i>Juniperus phoenicea</i> (3 m high)	Rascas is ² (Provence)	2008	17–18 June	M. Delaugerre
1 <i>Ee</i> found under the bark of <i>Eucalyptus</i> (Suppl. material 1: fig. S1c)	Giglio I ¹⁹ (Tuscany)	2008	August	S. Fattorini ^b
1 <i>Ee</i> climbing on <i>Anthyllis barba-jovis</i> , 3 m high	Cala Violina ¹⁴ (Tuscany)	2009	18 October	G. Radi
1 <i>Ee</i> head spotted among <i>Halimione portulacoides</i> (Suppl. material 1: fig. S1d)	Giraglia I ⁸ (N Corsica)	2012	6 October	M. Delaugerre
1 <i>Ee</i> found under the bark of <i>Juniperus phoenicea</i> ; and 1 <i>Ee</i> found under the bark of <i>Cupressus</i> sp.	Cavallo I ²² (S. Corsica)	2014	6 November	V. Rivière
1 <i>Ee</i> found hidden under the bark of <i>Quercus ilex</i>	La Paolina is ¹¹ (Tuscany)	2016	6 May	M. Delaugerre and C. Corti ^c
1 <i>Ee</i> on <i>Olea europaea</i> (1 m high)	Collelungo ¹⁸ (Tuscany)	2017	24 August	G. Radi
1 <i>Ee</i> foraging on the bark of <i>Juniperus Phoenicia</i> (1 m high)	Punta Ala ¹⁵ (Tuscany)	2019	30 September	G. Radi
1 <i>Ee</i> on <i>Anthyllis barba-jovis</i> (1.5 m high)	Punta Ala ¹⁶ (Tuscany)	2020	21 June	G. Radi
1 <i>Ee</i> on <i>Anthyllis barba-jovis</i> (2 m high)	Punta Ala ¹⁷ (Tuscany)	2020	5 October	G. Radi
>10 <i>Ee</i> , including a mating pair, observed by night on <i>Cistus</i> sp. and <i>Erica arborea</i> (2 m high; Fig. 2c)	Monte Albo ²⁴ (Sardinia)	2021	6–7 July	E. Berrilli, M. Garzia, D. Salvi, and V. Gomez
>10 <i>Ee</i> observed by night on <i>Cistus</i> sp. and <i>Erica arborea</i> (1.5 m high; Fig. 2d)	Monte Albo ²⁵ (Sardinia)	2021	8 July	D. Salvi, M. Garzia, and V. Gomez
2 <i>Ee</i> found under the bark of a fallen <i>Pinus halepensis</i> tree (geckos not active)	Positano ²⁷ (Campania)	2022	5 February	F. Russo ^d
>20 <i>Ee</i> active by night on trunks of <i>Quercus ilex</i> and <i>Juniperus phoenicea</i>	Dorgali ²⁶ (Sardinia)	2022	11–13 June	A. Macali and C. Pardo
1 <i>Ee</i> on a <i>Pinus halepensis</i> trunk, 2.5 m high	Montecristo I ²⁰ (Tuscany)	2022	14 June	G. Radi and M. Zuffi
1 <i>Ee</i> found on the nape of MD's neck while passing through <i>Malva arborea</i> (1.5 m high) 1 <i>Ee</i> on the nape of MD's neck	Giraglia I ⁹ (N Corsica)	2022	2 August	M. Delaugerre
2 <i>Ee</i> sighted within <i>Opuntia ficus indica</i> pads	Pomégue I ¹ (Provence)	2022	11 October	V. Lara and V. Rivière
1 <i>Ee</i> on <i>Asparagus aethiopicus</i> growing on a rock (Fig. 2e)	Tellaro ¹¹ (Liguria)	2022	12 December	G. Bruni
10 <i>Ee</i> on <i>Pistacia lentiscus</i> , close to the rock face (Fig. 2a, b, f)	Tellaro ¹² (Liguria)	2022	21 December	G. Bruni

1–27: sampling locality code as shown in Fig. 1.

^a: reported in Oneto et al. (2008, 4)

^b: reported in Corti et al. (2021, 54).

^c: reported in Fattorini (2010, fig. 3).

^d: reported in Di Nicola et al. (2022).

Table 2. Frequency of observations of *Euleptes europaea* on different plants and trees.

Species	N
<i>Anthyllis barba-jovis</i>	3
<i>Arbutus unedo</i>	2
<i>Asparagus aethiopicus</i>	1
<i>Cistus</i> sp.	2
<i>Cupressus</i> sp.	1
<i>Erica arborea</i>	4
<i>Eucalyptus</i> sp.	1
<i>Halimione portulacoides</i>	1
<i>Juniperus phoenicea</i>	4
<i>Lotus cytisoides</i>	2
<i>Malva arborea</i>	2
<i>Olea europaea</i>	1
<i>Opuntia ficus indica</i>	1
<i>Pinus halepensis</i>	2
<i>Pistacia lentiscus</i>	1
<i>Quercus ilex</i>	2
<i>Rosmarinus officinalis</i>	1
<i>Ulmus minor</i>	1

On vegetation this species rests, forages, and mates (Table 1). What is more important, *E. europaea* has been observed to move across branches and twigs with great agility, using its four members and its tail, and shows striking morphological adaptation to the arboreal lifestyle.

In low and thick vegetation, locomotion of *E. europaea* involves climbing, rather than crawling (on rock face), so that it may even “swim and vanish” in a puzzling way, in contrast to its relatively slow escape speed on rocks. In an arboreal context, it may perch head down as truly arboreal lizards do (Fig. 2a, c). Compared to rock faces, arboreal habitat is a much more structurally complex 3D array of branches and surfaces (Clark et al. 2021). This species’ prehensile tail has long been recognized (Fitzinger 1843 p. 95; Wiedersheim 1876; Camerano 1885 p. 501; Mourgue 1910; Eijdsen 1983; Bauer et al. 1997; Fig. 2c, d). Almost all adults bear a cartilaginous regenerated tail, turnip-like, enlarged for fat storage, that is still prehensile. The prehensile tail is a functional trait associated with the arboreal environment where this organ is used to grasp cork, branches, twigs and leaves (Mertens 1964; Eijdsen 1983; Alibardi and Bonfitto 2019). The prehensile tail is also flexible and is an asset for equilibrium and the guiding of escape jumps (Jusufi et al. 2008; Fleming and Bateman 2012) and this species is a good jumper. Its tail is not only prehensile, it also bears terminal adhesive pads, even when regenerated (Eijdsen 1983). In the arboreal context, tail pads supplement the muscular grappling of twigs or branches with dry adhesion of the sub caudal scansors (Fig. 2c, f and Suppl. material 1: fig. S1a, e). As stressed by Higham et al. (2017), in arboreal environments, when performing escape jumps or falls from trees, geckos land on leaves or smooth tree trunks. Arresting this fall requires a high loading of the adhesive system that is enhanced by an extra padded member, the tail. But these caudal adhesive systems are not the exclusive signature of arboreal lifestyle, since they are also used by geckos for climbing steep rock surfaces (Bauer 1998; Koppetsch

et al. 2020) and to prevent the animal falling backward (Bauer and Russell 1994; Jusufi et al. 2008). Therefore, it is likely that these ecomorphological adaptations allow the European Leaf-toed gecko to exploit both rocky and arboreal microhabitats. To explain the origin of such ecomorphological adaptations to arboreal lifestyle we could speculate that these have been inherited by *E. europaea* from its European ancestors living in subtropical forests environments during Miocene (Estes 1969; Müller 2001; Müller and Mödden 2001; Böhme 2003; Čerňanský and Bauer 2010; Daza et al. 2014; Čerňanský et al. 2022).

Our study opens further questions also on the use of vegetated and wooded habitat by *E. europaea*. Could there be a continuum from i) a simple nightly coming and going from rocky habitat to nearby vegetation, as suggested by some observations, ii) to more seasonal wandering movements, ultimately turning back to rocky habitat for winter; iii) to truly arboreal living all year round? The rupicolous habitat niche of *E. europaea* is primarily driven by thermal constraints (Delaugerre 1984). These are particularly severe for a strictly nocturnal ectothermic vertebrate, inhabiting a temperate region where it colonizes Alpine elevations up to 1500 m (Delaugerre 1992). Rocky substrates typically have larger thermal inertia and a greater capacity for heat storage than soil or wood (Huey et al. 1989). Of course, this gecko is an active and efficient thermoregulator that uses the expansion and retraction of melanophores to speed up the acquisition of heat and to slow down its loss (Delaugerre 1984). But being a strict thigmotherm (it never basks) it would still need the rocky substratum for the primary acquisition of heat, during periods of thermal deficit, 5 to 6 months a year. Our finding of arboreal behaviour *E. europaea* in autumn and winter with air temperatures as low as 11 °C (Table 1) is therefore surprising. A possible explanation is that this species is able to locate warm environmental microclimates that retain higher temperatures throughout the night, allowing them to loosen the bond to rocks. The ability to exploit rare microclimates is especially important for nocturnal species, as heterogeneity of environmental temperatures is reduced at night compared to the day (Nordberg and Schwarzkopf 2019). However, understanding the apparent discrepancy between the thermal requirements of the species and its arboreal habits will require dedicated investigation.

In conclusion, to answer the main question of this study: is *E. europaea* a strictly rock-dwelling species? Definitely it is not. This species efficiently uses vegetated and wooded habitats, both in hot summer and colder winter nights, and shows adaptations to climbing and clinging for arboreal locomotion. It is a climbing gecko (opposed to ground dwelling), occupying both saxicolous and arboreal microhabitats like other geckos (Pianka and Pianka 1976; Norris et al. 2021). This realization has important implications for designing fieldwork research and management strategies for conservation. The evolutionary origin and ecological drivers for such behaviour remain to be fully investigated.



Figure 2. *Euleptes europaea* on *Pistacia lentiscus* **a, b, f** locality Tellaro, Liguria; photo by G. Bruni; on *Cistus* sp. **c, d** locality M. Albo, Sardinia; photo by D. Salvi; and on *Asparagus aethiopicus* **e** locality Tellaro, Liguria; photo by G. Bruni.

Acknowledgements

We wish to thank our friends and colleagues M. R. Di Nicola, S. Fattorini, V. Rivière, M.A. L. Zuffi for sharing details on their field observations, and Aaron Bauer for fruitful discussion. A few geckos were manipulated under authorization by the Italian Ministry for the Environment (PNM-0008275) and by France's Préfet de Corse (2A-2020-06-23-003 and 2B-2020-06-16-004). Some studies were undertaken within the Port-Cros National Park and the Natural Reserves of Bouches-de-Bonifacio, Scandola and Iles du Cap Corse. The research was supported by a grant to DS from the Italian Ministry of University and Research (PRIN project 2017KLZ3MA).

References

- Alibardi L, Bonfitto A (2019) Morphology of setae in regenerating caudal adhesive pads of the gecko *Lygodactylus capensis* (Smith, 1849). *Zoology* 133: 1–9. <https://doi.org/10.1016/j.zool.2019.01.003>
- Bauer AM (1998) Morphology of the adhesive tail tips of carphodactylid geckos (Reptilia: Diplodactylidae). *Journal of Morphology* 235: 41–58. [https://doi.org/10.1002/\(SICI\)1097-4687\(199801\)235:1%3C41::AID-JMOR4%3E3.0.CO;2-R](https://doi.org/10.1002/(SICI)1097-4687(199801)235:1%3C41::AID-JMOR4%3E3.0.CO;2-R)
- Bauer AM, Russell AP (1994) Is autotomy frequency reduced in geckos with 'actively functional' tails. *Herpetological Natural History* 2: 1–15.
- Bauer AM, Good DA, Branch B (1997) The taxonomy of the southern african leaf-toed geckos (Squamata: Gekkonidae), with a review of old world. *Proceedings of the California Academy of Sciences* 49: 447–497.
- Böhme M (2003) The Miocene Climatic Optimum: evidence from ectothermic vertebrates of Central Europe. *Palaeogeography, Palaeoclimatology, Palaeoecology* 195: 389–401. [https://doi.org/10.1016/S0031-0182\(03\)00367-5](https://doi.org/10.1016/S0031-0182(03)00367-5)
- Camerano L (1885) *Monografia dei sauri italiani*. Memoria della Reale Accademia delle Scienze. Torino 2(41): 403–481.
- Caughley G (1994) Directions in conservation biology. *Journal of Animal Ecology* 215–244. <https://doi.org/10.2307/5542>
- Čerňanský A, Bauer AM (2010) *Euleptes gallica* Müller (Squamata: Gekkota: Sphaerodactylidae) from the Lower Miocene of North-West Bohemia, Czech Republic. *Folia Zoologica* 59: 323–328. <https://doi.org/10.25225/fozo.v59.i4.a8.2010>
- Čerňanský A, Daza JD, Smith R, Bauer AM, Smith T, Folie A (2022) A new gecko from the earliest Eocene of Dormaal, Belgium: a thermophilic element of the 'greenhouse world'. *Royal Society Open Science* 9: e220429. <https://doi.org/10.1098/rsos.220429>
- Clark J, Clark C, Higham TE (2021) Tail control enhances gliding in arboreal lizards: An integrative study using a 3D geometric model and numerical simulation. *Integrative and Comparative Biology* 61: 579–588. <https://doi.org/10.1093/icb/ibab073>
- Corti C, Cecchi L, Delaunay MJ (2021) Flore et Herpétofaune des Petites Îles et Îlots Autour des Îles d'Elba et de Pianosa (Arcipelago Toscano, Italie). Initiative PIM, Marseille, 60 pp. <http://initiative-pim.org/index.php/2022/01/27/flore-et-herpetofaune-des-petites-iles/>
- Daza JD, Bauer AM, Snively ED (2014) On the fossil record of the Gekkota: Gekkota fossil record. *The Anatomical Record* 297: 433–462. <https://doi.org/10.1002/ar.22856>
- Delaunay M (1981) Sur l'histoire naturelle de *Phyllodactylus europaeus* Gené, (Gekkonidae, Sauria, Reptiles). Port-Cros : étude d'une population naturelle. *Travaux scientifiques Parc. national Port-Cros* (1980) 6: 147–175.
- Delaunay M (1984) Sur l'écologie thermique des geckos *Phyllodactylus europaeus*, *Hemidactylus turcicus* et *Tarentola mauritanica*: rythmes d'activité, températures et activité, répartition altitudinale. *Travaux Scientifiques du Parc Naturel Régional de la Corse* 3: 96–121.
- Delaunay MJ (1992) *Phyllodactyle d'Europe Phyllodactylus europaeus* Gené, 1839. Atlas de Répartition des Batraciens et Reptiles de Corse-. Parc naturel régional Corse/Ecole pratique Hautes études, Pamplune, 60–63.
- Delaunay MJ, Corti C (2020) Tiny but "strong": the European Leaf-toed gecko, *Euleptes europaea*, a terrestrial vertebrate able to survive on tiny islets. *Israel Journal of Ecology and Evolution* 66: 223–230. <https://doi.org/10.1163/22244662-bja10017>
- Delaunay M-J, Sacchi R, Biaggini M, Lo Cascio P, Ouni R, Corti C (2019) Coping with aliens: how a native gecko manages to persist on Mediterranean islands despite the Black rat? *Acta Herpetologica* 14: 89–100. https://doi.org/10.13128/a_h-7746
- Deso G, Priol P, Reynier T, Renet J (2023) High occupancy of European leaf-toed gecko *Euleptes europaea* in two island stands of *Eucalyptus* sp.: tree selection, co-occurrence and habitat effect. *Ethology Ecology & Evolution*. [in press] <https://doi.org/10.1101/2023.02.08.527781> [preprint on bioRxiv]
- Di Nicola MR, Colombo M, Russo F (2022) First record of European leaf-toed gecko *Euleptes europaea* (Gené, 1839) (Squamata, Sphaerodactylidae) in Campania (Italy). *Rivista del Museo Civico di Scienze Naturali "Enrico Caffi", Bergamo* 35: 79–82.
- Eijssden EHT van (1983) Der haftfähige Greifschwanz des Europäischen Blattfingergekkos *Phyllodactylus europaeus* Gené. *Salamandra* (Frankfurt am Main) 19: 1–10.
- Estes R (1969) Die Fauna der miozänen Spaltenfüllung von Neudorf an der March (CSSR): Reptilia (Lacertilia). *Österreich Akad. Wiss. Mathem.-naturw. Kl. 1*, 178: 77–82.
- Fattorini S (2010) Influence of Recent Geography and Paleogeography on the Structure of Reptile Communities in a Land-Bridge Archipelago. *Journal of Herpetology* 44: 242–252. <https://doi.org/10.1670/09-046.1>
- Fitzinger LJ (1843) 1 Systema Reptilium. Braumüller & Seidel. Wien, 106 pp. <https://doi.org/10.5962/bhl.title.4694>
- Fleming PA, Bateman PW (2012) Autotomy, tail regeneration and jumping ability in Cape dwarf geckos *Lygodactylus capensis* (Gekkonidae). *African Zoology* 47: 55–59. <https://doi.org/10.3377/004.047.0110>
- Gamble T, Bauer AM, Greenbaum E, Jackman TR (2008) Evidence for Gondwanan vicariance in an ancient clade of gecko lizards. *Journal of Biogeography* 35: 88–104.
- Gené J (1839) Synopsis Reptilium Sardinia. *Memorie reale Accademia Scienze Fisiche Matematiche*. Torino 2: 257–286.
- Hanski I, Gyllenberg M (1997) Uniting two general patterns in the distribution of species. *Science* 275: 397–400. <https://doi.org/10.1126/science.275.5298.397>
- Higham TE, Russell AP, Niklas KJ (2017) Leaping lizards landing on leaves: escape-induced jumps in the rainforest canopy challenge the adhesive limits of geckos. *Journal of The Royal Society Interface* 14: e20170156. <https://doi.org/10.1098/rsif.2017.0156>

- Huey RB, Peterson CR, Arnold SJ, Porter WP (1989) Hot rocks and not so hot rocks: retreat site selection by garter snakes and its thermal consequences. *Ecology* 70: 931–944. <https://doi.org/10.2307/1941360>
- Jusufi A, Goldman DI, Revzen S, Full RJ (2008) Active tails enhance arboreal acrobatics in geckos. *Proceedings of the National Academy of Sciences* 105: 4215–4219. <https://doi.org/10.1073/pnas.0711944105>
- Koppetsch T, Böhme W, Büsse S, Gorb SN (2020) Comparative epidermal microstructure anatomy and limb and tail osteology of eyelid geckos (Squamata: Eublepharidae): Implications of ecomorphological adaptations. *Zoologischer Anzeiger* 287: 45–60. <https://doi.org/10.1016/j.jcz.2020.05.005>
- MacArthur RH, Pianka ER (1966) On optimal use of a patchy environment. *The American Naturalist* 100: 603–609. <https://doi.org/10.1086/282454>
- Mertens R (1964) Der Eideichschwanz als Haftorgan. *Senckenbergiana Biologica* 45: 117–122.
- Mourgue M (1910) Étude sur le Phyllodactyle d'Europe (*Phyllodactylus europaeus* Gené). *Feuille Jeune Naturaliste* (4) 40: 57–61.
- Müller J (2001) A new fossil species of *Euleptes* from the early Miocene of Montaigny, France (Reptilia, Gekkonidae). *Amphibia-Reptilia* 22: 341–348. <https://doi.org/10.1163/156853801317050133>
- Müller J, Mödden C (2001) A fossil leaf-toed gecko from the Oppenheim/Nierstein Quarry (Lower Miocene, Germany). *Journal of Herpetology*: 529–532. <https://doi.org/10.2307/1565978>
- Nordberg EJ, Schwarzkopf L (2019) Heat seekers: A tropical nocturnal lizard uses behavioral thermoregulation to exploit rare microclimates at night. *Journal of Thermal Biology* 82: 107–114. <https://doi.org/10.1016/j.jtherbio.2019.03.018>
- Norris J, Tingley R, Meiri S, Chapple DG (2021) Environmental correlates of morphological diversity in Australian geckos. In: Sandel B (Ed.) *Global Ecology and Biogeography* 30: 1086–1100. <https://doi.org/10.1111/geb.13284>
- Oneto F, Ottonello D, Salvadio S (2008) Primi dati sulla biometria di *Euleptes europaea* (Gené, 1839) dell'isola del Tino (La Spezia, Liguria) (Reptilia, Squamata, Gekkonidae). *Doriana* 8: 1–8.
- Pandit SN, Kolasa J, Cottenie K (2009) Contrasts between habitat generalists and specialists: an empirical extension to the basic metacommunity framework. *Ecology* 90: 2253–2262. <https://doi.org/10.1890/08-0851.1>
- Pianka ER, Pianka HD (1976) Comparative ecology of twelve species of nocturnal lizards (Gekkonidae) in the Western Australian desert. *Copeia* 1976: 125–142. <https://doi.org/10.2307/1443783>
- Pulliam HR, Danielson BJ (1991) Sources, sinks, and habitat selection: a landscape perspective on population dynamics. *The American Naturalist* 137: S50–S66. <https://doi.org/10.1086/285139>
- Russell AP, Delaunoy M-J (2017) Left in the dust: differential effectiveness of the two alternative adhesive pad configurations in geckos (Reptilia: Gekkota). *Journal of Zoology* 301: 61–68. <https://doi.org/10.1111/jzo.12390>
- Salvadio S, Oneto F (2008) Density regulation in the Mediterranean leaf-toed gecko *Euleptes europaea*. *Ecological Research* 23: 1051–1055. <https://doi.org/10.1007/s11284-008-0465-5>

- Salvadio S, Lanza B, Delaunoy MJ (2010) *Euleptes europaea* (Gené, 1839). *Fauna d'Italia- Reptilia*. Edizioni Calderini de Il Sole 24 ORE, Milano, 869 pp. <https://hal.science/hal-03712438>
- Wiedersheim R (1876) Zur Anatomie und Physiologie des *Phyllodactylus europaeus* mit besonderer Berücksichtigung des Aquaeductus vestibuli der Ascalaboten im Allgemeinen. Zugleich als zweiter Beitrag zur Inselfauna des Mittelmeeres. *Morphologisches Jahrbuch* 1: 495–534.

Supplementary material 1

Euleptes europaea on various plant items

Authors: Daniele Salvi, Emanuele Berrilli, Giacomo Bruni, Matteo Garzia, Veronica Gomes, Giacomo Radi, Michel-Jean Delaunoy

Data type: figure (JPG image)

Explanation note: *Euleptes europaea* on *Rosmarinus officinalis* (a; locality Port-Cros island, Provence; photo by M. Delaunoy); on *Eucalyptus* (b; locality Giglio island, Tuscany; photo by S. Fattorini); on *Lotus cytoides* (c; locality Gabinière islet, Provence; photo by M. Delaunoy); on *Halimione portulacoides* (d; locality Giraglia island, Corsica; photo by M. Delaunoy); on *Juniperus phoenicea* (e; locality Punta Ala, Tuscany; photo by G. Radi).

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/herpetozoa.36.e103465.suppl1>

Supplementary material 2

Observations of arboricolous behaviour of *Euleptes europaea*

Authors: Daniele Salvi, Emanuele Berrilli, Giacomo Bruni, Matteo Garzia, Veronica Gomes, Giacomo Radi, Michel-Jean Delaunoy

Data type: .docx / table

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/herpetozoa.36.e103465.suppl2>

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Herpetozoa](#)

Jahr/Year: 2023

Band/Volume: [36](#)

Autor(en)/Author(s): Salvi Daniele, Berrilli Emanuele, Bruni Giacomo, Garzia Matteo, Gomes Veronica, Radi Giacomo, Delaugerre Michel-Jean

Artikel/Article: [The secret life of a rock-dweller: arboreal acrobatics observed in the European leaf-toed gecko Euleptes europaea 135-141](#)