

Neue Formen in den „Musci Frondosi Archipelagi Indici et Polynesiaci“.

XI. Serie. (1928.)

Von Max Fleischer.

501. ***Sphagnum Junghuhnianum*** var. ***gedeanum*** n. f. ***roseum***. Caespites dense vel altiores laxique, capitula rosea vel roseo-subfusca colorata.
507. ***Pinnatella Kühliana*** n. f. ***coarctata***. A forma typica differt: statura minore, planta 1—2 cm alta, dense ramosa, rami coarctati.
508. ***Sympyodon Perrottetii*** n. f. ***robusta***. A forma genuina statura robustiore diversa; caulis secundarius dense pinnatim ramulosus.
515. ***Thuidium glauceinoides*** n. f. ***pendulum***. Planta tenera, caulis pendulus flexuosus, laxe pinnatus.
521. ***Thuidium dodabettense*** Flsch. n. sp. Dioicum. A specie *T. cymbifolio* Bryol. jav. affini atque valde simili foliis pinnularum ovato-lanceolatis, minus concavis, longius acuminatis, dorsoque grosse papillosum, nervo dorso carinato, apicem versus remote spinuloso diagnosticum. Cetera ut in *T. cymbifolio*, sporogonium ignotum.
531. ***Trismegistia rigida*** n. f. ***viridis***. A forma genuina colore flavoviridiore diversum.
534. ***Warburgiella hygrophila*** n. f. ***minor***. A forma typica differt statura minore, caespite densiore.
535. ***Sematophyllum caespitosum*** n. f. ***monsuniense***. A forma typica americana differt statura plus minusve humiliore, seta breviore.
540. ***Ectropothecium hyalinum*** n. f. ***rufescens***. Plantae plus minusve pulchre aureo-rufulae, nitidae.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Hedwigia](#)

Jahr/Year: 1929

Band/Volume: [69_1929](#)

Autor(en)/Author(s): Fleischer Max

Artikel/Article: [Neue Formen in den "MUSCI FRONDOSI ARCHIPELAGI INDICI ET POLYNESIACI". 94](#)