

**First record of *Aphanus rolandri* (LINNAEUS, 1758) in Albania (Hemiptera:
Heteroptera: Rhyparochromidae: Rhyparochrominae: Gonianotini)**

**Erstnachweis von *Aphanus rolandri* (LINNAEUS, 1758) in Albanien (Hemiptera:
Heteroptera: Rhyparochromidae: Rhyparochrominae: Gonianotini)**

TORSTEN VAN DER HEYDEN

Abstract:

The first record of *Aphanus rolandri* (LINNAEUS, 1758) in Albania is reported. Additional information on the distribution of this species is given.

Zusammenfassung:

Der Erstnachweis von *Aphanus rolandri* (LINNAEUS, 1758) in Albanien wird gemeldet. Ergänzende Informationen zur Verbreitung dieser Art werden gegeben.

Key words: *Aphanus rolandri*, Rhyparochromidae, Heteroptera, Hemiptera, distribution, first record, Albania

Introduction

The genus *Aphanus* LAPORTE, 1833 belongs to the family Rhyparochromidae AMYOT & SERVILLE, 1843 – established by HENRY (1997) – within the suborder Heteroptera of the Hemiptera. 22 species within this genus have been described (DELLAPÉ & HENRY 2018).

Aphanus rolandri (LINNAEUS, 1758) is an unmistakable black ground bug with a characteristic orange or red spot at the base of the wing membrane and a length of 6 to 8 mm.

The species has an Euro-Mediterranean and Pontic distribution. It has been reported from the following European countries: Austria, Belgium, Bosnia and Herzegovina, Bulgaria, Croatia, Cyprus, the Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Italy, Latvia, Liechtenstein, Luxembourg, Macedonia, Malta, Moldova, Montenegro, the Netherlands, Norway, Poland, Portugal (including the Azores and Madeira), Romania, Russia, Serbia, Slovakia, Slovenia, Spain (including the Canary Islands), Sweden, Switzerland, Turkey, Ukraine and the United Kingdom (including the Channel Islands) (HOBERLANDT 1952; PÉRICART 1998, 2001; PROTIĆ 2001, 2016; LEZHENA et al. 2009; ÖNCÜL ABACIGİL et al. 2010; SKIPPER & TOLSGAARD 2013; HANSEN et al. 2015). Furthermore, *A. rolandri* has been reported from Algeria, Armenia, Azerbaijan, Egypt, Georgia, Iran, Israel, Jordan, Libya, Morocco, Syria, Tunisia and Turkmenistan (PÉRICART 1998).

First record of *Aphanus rolandri* (LINNAEUS, 1758) in Albania

On 30.06.2017, ALEKSANDER GOLEMAJ was able to photograph a specimen of *A. rolandri* in the city of Vlorë in southern Albania (Fig. 1).

As *A. rolandri* has not been reported from Albania in scientific publications yet, the record mentioned in this paper is the first one for this country.

Acknowledgements:

I would like to thank ALEKSANDER GOLEMAJ (Vlorë, Albania) for allowing me to use his photograph of *A. rolandri* to illustrate this paper.

References:

- DELLAPÉ, P. M. & HENRY, T. J. (2018): Lygaeoidea Species File. Version 5.0/5.0. (Online database: <http://lygaeoidea.speciesfile.org>). Genus *Aphanus* LAPORTE & DE CASTELNAU, 1832. Available from: <http://lygaeoidea.speciesfile.org/Common/basic/Taxa.aspx?TaxonNameID=1213299>. (Accessed: 06.01.2018).
- HANSEN, M. D. D., OLSEN, K. & SECHER JENSEN, T. (2015): Nye arter i Danmark. Terrestriske arthropoder og vertebrater. Naturhistorisk Museum, Aarhus, 90 pp.
- HENRY, T. J. (1997): Phylogenetic Analysis of Family Groups within the Infraorder Pentatomomorpha (Hemiptera: Heteroptera), with Emphasis on the Lygaeoidea. – Annals of the Entomological Society of America **90**, 275-301.

- HOBERLANDT, L. (1952): On some Hemiptera-Heteroptera of Cyprus. – Acta Entomologica Musei Nationalis Pragae **28 (403)**, 109-116.
- LEZHENINA, I. P., GRAMMA, V. N., TCHAPLIGINA, A. B. & SAVINSKAYA, N. A. (2009): Arthropoda in nests of *Muscicapa albicollis* in highland oak-grove (Kharkiv Region). – Belgorod State University Scientific bulletin, Natural Sciences **3 (58)**, Issue **8**, 95-100. [In Russian.]
- ÖNCÜL ABACIGİL, T., VARLI, S.V. & TEZCAN, S. (2010): Edremit (Balıkesir) Körfezi çevresindeki zeytin bahçelerinde kışlak tuzaklarla saptanan Heteroptera türleri. – Türkiye Entomoloji Dergisi **34**, 105-115.
- PÉRICART, J. (1998): Faune de France. France et régions limitrophes. – 84 C –. Hémiptères Lygaeidae euro-méditerranéens. Volume 3. Systématique: troisième partie. Rhyparochrominae (2). Fédération Française des Sociétés de Sciences Naturelles, Paris, vi + 487 pp.
- PÉRICART, J. (2001): Family Lygaeidae Schilling, 1829 – Seed-bugs. In: AUKEEMA, B. & RIEGER, C. (eds.): Catalogue of the Heteroptera of the Palaearctic Region. Vol. 4. Pentatomomorpha I. Netherlands Entomological Society, Amsterdam, xiv + 346 pp.
- PROTIĆ, L. (2001): Distribution of Heteroptera in various biomes of Serbia. – Acta entomologica serbica **6**, 1-24.
- PROTIĆ, L. (2016): Checklist of Heteroptera of Montenegro. – Ecologica Montenegrina **7**, 350-393.
- SKIPPER, L. & TOLSGAARD, S. (2013): Danmarks tæger - en oversigt. In: SKIPPER, L.: Danmarks blomstertæger. Danmarks Dyreliv, Vol. 12. Apollo Booksellers, Vester Skerninge, 407 pp.

Address of the author:

Torsten van der Heyden, Immenweide 83, D-22523 HAMBURG, e-mail tmvdh@web.de



Fig. 1: *Aphanus rolandri* (LINNAEUS, 1758) (Hemiptera: Heteroptera: Rhyparochromidae),
Vlorë, Albania, 30.06.2017. (Photograph: ALEKSANDER GOLEMAJ.)

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Heteropteron - Mitteilungsblatt der Arbeitsgruppe
Mitteleuropäischer Heteropterologen](#)

Jahr/Year: 2018

Band/Volume: [51](#)

Autor(en)/Author(s): Heyden Torsten van der

Artikel/Article: [First record of Aphanus rolandri \(LINNAEUS, 1758\) in Albania
\(Hemiptera: Heteroptera: Rhyparochromidae: Rhyparochrominae: Gonianotini\)
Erstnachweis von Aphanus rolandri \(LINNAEUS, 1758\) in Albanien \(Hemiptera:
Heteroptera: Rhyparochromidae: Rhyparochrominae: Gonianotini\) 32-33](#)