



THE LIFE CYCLE AND NYMPHAL FEEDING OF *CAPNIONEURA PETITPIERREAE* AUBERT, 1961 (PLECOPTERA, CAPNIIDAE)

Daniel Navarro-Martínez, Manuel J. López-Rodríguez & J. Manuel Tierno de Figueroa*

Departamento de Biología Animal, Facultad de Ciencias, Universidad de Granada, 18071, Granada, Spain

*Author for correspondence: J.M. Tierno de Figueroa, E-mail: jmtdef@ugr.es

ABSTRACT

The life cycle and nymphal feeding of *Capnioneura petitpierreae* Aubert in a temporary stream in the Southern Iberian Peninsula was studied. This collector-gatherer species had a univoltine life cycle with fast growth. In fact, nymphs are only present during five months, from November to March, showing remarkable size variation in all months. The life cycle is compared with that of other Plecoptera from temporary streams and with other species of Capniidae.

Keywords: Plecoptera, stonefly, nymphal biology, temporary stream, Spain

INTRODUCTION

Feeding habits and growth of nymphs are often highly variable among Plecoptera, even within the same genus or family, depending on environmental conditions. Thus, trophic and developmental ecology of stonefly nymphs cannot always be inferred from general patterns at higher taxonomic levels (Stewart & Stark 1993). Studies of these topics are not very numerous, and they are particularly scarce in the European and North African Mediterranean area, especially in temporary streams from Southern Europe. These streams present a pronounced seasonality (summers with low or no water flow) that highly conditions the life cycle strategies of the species living in them.

In stoneflies, only the life cycle and sometimes feeding habits of a few species have been studied in temporary waters (e.g. Agüero-Pelegrín & Ferreras-Romero 2002; López-Rodríguez & Tierno de Figueroa 2005, 2006). Thus more studies are needed to enhance our understanding of the Plecoptera nymphal adaptations to this particular habitat. Because no

studies exist on life cycle and nymphal feeding in Capniidae from European Mediterranean temporary waters, our objective is to elucidate these aspects of the *Capnioneura petitpierreae* Aubert biology. This species is only present in some streams from the Southern Iberian Peninsula (Cádiz and Málaga provinces) and from northern Africa (Tierno de Figueroa et al. 2003). Limited biological data are available on the flight period of Spanish populations (Tierno et al. 1996) and nymphal feeding of a population from Morocco (Azzouz & Sánchez-Ortega 2000).

MATERIALS AND METHODS

Nymphs were collected in Bolaje stream, a tributary of the Genal River, in the Serranía de Ronda (Málaga, Spain), U.T.M. coordinates 30SUF085545, 680 m. The study stream has a temporary water regime with low flow during July and August when only a few pools remain. Monthly samples were taken from February 2006 to January 2007, but no nymphs were taken from April through October.

Collections were made using a kick net with mesh size of 300 μm and specimens were preserved in 70% ethanol. Water temperature, measured during each sample period, varied between 8 and 13.5° C.

Growth data were obtained by measuring pronotal width and outer, right hind femoral length. Measurements were made using an ocular micrometer on an Olympus binocular microscope at 30-40x magnification. In the study of nymphal feeding, 200 individuals (40 per sample period) were randomly selected. Gut contents were analyzed using the methodology of Bello & Cabrera (1999) as modified by, for example, Tierno de Figueroa & Sánchez-Ortega (1999, 2000), Derka et al. (2004) and López-Rodríguez & Tierno de Figueroa (2006). Nymphs were placed in a vial with Herwitg's solution (a variation of Hoyer's media) and

maintained in an oven at 65° C for 24 hours. Afterwards, cleared individuals were collocated on a glass slide and studied with an Olympus microscope at 40x magnification to estimate absolute content percentage (measured as percentage of occupied area), and at 400x magnification to estimate relative content percentage of each component.

A Kolmogorov-Smirnov normality test was used to evaluate distribution of life cycle data. Non-parametric tests were used due to the non-normality of data. Spearman R correlations between femur length and pronotum width were established to represent nymphal growth. For feeding data we determined mean, standard deviation, minimum and maximum, median and quartile range for each component in the diet.




Fig. 1. Life cycle of *Capnioneura petitpierreae*. Box width is proportional to the number of nymphs.

RESULTS AND DISCUSSION

A total of 412 *C. petitpierreae* nymphs were collected from February to March 2006 and from November 2006 to January 2007. Pronotal width and femoral length were highly correlated ($R = 0.96$; $p < 0.05$), consequently only pronotal width was used for construction of graphs (Figs. 1-2). The data indicate nymphal growth extends over five months, with egg hatching beginning in October, after a probable spring-summer diapause. The remarkably high size variability (reflecting asynchronous growth) for a given sample date is typical of reports for other

aquatic insect populations in temporary water (Dieterich & Anderson 1995). Most full grown nymphs were found in February-March samples but a few were found in each sample which probably accounts for the extended flight period reported for this species in this area (Tierno et al. 1996). Thus, *C. petitpierreae* shows a univoltine life cycle typical for most Capniidae (Hynes 1976; Stewart & Stark 1993), and following Hynes (1970) it would be classified as a fast seasonal species due to the short developmental period and extended diapause.

The life cycle of *C. petitpierreae* is similar to that




Fig. 2. Box plots representing *Capnionera petitpierreae* life cycle.

reported for other univoltine species from temporary waters, such as *Rhabdiopteryx christinae* Theischinger and *Nemoura lacustris* Pictet (López-Rodríguez & Tierno de Figueroa 2005, 2006), but it is also similar to that of other *Capnionera* species found in permanent waters of southern Spain (Sánchez-Ortega & Alba-Tercedor 1990). *Capnionera mitis* Despax, for example, occurs in permanent streams but shows a

univoltine fast cycle with extended flight period (Sánchez-Ortega & Alba-Tercedor 1990; Tierno de Figueroa et al. 2001). However, this does not appear to be a general pattern for all *Capnionera* species because *C. brachyptera* Despax is known to have a two year life cycle with slow growth in the first year followed by more rapid growth in the second summer (Lavandier 1975).

	Valid N	Mean	Std.Dev.	Minimum	Maximum	Median	Quartile Range
% abs	200	31.79	26.56	0	90	30	46
% detritus	151	67.23	28.20	5	100	70	45
% diatoms	151	20.26	25.31	0	95	8	30
% hyphae	151	5.42	12.03	0	80	1	5
% spores	151	6.46	14.37	0	85	1	4
% phyllidia	151	0.02	0.24	0	3	0	0
% pollen	151	0.22	0.54	0	3	0	0

Table 1. Gut content composition of *Capnionera petitpierreae* nymphs.

Regarding feeding habits, 151 of 200 individuals examined for diet had gut content (Table 1), and based on the gut composition, this population of *C. petitpierreae* can be considered as a gatherer-collector (feeding mainly on detritus and, to a lesser extent, on diatoms). Other Capniidae are often listed as shredder species (e.g. Tachet et al. 2000), or as shredders and collectors (e.g. Hynes 1976) and the Morocco population of this species studied by Azzouz & Sánchez-Ortega (2000) had a slightly higher quantity of CPOM than was found in our study. This suggests, as does many other studies, that multiple studies on multiple species and multiple populations within species are necessary to gain a more complete understanding of aquatic insect feeding and life cycle adaptations. No general patterns can be applied to particular species living in streams with different biotic and abiotic conditions. The Capniidae, in particular exhibit great diversity in different aspects of life history (e.g. oviparity vs ovoviviparity; lotic vs lentic habitat; univoltinism vs semivoltinism, etc.) showing it is a very heterogeneous group.

ACKNOWLEDGMENTS

The authors want to thank the valuable comments of doctors Ignac Sivec, Bill Stark and John Brittain that notably improved the original manuscript.

REFERENCES

- Agüero-Pelegri, M. & M. Ferreras-Romero. 2002. The cycle of *Guadalgenus franzi* (Aubert, 1963) (Plecoptera: Perlodidae) in the Sierra Morena Mountains (Southern Spain): Semivoltinism in seasonal streams of the Mediterranean Basin. *Aquatic Insects*, 24:237-245.
- Azzouz, M. & A. Sánchez-Ortega. 2000. Feeding of the nymphs of nine stonefly species (Insecta: Plecoptera) from North Africa (Rif Mountains, Morocco). *Zoologica Baetica*, 11:35-50.
- Bello, C.L. & M.I. Cabrera. 1999. Uso de la técnica microhistológica de Cavender y Hansen en la identificación de insectos acuáticos. *Boletín Entomológico Venezolano*, 14:77-79.
- Derka, T., J.M. Tierno de Figueroa & I. Krno. 2004. Life cycle, feeding and production of *Isoptena serricornis* (Pictet, 1841) (Plecoptera, Chloroperlidae). *International Review of Hydrobiology*, 89:165-174.
- Dieterich, M. & N.H. Anderson. 1995. Life cycles and food habits of mayflies and stoneflies from temporary streams in western Oregon. *Freshwater Biology*, 43:47-60.
- Hynes, H.B.N. 1970. *The ecology of running waters*. Liverpool University Press, Liverpool.
- Hynes, H.B.N. 1976. *Biology of Plecoptera*. *Annual Review of Entomology*, 21:135-153.
- Lavandier, P. 1975. Cycle biologique et production de *Capnionera brachyptera* D. (Pléoptères) dans un ruisseau d'altitude des Pyrénées centrales. *Annales de Limnologie*, 11:145-156.
- López-Rodríguez, M.J. & J.M. Tierno de Figueroa. 2005. Ciclo de vida y composición de la dieta de *Nemoura lacustris* Pictet, 1865 (Plecoptera, Nemouridae). *Boletín de la Asociación Española de Entomología*, 29:87-97.
- López-Rodríguez, M.J. & J.M. Tierno de Figueroa. 2006. Life cycle and nymphal feeding of *Rhabdiopteryx christinae* Theischinger, 1975 (Plecoptera, Taeniopterygidae). *Annales de la Société Entomologique de France*, 42:57-61.
- Sánchez-Ortega, A. & J. Alba-Tercedor. 1990. Life cycles of some species of Plecoptera in Sierra Nevada (South of Spain). Pages 43-52. *In*. I.C. Campbell [editor]. *Mayflies and stoneflies: life history and biology*. Kluwer Academic Publications, Dordrecht.
- Stewart, K.W. & B.P. Stark. 1993. *Nymphs of North American stonefly genera (Plecoptera)*. University of North Texas Press, Denton.
- Tachet, H., P. Richoux, M. Bournaud & P. Usseglio-Polatera. 2000. *Invertébrés d'eau douce. Systématique, biologie, écologie*. CNRS Éditions, Paris.
- Tierno, J.M., T. Pérez & A. Sánchez-Ortega. 1996. Composición faunística y fenología de los Plecópteros (Insecta: Plecoptera) de la Serranía de Ronda (Málaga). *Boletín de la Asociación Española de Entomología*, 20:47-58.
- Tierno de Figueroa, J.M., J. Luzón-Ortega & A. Sánchez-Ortega. 2001. Fenología de los plecópteros (Insecta, Plecoptera) de Sierra Nevada (Granada, España). *Zoologica Baetica*, 12:49-70.
- Tierno de Figueroa, J.M. & A. Sánchez-Ortega. 1999. Imaginal feeding of certain systellognathan stonefly species (Insecta: Plecoptera). *Annals of the Entomological Society of America*, 92:218-221.
- Tierno de Figueroa, J.M. & A. Sánchez-Ortega. 2000.

Navarro-Martínez, D., M. J. López-Rodríguez & J. M. Tierno de Figueroa 2007. The life cycle and nymphal feeding of *Capnioneura petitpierreae* Aubert, 1961 (Plecoptera, Capniidae). *Illiesia*, 3(8):65-69. Available online: <http://www2.pms-lj.si/illiesia/Illiesia03-08.pdf>

Imaginal feeding of twelve nemouroidean stonefly species (Insecta, Plecoptera). *Annals of the Entomological Society of America*, 93:251-253.

Tierno de Figueroa, J.M., A. Sánchez-Ortega, P. Membiela Iglesia & J.M. Luzón-Ortega. 2003. Plecoptera. 404 pages. *In*. M.A. Ramos et al. [editors]. *Fauna Ibérica*, vol. 22. Museo Nacional de Ciencias Naturales, CSIC, Madrid.

Received 8 May 2007, Accepted 16 July 2007, Published 17 July 2007

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Illiesia](#)

Jahr/Year: 2007

Band/Volume: [03](#)

Autor(en)/Author(s): Navarro-Martínez Daniel, López-Rodríguez Manuel J., Tierno de Figueroa José Manuel

Artikel/Article: [The life cycle and nymphal feeding of *Capnioneura petitpierreae* Aubert, 1961 \(Plecoptera, Capniidae\). 65-69](#)