

Gauge data and water temperatures in the RITRODAT area 1991 and 1992 and long year averages of climatic data.

Gernot BRETSCHKO

Keywords: Discharge, water-temperature, climatic data

Water level is registered with a GANSER water-stage recorder. Staff gauge and recorder are positioned on the right bank below the RITRODAT-bridge.

The hydrological year 1991 was dominated by an extreme flood (30–50 years recurrence) with discharge maxima on the 28th of July and 2nd of August. The highest gauge mark was 120 cm. The actual water level was somewhat higher than that but it surpassed the measuring capacity of the equipment. Waterlevel as means of decades and extremes are set out in Tab. 1.

		Mean	Extremes Max	Min			Mean	Extremes Max	Min
1		32.2	56.0	21.5	1		25.4	48.5	17.5
2	J	23.3	34.0	16.0	2	J	32.8	81.0	14.5
3		12.9	16.0	10.0	3		49.9	120.0	19.0
1		6.5	10.0	1.5	1				
2	F	6.1	8.5	2.0	2	A			
3		9.8	15.0	2.5	3				
1		21.3	42.0	10.0	1		89.2	91.5	88.0
2	M	28.7	45.5	21.5	2	S	88.4	89.0	88.0
3		28.6	43.0	16.0	3		88.4	89.0	87.0
1		24.5	30.0	16.0	1		94.0	100.5	88.0
2	A	22.8	28.5	18.0	2	O	86.8	90.0	86.0
		22.8	39.5	15.5	3		87.2	88.0	86.5
1		32.0	41.0	25.0	1		93.9	104.5	87.5
2	M	38.0	50.0	31.0	2	N	101.3	106.0	99.0
3		43.1	69.0	30.5	3		100.0	102.0	97.5
1		45.3	73.0	33.5	1		93.0	97.5	89.5
2	J	31.4	49.0	22.0	2	D	89.7	101.5	87.5
3		24.6	49.5	17.0	3		122.5	191.5	101.5
ANNUAL			MEAN		MAX			MIN	
-			(21.8)		(120)			2	
MONTH/DATE			-		VII/3			II/1	

Table 1: Water level means of decades and extremes, 1991. Gauge station RITRODAT-bridge. August data are missing because of extreme flood. Values in brackets: approximations only.

Gernot Bretschko




Figure 1: RITRODAT study area. Shore line is bankfull line (gauge RITRODAT-bridge: 70 cm). Hatched areas: deeper than 25 cm Black areas: higher than 25 cm relative to gauge station.

The summer flood changed greatly the topography of the channel (Fig. 1). Severe erosions happened on the right bank between profile 7 and 14. In the absence of the fortified dam, the brook would have changed its entire course towards north, migrating into the valey. Aggregations occurred on the right bank between profiles 0 and 4 and in the left part of the channel between profiles 0 and 15 (Fig. 2). The effects of these flood induced aggregation/erosion processes are very obvious in the development of the mean longitudinal profile (Fig. 3).

Because of these topographic changes, a new rating curve had to be established for the gauge RITRODAT-bridge. In the praeflood situation (1989) one typical power function describes the regression between water level and discharge (Fig. 4):

$$y = 0.00018 x^{2.584} \quad r^2 = 0.988 \quad n = 11$$

After the flood two power functions are necessary to describe the regression with the necessary accuracy (Fig. 4):

gauge reading lower than 135 cm:

$$y = 3.266^{-18} x^{8.474} \quad r^2 = 0.946 \quad n = 18$$

gauge reading higher than 135 cm:

$$y = 1.045^{-9} x^{4.440} \quad r^2 = 0.992 \quad n = 10$$

Gauge data and water-temperatures




Figure 2: RITRODAT-study area. Cross-sectional profiles, 1991-08-27. Reference: Gauge RITRODAT-bridge, 150 cm. Open areas show erosions and hatched areas show aggregations in comparison to the praeflood situation (1990-10-17, Fig. 1).

Gernot Bretschko




Figure 3: RITRODAT- study area. Mean level of cross-sectional profiles at mean discharge. Reference: Gauge RITRODAT-bridge, 150 cm. Hatched areas: 95 % Confidence intervals (from BRETSCHKO, in press).




Figure 4: Gauge RITRODAT-bridge, rating curves. Broken line: pre-flood condition (1989). Full curve: after-flood condition (1992).

Gauge data and water temperatures

The deep erosion in the area of the gauge station made it necessary to elongate the rod gage by 100 cm downwards. Based on equal discharges a regression has been established between the prae- and afterflood gauge readings (Fig. 5):

$$y = -47.726 + 0.662 x \quad r^2 = 0.999 \quad n = 17$$

Because of the linearity of the regression, the difference between the two gauge readings is constant and amounts

$$\underline{\underline{81.5 \text{ cm}}}$$

The water level data for 1992 are shown in Table 2.




Figure 5: Gauge RITRODAT-bridge. Regression between prae- (ordinate) and afterflood (abscissa) gauge readings. Based on equal discharges.

The temperature of the surface water is measured with a Pt-100-probe at the upstream end of the RITRODAT-study area (0C1↑1) and continuously registered with a SCHENK analog recorder. The data (means of decades and extremes) are set out in table 3.

Since the establishment of the Biological Station Lunz (1906) various climatic data have been collected continuously. So far, a 32-years average (1927–1959) is used to characterize the climatic conditions in Lunz. A more recent 30-years average (1960–1989) is set out in table 4.

For the compilation of the annual data I have to thank Mr. A. AIGNER.

Gernot Bretschko

1991

MEAN					EXTREMES		MEAN					EXTREME		
					MAX	MIN						MAX	MIN	
1	J	5.1	5.5	4.7	1	J	8.1	9.7	6.8	1	A	7.6	9.0	6.8
2		4.4	5.4	3.3	2		8.1	10.4	6.7	2		9.0	10.7	7.9
3		3.8	5.6	2.9	3		7.4	9.3	6.6	3		10.0	12.1	7.7
1	F	2.4	4.5	0.6	1	S	9.3	12.7	8.0	1	O	8.5	10.8	7.3
2		2.2	4.1	0.4	2		9.0		8.2	2		8.5		7.0
3		3.9	5.2	2.6	3		9.2	12.7	8.2	3		5.8	9.6	3.7
1	M	4.9	5.8	3.6	1	N	6.0	7.4	3.4	1	D	3.8		1.5
2		5.5	6.1	4.8	2			9.7		2		6.1		5.2
3		5.8	6.9	4.6	3			9.6		3		6.0	7.0	5.4
1	A	6.0	7.0	5.1	1	J	6.7	8.0	6.1	1	J	7.5	9.1	6.4
2		5.8	7.2	3.9	2		7.5	9.1	6.4	2		2.8		1.6
3		5.8	7.1	3.5	3		7.9	9.4	6.8	3		4.6		3.2
1	M	6.5	7.5	5.7	1	D	3.8		1.5	1	D	3.8		1.5
2		6.2	7.2	4.9	2					2		2.8		1.6
3		6.3	7.2	5.6	3					3		4.6		3.2
1	J	6.7	8.0	6.1	1	D	3.8		1.5	1	D	3.8		1.5
2		7.5	9.1	6.4	2					2		2.8		1.6
3		7.9	9.4	6.8	3					3		4.6		3.2
ANNUAL:					MEAN		MAX					MIN		
-					6.3		12.7					0.4		
MONTH/DECADE:					-		IX/1					II/2		

1992

MEAN					EXTREMES		MEAN					EXTREME		
					MAX	MIN						MAX	MIN	
1	J	4.6	5.6	3.4	1	J	9.2	10.4	8.4	1	A	10.0	11.4	9.2
2		4.9	5.4	3.8	2		8.8	10.4	7.9	2		10.0	11.3	8.8
3		4.4	5.4	3.2	3		9.7	11.2	8.4	3		10.0	11.2	9.1
1	F	4.7	5.6	3.7	1	S	8.5	10.0	7.3	1	O	7.8	8.2	7.2
2		4.7	5.6	3.7	2		9.0	9.7	8.3	2		7.4	8.3	6.5
3		4.6	5.7	3.3	3		9.0	9.7	7.3	3		6.8	7.8	6.2
1	M	5.2	6.2	4.3	1	N	6.6	7.1	6.1	1	D	5.9	6.5	5.3
2		4.9	6.2	3.9	2		9.0	9.7	8.3	2		6.0	7.0	5.1
3		5.5	6.6	4.5	3		9.0	9.7	7.3	3		6.3	6.9	5.1
1	A	6.2	7.1	5.0	1	J	7.4	9.0	6.5	1	D	5.9	6.5	5.3
2		6.3	7.3	5.2	2		7.5	9.4	6.5	2		5.3	5.9	4.4
3		6.7	7.7	5.6	3		8.5	10.3	7.6	3		4.3	5.7	2.7
1	M	6.7	7.9	6.2	1	N	6.6	7.1	6.1	1	D	5.9	6.5	5.3
2		6.8	7.8	6.0	2		6.0	7.0	5.1	2		6.0	7.0	5.1
3		7.3	8.7	5.9	3		6.3	6.9	5.1	3		6.3	6.9	5.1
1	J	7.4	9.0	6.5	1	D	5.9	6.5	5.3	1	D	5.9	6.5	5.3
2		7.5	9.4	6.5	2		5.3	5.9	4.4	2		5.3	5.9	4.4
3		8.5	10.3	7.6	3		4.3	5.7	2.7	3		4.3	5.7	2.7
ANNUAL:					MEAN		MAX					MIN		
-					6.9		11.4					2.7		
MONTH/DECADE:					-		VIII/1					XII/3		

Table 3: Temperatures of surface water (means of decades and extremes): measured at the upstream end of the RITRODAT-study area (0C1†1).

Gauge data and water-temperatures

T E M P E R A T U R E				
	1927/59		1960/89	
Annual mean	6.36.6 ± 0.4			
Mean Maximum	33.232.8 ± 1.6			
Mean Minimum	– 25.1– 23.0 ± 5.0			
	1927/59		1960/89	
	First	Last	First	Last
hoar-frost	7.X.	3.V.	27.IX.	24.V.
Frost	13.X.	4.V.	27.IX.	29.V.
Snow cover	9.XI.	9.IV.	19.X.	5.V.
Snow fall	1.XI.	–	8.X.	11.V.
Max. snow cover (cm)	150 (1988–03–13)			

P R E C I P I T A T I O N		
	1927/59	1960/89
Annual mean total (mm)	1629.01514.8 ± 65.9	
Mean highest daily total	72.062.4	

Table 4: 30-years averages of climatic data (1927/59 and 1960/89). Biological Station Lunz (615 m asl; 47°51' N; 15°04' E).

References:

BRETSCHKO, G., in press: Niederschlagsereignisse, Hochwässer und Fließgewässerökologie.– Wiener Mitteilungen.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Jahresbericht der Biologischen Station Lunz](#)

Jahr/Year: 1993

Band/Volume: [1991-92_014](#)

Autor(en)/Author(s): Bretschko Gernot

Artikel/Article: [Gauge data and water temperatures in the RITRODAT area 1991 and 1992 and long year averages of climatic data. 41-47](#)