## Upper Eocene to Early Miocene Planktonic Foraminifera from the Subsurface Sediments in Cauvery Basin, South India\*)

By D. S. N. RAJU

With 13 plates, 14 figures, 1 table

Palaeontology Laboratory, O. N. G. Commission, Baroda-9, India

## Abstract

Fifty species and subspecies of planktonic foraminifera, belonging to the genera Globigerina, Globigerinoides, Globigerinita, Globigerapsis, Globoquadrina, Globorotalia, Globigerinopsis, Pseudohastigerina, Hantkenina, Cribrohantkenina, Cassigerinella and Chiloguembelina, are systematically described, for the first time, from the Upper Eocene, Oligocene and early Miocene subsurface sediments in Cauvery basin, South India. Two new species, Globigerina sastrii and Globigerinopsis gubai, are described.

Seven planktonic foraminiferal zones are proposed for the Upper Eocene to early Miocene sequence. The zones, in order from bottom to top, are: Globigerapsis mexicana zone, Globorotalia cerroazulensis zone, Globigerina gortanii zone, Globigerina sastrii zone, Globigerina ampliapertura zone, Globigerina angulisuturalis zone and Globorotalia kugleri/ Globigerinoides primordius zone.

The significance of these findings in correlation of the sections in Cauvery basin with those from other tropical regions is discussed. The Eocene — Oligocene limit is placed at the top of the Globorotalia cerroazulensis zone. The Oligocene — Miocene limit is placed at the lower boundary of the Globorotalia kugleri/Globigerinoides primordius zone.

## Introduction

The last two decades are marked by a stupendous increase in our knowledge on planktonic foraminifera and a rapid growth of their importance in biostratigraphic classification and intercontinental correlations. Researches of BOLLI (1957) and several others in Caribbean region have led to an elaborate planktonic foraminiferal zonal classification of the Cretaceous and Cenozoic marine sediments. These works were followed by attempts to recognize the Trinidad zones in other parts of the world. Much of additional knowledge on the mid-Tertiary planktonic foramini-

<sup>\*)</sup> Published with the permission of the Director of Geology, Oil & Natural Gas Commission, Debradun, India.

The views expressed in the paper are of the author and not necessarily those of the O. N. G. Commission.

fera was brought out by BLOW & BANNER (1962) and BLOW (1969) from sections in East Africa and other tropical regions.

Some preliminary studies carried out during the last six years on Indian localities have indicated the presence of rich assemblages of planktonic foraminifera. However, most of the species were not systematically described and the knowledge on stratigraphic distribution of already known species is very limited.



The present study is an attempt to systematically describe the Upper Eocene, Oligocene and early Miocene planktonic foraminifera from the subsurface sections in Cauvery basin, South India; to define a sequence of biostratigraphic zones and to evaluate their value in correlation with sections outside this region.

## Previous work

The Upper Eocene to Lower Miocene planktonic foraminifera are known in Cauvery basin only from the deep exploration wells drilled by Oil & Natural Gas Commission since 1964. The occurrence of some diagnostic species was reported by RAJU (1966). Later, RAJU (1967, 1968) recognized some planktonic foraminiferal zones in the Late Cretaceous to Oligocene subsurface sediments. However, the planktonic species were not systematically described so far. The knowledge on the mid-Tertiary planktonic foraminifera from other sedimentary basins in India is very limited.

Present work

The present study is essentially limited to the core samples. Data from the following intervals of the deep wells from Cauvery basin constitute the main theme of the present study.

Well No.	Depth interval	Number of cores studied
KKL-1	8001600 m.	7
KKL-2	1000—1530 m.	6
KKL-4	8001800 m.	15
KKL-5	1000—1800 m.	5
KKL-6	1000—1800 m.	4
NGT-1	1600—2000 m.	2
Madanam-1	800—1300 m.	3
TPD-1	450— 950 m.	5

The following abbreviations, KKL for Karaikal, NGT for Nagapattinam, TPD for Tirupundi, are used in this paper (see fig. 1).

One to six samples are studied from each core depending on the length recovered. Besides these almost all the cutting samples representing every three metres interval of the well are also studied. There are considerable gaps between the cores and as such the complete distribution of many of the species could not be satisfactorily determined during this study. Some data from well cutting samples are incorporated in preparing the range chart, only to indicate last occurrences (extinctions) of some diagnostic species.

## Resume of the general stratigraphy of the Cauvery basin

The oldest sediments known, which overlie the Archaean metamorphics in Cauvery basin, are the so-called "Upper Gondwanas" exposed as small patches along the western fringes of the basin. They are essentially lacustrine to brackish water deposits of ? Jurassic to Lower Cretaceous age. Overlying the "Upper Gondwanas" are the well known Cretaceous (Albian to Maestrichtian) sediments, which are widely distributed in the Cauvery basin (see Figure 1). These marine Cretaceous sediments were the source of many palaeontological works during the last 90 years.

The Cretaceous rocks are overlain by marine Paleocene-early Eocene sediments particularly in the Pondicherry and Trichinopoly areas. The Paleocene-early Eocene sediments are overlain, on the surface, by continental deposits known as "Cuddalores" of Neogene age. Large areas in the eastern part of the basin are covered by Alluvium or laterites.

Marine sediments younger than the Lower Eocene are known only in the subsurface, deep well sections. Recent Petroleum exploration activities, geophysical surveys and deep drilling by Oil & Natural gas Commission, have added considerable knowledge on the subsurface stratigraphy and structure of this basin. For some of these details reference is made to RAMANATHAN (1968).

#### Acknowledgements

The author is thankful to Prof. Dr. H. KÜPPER, Director, Dr. M. E. SCHMID, micropaleontologist and other staff members of the Geological Survey of Austria; to Shri D. K. GUHA, Senior Geologist, Shri C. K. R. SASTRY, Chief Geologist, Shri S. N. TALUKDAR, Additional Director of Oil & Natural Gas Commission, India for providing various facilites during the study in Vienna (Austria) and Baroda (India). He is thankful to Prof. H. M. BOLLI, Zürich (Switzerland) for checking and commenting on many of the species described in this paper. He is grateful to Prof. A. PAPP, University of Vienna; Dr. P. MARKS and Prof. C. W. DROOGER of the University of Utrecht (Holland) for many suggestions. He is also thankful to Dr. C. G. ADAMS for providing facilities to examine some of the types deposited in British Museum (Natural History), London.

## Zonation

Most of the Upper Eocene to early Miocene planktonic foraminiferal species, reported from other tropical regions, are now recorded from Cauvery basin. However, the recognition of the zonal boundaries within the subsurface sections in Cauvery basin is hamphered to a great extent due to the absence of continuous core samples. The boundary between the *Globorotalia cerroazulensis* zone and *Globigerina gortanii* zone falls within cored interval of a well. The microfaunal changes at the other zonal boundaries could not be precisely defined. Six of the zonal boundaries are defined here by horizons of extinction of markers, while two are defined by the first appearance level. The horizons of extinction, used here in definition PLATE 18



G. semiinvoluta-G. barri zone, G. cerroazulensis zone and C. cubensis-G. ampliapertura euapertura zone. The present study has resulted in providing additional data from the G. mexicana zone (= G. semiinvoluta zone) and G. cerroazulensis zone, subdivision of "C. cubensis-G. ampliapertura euapertura zone" into three zones and in recognition of two additional zones in the Late Oligocene-early Miocene sequence.

## Globigerapsis mexicana zone

Author: BOLLI, 1957 (given as Globigerapsis semiinvoluta zone).

Reference section: Core between 1904. 2 and 1906. 5 m. inNGT-1.

This zone is defined as the interval from the last occurrence of Truncorotaloides rohri BRÖNNIMANN & BERMUDEZ to the last occurrence of Globigerapsis mexicana (CUSHMAN). A very rich assemblage of planktonic foraminifera is known from this interval in NGT-1. The assemblage includes Globigerapsis mexicana, G. cf. tropicalis BLOW & BANNER, Globigerinatheka barri BRÖNNIMANN, G. cf. lindiensis BLOW & BANNER, Pseudohastigerina micra (COLE), Hantkenina alabamensis CUSHMAN, H. suprasuturalis BRÖNNIMANN, H. trinitatensis BRÖNNIMANN, H. cf. thalmanni BRÖNNIMANN, Hantkenina sp., Globigerina corpulenta SUBBOTINA, G. pseudocorpulenta CHALILOV, G. gortanii praeturritilina BLOW & BANNER, G. angiporoides HORNIBROOK, Chiloguembelina martini (PIJPERS), C. cubensis (PALMER) and Globigerinita spp. A poorly preserved form with doubtful cribrate aperture, referred here to as Cribrohantkenina cf. inflata (HowE) is also found in this zone.

A rich assemblage of larger foraminifera including *Pellatispira* madaraszi, Nummulites cf. fabianii and Discocyclina spp. are recorded from core samples, at a depth of 1230 m. in Madanam-1, belonging to this zone.

The extinction level of T. rohri was marked in almost all the wells drilled in Cauvery basin and found to be a very useful horizon in well to well correlation. The extinction level of G. mexicana was marked in only a few wells so far.

## Globorotalia cerroazulensis zone

Author: BOLLI, 1957 (given as Globorotalia cocoaensis zone).

Reference section: Cores between 1689 and 1720 m. in KKL-4.

This zone is defined as the interval between the last occurrence of Globigerapsis mexicana and the last occurrence of Globorotalia cerroazulensis (COLE). Cribrobantkenina inflata (HOWE), Hantkenina alabamensis, Globorotalia cerroazulensis, G. centralis CUSHMAN & BERMUDEZ and several other species shown in table I are common in this zone. The extinction level of G. cerroazulensis was marked in all the wells drilled near Karaikal, Nagapattinam and Tirupundi and this horizon is one of the most useful levels in well to well correlation of the subsurface sequence and in intercontinental correlation. In Karaikal well-6, the top of G. cerroazulensis zone falls within a core and the foraminiferal assemblages above and below this horizon do not indicate any major paleoenvironmental change.

## Chiloguembelina cubensis assemblage

The present author (RAJU, 1968) proposed, tentatively, the C. cubensis-G. ampliapertura euapertura assemblage zone from the interval from the last occurrence of G. cerroazulensis to the last occurrence of Chiloguembelina cubensis (PALMER). The present study has led to the recognition of three zones within this interval.

## Globigerina gortanii zone

Authors: BLOW & BANNER, 1962, emended here.

Reference section: Core between 1800 and 1803 m. in NGT-1.

This zone is defined as the interval from the last occurrence of G. cerroazulensis to the first appearance of Globigerina sastrii n. sp. The common species in this zone are Globigerina gortanii gortanii (BORSETTI), G. gortanii praeturritilina BLOW & BANNER, G. cf. pseudoampliapertura BLOW & BANNER, G. ampliapertura BOLLI, G. prasaepis BLOW, Globorotalia increbescens (BANDY), Pseudohastigerina barbadoensis BLOW and Cassigerinella chipolensis CUSHMANN and PONTON. Cassigerinella chipolensis and P. barbadoensis occur together even in the very basal part of this zone. No forms referable to Globigerina sellii BORSETTI are found in this zone.

## Globigerina sastrii zone

Author: Proposed here.

Type section: Cores between 1579 and 1586 m. in KKL-4.

This zone is defined by the range of Globigerina sastrii n. sp. The planktonic foraminifera recorded from the type section of this zone include Globigerina sastrii n. sp., G. gortanii gortanii, G. sellii BORSETTI, G. ampliapertura, G. tripartita tripartita KOCH, G. tripartita rohri BOLLI, G. prasaepis, Globorotalia cf. opima opima BOLLI, G. opima nana BOLLI, G. gemma JENKINS, Cassigerinella chipolensis, Chiloguembelina cubensis and other forms. Rare forms of Lepidocyclina (Nephrolepidina) cf. isolepidinoides are also found in this zone in KKL-4 and TPD-1. Globigerina sastrii n. sp. is very common in Cauvery basin and has a short range. However, it may be of local value only as this species was not described from sections outside the Cauvery basin.

## Globigerina ampliapertura zone

Author: BOLLI, 1957, emended.

Reference section: Drill cutting samples from section immediately overlying the G. sastrii zone in KKL-4.

This zone is tentatively defined as the interval from the last occurrence of *Globigerina sastrii* n. sp. to the last occurrence of *Globigerina ampliapertura*. The study of the drill cutting samples indicate that *G. ampliapertura* extincts approximately at the same level as that of *Chiloguembelina cubensis* in Karaikal wells.

## Globigerina angulisuturalis zone

Author: BLOW, 1969, emended.

Reference section: Cores between 1450 and 1500 m. in KKL-2.

This zone is defined as the interval from the last occurrence of Globigerina ampliapertura to the first appearance of Globigerinoides primordius BLOW & BANNER. The planktonic foraminifera recorded from this zone include Globigerina angulisuturalis BOLLI, G. ciperoensis BOLLI, G. angustiumbilicata BOLLI, G. woodi woodi JENKINS, Globorotalia siakensis (LEROY), G. opima nana BOLLI, Cassigerinella chipolensis, Globoquadrina cf. larmeui AKERS and very rare forms referable to Globigerina gortanii gortanii (BORSETTI). Globoquadrina first appears in this zone. Larger foraminifera Lepidocyclina (Eulepidina) sp. and L. (Nephrolepidina) are also recorded within this zone in KKL-2.

## Globorotalia kugleri / Globigerinoides primordius zone

Author: BLOW, 1969, Zone N 4.

Reference section: Cores between 1126 and 1425 m. in KKL-2.

This zone is defined as the interval from the first appearance of Globigerinoides primordius BLOW & BANNER to the last occurrence of Globorotalia kugleri BOLLI. The planktonic species recorded from the lower part of this zone include Globigerina angulisuturalis, G. ciperoensis, Globigerinoides primordius, Globorotalia siakensis, G. opima nana and Globoquadrina cf. larmeui. In the middle part of this zone, though Globigerinoides primordius, Globorotalia siakensis and a few other forms are recorded, the planktonic forms are rare. Again, the planktonic foraminifera are abundant in the upper part of this zone. The assemblage includes Globorotalia kugleri BOLLI, G. siakensis, G. obesa BOLLI, Globoquadrina praedehiscens BLOW & BANNER, G. altispira globosa BOLLI, Globigerinoides cf. altiaperturus BOLLI, Globigerinopsis gubai n. sp., Globigerinita dissimilis (CUSHMAN), G. stainforthi (BOLLI, LOEBLICH & TAPPAN), Globigerina venezuelana HEDBERG and rare forms of Globigerina binaiensis KOCH.

## Globigerinoides trilobus assemblage

Planktonic foraminiferal zones are not successfully recognized so far in the interval above the *G. kugleri/G. primordius* zone in the subsurface sections in Cauvery basin. In the sequence above the *G. kugleri/G. primordius* zone, the larger foraminifera Miogypsinidae and Lepidocyclina are common and planktonics are not adequately represented.

#### FORAMINIFERA IN CAUVERY BASIN

#### TABLE-I THE DISTRIBUTION OF SOME STRATIGRAPHICALLY IMPORTANT MIDDLE ECCENE TO LOWER MICCENE PLANKTONIC

			_		_					_		_		_	_				_	_		_								_							_					_					
BIOST & AT IGRAPHIC ZONES	RÉFERÈNCE SECTIONS Well Nº. / DÉPTHS In Meters	PROBABLE PALEOEMVIRONMENTS	H. Aunblei H. trindatantis	H.aloboma nais V = mirziscinicatie	Cribrohantkening Inflata	1 Etimenerita	G. yegudenáis	6. frontosa A orendocessaritesta	G.corpulerio	G.gortonic praeturritiling	G.gorlani gartanii B. cf. settii	G, south	G. amplio perntura	G. on gusti und Hicata	G. ciperoensis	G.woodi woodi	G. venezuelono			:	G labrati	G.crassata	G. bolivanaria	G.centralis	6. cerradzulensia	6. opiena hana			G. trilobus		G. cf. Harmuk	G. proedehiscens	G. deniscens	G. 40+1\$#44 9150664	Preudohastigerina micrá	Pesudohastiger he borbodosnais	Cassigeriaella ahibolensis	Chiloguembeling mertini	Chiloguembellno cubensis	Globigerinopsis guret	Treacorototáles topliansis	Giobigerapsis kugleri	Globigerapeis menicana	V.I.O.I.O.I.MOTHERA DOIRI	Urbumpiges Deckinging I Clubidarinita die almiks	Glabigerlaita stainfacthi	Truncarotoloides robri
												-		ſ															1								Γ	Γ		Π				Τ	Ι		Γ
G.TRILOBUS ASSEMBLAG	e	MIDDLE TO		1							_				Γ	T				1							_		1					_			П								T	T	
	KKL	OUTER SHELF			L																					_		1	· II			٠	Ц	•			4			•	Ц				÷	÷	
G. KUGLERI-	KKL_2, #28_ #25		$\square$				11	4	1	$\square$	4		$\downarrow$	4	•	$\square$	4	_	1	4	+	1	Ľ		_	4	-	ŧĹ	• 61	$\vdash$		• •	-	-	1	Ĺ	#		Ц	4	Ц	4	4	+	4	÷	L
6. PRIMORDIUS	KKL_2,1353_1349	4.	L				$\perp$		╘						•		•	_		4	1	_			_	4	1	ŧĻ	+			:1	4	∔		┡	₩	Ł	Ľ	4	$\vdash$	_	4	∔	Щ	4	⊢
	KK1_2,14216_1415.6	15 1					$\downarrow$		1	$\square$				ĮĮ,	· ·	•	cf	_	-	4		⊢				Щ	•	ŧļ.		1		4	4	⊥		Ļ.	Щ	L		Ц		4	┛	4	뿌	4	L
	KKL_2,1425.4_1421.6	1 2 2												11		•				⊥		L			_	Ц		L					1				Щ	Ι.	L			┛	┙	1	4	4	L
G.ANGULISUTURALIS	KKL_2,1460.5_1457.5	828	$\square$		+		$\downarrow$		$\perp$	Ц	╇		Ц	Ц	•		1.	_	4	+					_	4	$\downarrow$	4				4	$\downarrow$	+		1	Щ			H		_	$\perp$	4	4	4	L
	KKL-2,1500-5-14945		$\vdash$				+-+			Ц			1	4			H	_	4	$\downarrow$	1	Ļ.				#	4	┥	+	Ŀ	Ц		$\downarrow$	1			4		Ц	Ц		4	4	$\downarrow$	₽	4	L
	KKL_4,1609_1503	5 S	Ц.								⊥			1	·				4				L.			1		1			L		$\downarrow$	1		1	4				Ц	⊢	4	4	4	4	Ļ
G AMPLIAPERTURA	KKL_4,1569_1549	ELF	$\vdash$	H	+	+	┟┤		╉	H	$\frac{1}{1}$	$\frac{1}{1}$	H	∦	╀			+	ł	+	╀	┝	-		1	╢	╉	╈	╀	╉		┥	╉	╞	╉	t	Ħ			Н	H	+	╉	╡	╬	╀	ŀ
	KKL_4,1582_ 1579	15.2		T			T		1	Π	11.	Π	Π	T	Ţ	Π			Т	Т	I	1				П	Т	Т		Τ			T	ŀ	Т	Т	Π	Γ	П	П	П	Т	T		Ti	T	Γ
	KKL_4,1586_1582	] ដ∔ ដ		П			Π		Т	Π	Iŀ	Π	П	Т	Т			Т	Т	Τ	Τ					I		Т				Т	Т	Т	1		Π		П	П	П	Т	Т	Τ	Tt	Æ	ſ
6. GORTANII	NGT_1,1803_1800	15 5					Π	T	1	1	П	1			Τ	Π				Τ				:		Π		Τ									Π		Π	Π	Π		Τ	Τ	T	í T	Γ
	KRL _6,1746 _ 1749	<u> </u>	$\Box$	T		1	1			1		[	!!							Τ						Π	T	1	1		:	T	T	T			T	1	П		$\square$		Τ	Τ	Ti		ſ
-	KKL _ 6, 1749 _1750	]	Ĺ	П	1		1			Ī	?		Π		Ι					Τ	1				Т	I								Τ	Π	•	ş	Π					I	Ι		ī	ſ
	KKL _ 4,1689 _ 1665	] ; ;	T	Ш	Ш		1			I					T							1				Π		T				T			$\Pi$	Ţ		Π		$\Box$				T	Ti	1	ſ
	KKL _ 4,1692.6_1689	1		I	Π		1	¢	f	1	Ι				Γ			T	Ι	Ι				I	Π	ſ	T	Ι		Γ		T	T	Ι	Π	ſ	Γ	Π	Π	Π	Π	Т	J	T	Π	T	ſ
G. CERROA ZUZENSIS	KKL_4,1705.5_1702	] 🕺 🖏		Ш	Ш	Ĩ	1		L	1															Π	1				Γ					Π		L			$\Box$						<u>.</u>	ſ
	KKL _ 4, 1709 _ 1705.5			Ш	Π		1		T	t	T		П			$\square$			1	I	1			1	Π	Ш	1								Ц		L	ļ	Ш		Ц				Ш	4	Ļ
	KKL _ 4, 1712 5_ 1709	_ <u>~</u>		Ш		L I	4			U		L	Ц											L		Ш				1.					$\downarrow$			Ш	H		Ш			1	Ψ	ц.	L
	KKL _ 4, 1716 _1712.5		Ш	Ш			1		Ē	Ł			П	_											1													1L	Ш						Т	i L	
	KKL _ 4,1719_1716	, ä	61	1	III	ĽÌ	1	¢	t _	1		Ľ	Ц																Γ						П	1	L	Ш	Ш						J	Ľ	ſ
G, MEXICANA	NGT_ 1, 1906,5_1904.2	UPPER BATHYL	• •	Ш	cf		ILL	ŀ	• •	ź		Γ			Γ					Ι	Γ	Γ	¢f	ŀ	Π	¢1						T	Ι				Γ	ΤΓ	Π				П	IJ	Т	Τ	I
T.ROHRI	KKL _ 4,1764_1734	MICOLE TO					1		T		Т		Π					T	T	T	Π	Ŀ	•		Π	Ţ						T			$\square$			Π	P	$\Box$	IJ	J		Ц	0	n	ſ
O.BECKMANN	KKL _ 4,1770 _ 1764	OUTER SHELF		T	Т			F	¶_	ГТ	- [		IΤ	Т	Т	Т	ΓT	Т	T		П	Ŀ	•		Т	Т	T	F	1	I_		T	T	Г	П	1	Г	١F	17	17	10	1	i T	I	j þr	1	Г

The biostratigraphic unit Globigerinoides trilobus assemblage is proposed here, provisionally, to denote the interval in Cauvery basin from the top of G. kugleri/G. primordius zone to the level of the first appearance of Globigerinoides sicanus DE STEFANI (= G. bisphericus TODD). The planktonic foraminifera recorded from this interval include Globigerinoides trilobus trilobus (REUSS), G. altiaperturus BOLLI, G. subquadratus BRÖNNI-MANN, Globoquadrina altispira s. l., G. debiscens debiscens (CHAPMAN, PARR & COLLINS), Globorotalia obesa, G. siakensis and rare forms of Globorotalia peripheroronda BLOW & BANNER. Globigerina binaiensis KOCH and Globoquadrina praedebiscens are found in the lower part of this unit.

In KKL-4 Miogypsina (Miogypsinoides) dehaarti-M. (Miogypsina) tani, M. (Miogypsina) globulina and M. (Miogypsina) antellia are recorded from the sequence between 940 and 246 m. In Madanam well-1 Miogypsina (Miogypsinoides) bantamensis-M. (Miogypsina) gunteri, M. (Miogypsina) cf. mediterranea and M. (Miogypsina) antellia are recorded from the sequence between 1050 and 250 m. Further study on the planktonics and Miogypsinidae from the interval above the G. kugleri/G. primordius zone is under progress.

## Correlations

The distribution of the Upper Eocene to early Miocene planktonic foraminifera from the tropical regions, particularly as worked out by BOLLI (1957) from Trinidad, by BLOW & BANNER (1962) from Lindi area, East Africa and by BLOW (1969) from several tropical regions, and the data from the present investigation in Cauvery basin indicates that a detailed correlation of these sections is possible. Various successful attempts were made during the last few years to correlate the sections from different continents with the zonal scheme proposed by BOLLI (1957) and or by BLOW & BANNER (1962). The planktonic zones have found a wide spread acceptance and applicability in several regions. However, there is no agreement as to the usage of the planktonic zonal scheme. In general different zones were proposed for different parts of the world, obviously due to the reasons of geographical distribution of the planktonic foraminifera. BANNER & BLOW (1965) and BLOW (1969) have proposed a very detailed zonal scheme for the Upper Eocene to Recent tropical regions. On the other hand, a simplified planktonic foraminiferal zonal scheme applicable also to temperate Mediterranean Neogene was proposed as a result of the coordination of some 22 specialists, CATI et al. (1968).

The correlation of the zones from Cauvery basin with those proposed by BOLLI (1969) and BLOW (1969) is presented in figure 3. The correlations are based on the distribution of the planktonic foraminifera within the regions concerned with the assumption that the levels of the initial appearance and extinction of the species are synchroneus events in these regions. In general the successive levels of appearance and extinction of species are comparable.

However, a few minor discrepancies in the ranges of the presently recorded species could be noted from the published accounts. But it may be mentioned that the present observations are very close to those of BLOW (1969).

1. Cassigerinella chipolensis: BOLLI (1957) has shown that Cassigerinella chipolensis first appears at the base of G. ampliapertura zone in Trinidad. BLOW & BANNER (1962) traced its range from the base of G. oligocaenica zone in East Africa. BLOW (1969) gave the range of C. chipolensis from the Zone P 18 (= Globigerina tapuriensis zone) to Zone

Zones proposed here for Cauvery Basin	Blow	W. H. (1969)	Bolli H. M. (1966)
Globigerinoides trilobus	N. 6	Globigerinatella insueta / Globigerinita dissimilis	Catapsydrax stainforthi
assemblage	N. 5	Globoquadrina dehiscens praedehiscens-G. dehiscens dehiscens	Catapsydrax dissimilis
Globorotalia kugleri / Globigerinoides primordius	N. 4	Globigerinoides quadrilobatus primordius / Globorotalia kugleri	Globorotalia kugleri
Globigerina angulisuturalis	N. 3 (P. 22)	Globigerina angulisuturalis	Globigerina cipero- ensis ciperoensis
	N. 2 (P. 21)	Globigerina angulisuturalis / Globorotalia (T) opima opima	Globorotalia opima opima
Globigerina ampliapertura	N. 1 (P. 20)	Globigerina ampliapertura	Globigerina ampliapertura
Głobigerina sastrii	P. 19	Globigerina sellii / Pseudohastigerina barbadoensis	Cassigerinelia
	P. 18	Globigerina tapuriensis	Hastigerina micra
Globigerina gortanii	P. 17	Globigerina gortanii gortanii / Globorotalia (T) centralis	
Globorotalia cerroazulensisis	P. 16	Cribrohantkenina inflata	Globorotalia cerroazulensis
Globigerapsis mexicana	P. 15	Globigerapsis mexicana	Globigerapsis semiinvoluta
Truncorotaloides rohri	P. 14	Truncorotaloides rohri / Globigerinita howei	Truncorotaloides 10hri
Orbulinoides beckmanni	P. 13	Orbulinoides beckmanni	Porticulasphaera mexicana

Figure 3. Relationship of the zones proposed here for Cauvery basin to those given by BOLLI (1966) and BLOW (1969).

N 13 and gave the range of Cassigerinella eocaena CORDEY from Zone P 16 to Zone P 19. In Cauvery basin Cassigerinella chipolensis is found at the very basal part of the G. gortanii zone.

2. Globigerinoides primordius: This species appears in Cauvery basin at the base of G. kugleri/G. primordius zone, at a level before the first appearance of Globorotalia kugleri and the extinction of Globigerina angulisuturalis. BLOW (1969) has already noted the concurrence of the ranges of G. primordius and G. angulisuturalis in the lower part of Zone N 4. However, there are some minor discrepancies in the published accounts regarding the first appearance of Globigerinoides.

3. Globoquadrina appears in Cauvery basin in the G. angulisuturalis zone.

4. Many planktonic species viz. Globoquadrina altispira globosa, Globigerinita stainforthi appear in Cauvery basin in the uppermost part of the G. kugleri/G. primordius zone. In Trinidad these species are known to appear at the base of C. dissimilis zone (BOLLI, 1957). BLOW (1969) has shown that Globorotalia kugleri overlaps the ranges of G. stainforthi and G. altispira globosa.

Blow, 1969	Bolli, 1966	Figure 4.
N 5	C. dissimilis	 М Волл, 1957.
N 4	G. kugleri	- O M "Comité du Néosène"
N 3 (= P 22)	G. ciperoensis	M Connic du Neogene ,   O Blow, 1969; CATI et al., 1968.   M JENNING 1966
N 2 (= P 21)	G. opima opima	M BERGGREN, 1969.
N 1 (= P 20)	G. ampliapertura	О М Елмеs et al., 1962.
P 19	Cassigerinella	О Елмеs et al., 1962,
P 18	Hastigerina micra	E Eocene Colloquium, Paris, 1968.
P 17		$-\frac{O}{E}$ Bolli, 1957; Bandy, 1964.
P 16	G. cerroazulensis	$\frac{M}{O} = Oligocene / Miocene boundary$
P 15	G. semiinvoluta	$\frac{O}{E} = \frac{\text{Eocene / Oligocene}}{\text{boundary}}$

## Limits of Oligocene

The problem of marking the limits of Oligocene is topic of many debates and discussions during the last one and half decades. Recent attempts to relate the European Stages to the tropical planktonic zonal sequence have led to many controversial conclusions (EAMES et al, 1962; DROOGER, 1964; BLOW, 1969; BERGGREN, 1969, and others). The interpretations by various authors have led to controversies in marking the Eocene-Oligocene and Oligocene-Miocene boundaries. Some of the interpretations are summarized in figure 4.

Eocene-Oligocene boundary

The top of the Globorotalia cerroazulensis zone was taken to mark the Eocene-Oligocene boundary by many authors (BOLLI, 1957, 1966; BANDY, 1964, and others). The present author (RAJU, 1968) has earlier supported this view particularly taking account of the subsurface sections in Cauvery and Cambay basins, India.

The delegates to the Eocene Colloquium, Paris, 1968 (in Press, see BRABB, 1969) unanimously agreed for placing the Eocene-Oligocene limit between the Nummulites fabianii and N. intermedius zones for the Nummulites, between the Globigerina gortanii and Hastigerina micral Cassigerinella chipolensis zone (= Globigerina sellii zone) for the planktonic foraminifera and between the Isthmolithus recurvus and Ellipsolithus subdistichus zones for the nannoplanktons.

In Cauvery basin, in three of the wells, namely Madanam-1, TPD-1 and Tiruttaraipundi-1, where the larger foraminifera are dominant, the Eocene-Oligocene boundary is marked at the extinction level of *Pellatispira*. This level could be interpreted as marking the top of the Priabonian Stage. However, in the well sections around Karaikal and Nagapattinam the Upper Eocene larger foraminifera are not present and it is difficult to mark a horizon in the planktonic scale time equivalent to extinction level of *Pellatispira*. In these sections the extinction level of *G. cerroazulensis* and *Hantkenina* could be confidently marked on the basis of either core or cutting samples.

BLOW (1969) considered that the Eocene-Oligocene boundary will be approximately between the Zone P 17 (= G. gortanii/G. [T.] centralis zone) and Zone P 18 (= G. tapuriensis zone). BLOW (1969) and CLARKE & BLOW (1969) have shown that the characteristic Eocene foraminifera Pellatispira, Discocyclina, Hantkenina, and Globorotalia cerroazulensis all range only to the lower part of Zone P 17. It may be noted that the lower part of Zone P 17 of BLOW (1969) is equivalent to the uppermost part of G. cerroazulensis.

BAUMANN & ROTH (1969) observed that the Ericsonia subdisticha (= Ellipsolithus subdistichus) zone is restricted to the lower part of the Globigerina gortanii gortanii zone and further noted that the Latdorfian belongs to the E. subdisticha zone, and to the very base of the Cyclococco-

*lithus margaritae* zone. These evidences indicate that the Eocene-Oligocene boundary could be best placed at the top of the *G. cerroazulensis* zone.

Oligocene-Miocene boundary

This problem was discussed in detail by EAMES et al. (1962), DROOGER (1964), JENKINS (1966), RAJU (1968), BERGGREN (1969), and BLOW (1969). BERGGREN (1969) has drawn, provisionally, the Oligocene-Miocene boundary at the N 1/N 2 zonal boundary. BLOW (1969) accepted the recommendations of the "Comité du Néogène" in placing the Oligocene-Miocene boundary at the *Globigerinoides* datum, which coincides with the base of Zone N 4.

The Committee on Mediterranean Neogene stratigraphy has defined, in 1964 and 1967, the lower limit of the Miocene by the first appearance of Globigerinoides and Miogypsina gunteri (see PAPP et al., 1968). In Cauvery basin this level coincides with the lower boundary of G. kugleri/ G. primordius zone. In Madanam well M. (Miogypsina) gunteri-M. (Miogypsinoides) bantamensis assemblage is found. Thus in Cauvery basin the Oligocene-Miocene boundary could be marked on the basis of the criteria set forth by the Committee on Mediterranean Neogene Stratigraphy.

## Systematics

General remarks

Planktonic foraminifera are very rich in the Upper Eocene to early Miocene sediments in parts of Cauvery basin. The preservation of the specimens is good and the morphological details could be easily observed. Some specimens from the G. sastrii zone are, however, poorly preserved.

The specific determination was made first with the aid of the description and illustrations of the types. Most of the specimens were sent to Prof. H. M. BOLLI (Zürich) in 1968 for checking the identifications. Prof. BOLLI has kindly gone through the specimens and opined that the identications are also in accordance with his views. The present author has compared the specimens of Cauvery basin with the types deposited by BLOW & BANNER (1962) in the British Museum (Natural History), London. However, the determination of a few species viz. *Globigerina corpulenta* SUBBOTINA, *G. pseudocorpulenta* CHALILOV, is only on the basis of literature.

No attempts are made during the present work to trace the lineages as the available samples do not represent a complete subsurface sequence.

, The following five species were illustrated earlier by the present author (RAJU, 1968) and they are not repeated here. Globorotalia centralis CUSHMAN & BERMUDEZ, G. cerroazulensis (COLE), Chiloguembelina cubensis (PALMER), C. martini (PIJPERS) and Cassigerinella chipolensis (CUSHMAN & PONTON).

A few of the species viz. Globigerina officinalis SUBBOTINA, G. praebulloides group, G. ouachitaensis group, Gobigerinita spp. and Globorotalia gemma JENKINS, though determined during the present study, are not described here.

All the illustrated types are deposited in the Palaeontology Laboratory, O. N. G. Commission, Baroda, India.

Superfamily GLOBIGERINACEA CARPENTER, PARKER & JONES, 1862

## Family GLOBIGERINIDAE CARPENTER, 1862

Subfamily Globigerininae CARPENTER, PARKER & JONES, 1962

#### Genus Globigerina D'ORBIGNY, 1826

Globigerina ampliapertura BOLLI (Plate I, figs. 1 a—c)

Globigerina ampliapertura BOLLI, 1957, p. 108, pl. 22, figs. 5 a-7 b (not figs. 4 a-b).

Globigerina ampliapertura BOLLI, BLOW & BANNER, 1962, pp. 83-84, pls. XI A-D, XVII c, fig. 12 b.

Globigerina ampliapertura ampliapertura BOLLI, RAJU, 1968, pl. 4, fig. 7.

Remarks: The forms from Cauvery basin are well comparable with the type description and illustrations.

Stratigraphic range: BOLLI (1957) gave the range of this species from G. cocoaensis zone to G. opima opima zone in Trinidad. BLOW (1969) gave its range from the base of Zone P 17 to the basal part of Zone P 21(= N 2). In Cauvery basin this species is known from G. gortanii zone to the G. ampliapertura zone.

Globigerina angulisuturalis BOLLI (Plate I, figs. 2, 3 a, b)

Globigerina ciperoensis angulisuturalis Bolli, 1957, p. 109, pl. 22, figs. 11 a–c. Globigerina angulisuturalis Bolli, Blow & BANNER, 1962, p. 84, pl. IX, figs. A a–C c.

Remarks: The forms from Cauvery basin are well comparable with the type description and illustrations. These forms are compared with the hypotypes of BLOW & BANNER, 1962, from the Lower Ragusa Limestone, S. E. Sicily, deposited in British Museum (Natural History), London.

Stratigraphic range: BOLLI (1957) gave the range of this species from the base of the G. opima opima zone to the top of G. ciperoensis ciperoensis zone in Trinidad. It has also been reported from the Type Aquitanian of France (JENKINS, 1966), Israel (REISS & GVIRTZMANN, 1966), south-east Australia and Newzealand (JENKINS, 1960, 1965). BLOW (1969) gave its range from the base of Zone P 21 (= N 2) to lower part of Zone N 4.

#### Globigerina angustiumbilicata Bolli

(Plate I, fig. 6)

Globigerina ciperoensis angustiumbilicata BOLLI, 1957, p. 109, pl. 22, figs. 12 a-13 c, p. 164, pl. 36, figs. 6 a-b.

Globigerina angustiumbilicata BOLLI, BLOW & BANNER, 1962, p. 85, pl. IX x-z, figs. 9 (iv), 16 (vi, vii).

Remarks: In some of the forms from Cauvery basin the last chamber is smaller in size than the penultimate chamber, otherwise well comparable with the types.

Stratigraphic range: BOLLI (1957) gave its range from the G. ampliapertura zone to G. dissimilis zone in Trinidad. BLOW (1969) gave its range from the upper part of Zone P 16 to Zone N 22. In Cauvery basin this species is common and known from the G. cerroazulensis zone to a level above the G. kugleri/G. primordius zone.

Globigerina binaiensis KOCH (Text-figures 5, 6, 7, 8)

Globigerina binaiensis KOCH, 1935, nom. nov. for "G. aspera KOCH, 1926" (non EHREN-BERG), type figures in KOCH, 1926, Ecologae Geol. Helv., vol. 19 (1925-1926), no. 3, p. 745, figs. 22 a-c, 23 a-c.

Globigerina binaiensis Koch, Blow, 1969, p. 316, pl. 13, figs. 1, 2.

Remarks: This species is characterized in having three chambers in the last whorl and a large flattened apertural face of the last chamber. The specimens from Cauvery basin are very well comparable with the illustrations given by BLOW (1969).

Stratigraphic range: BLOW (1969) gave the range of G. binaiensis from the Zone P 22 (= N 3) to Zone N 5, and noted that it appears to be restricted to the Indo-Pacific Province. In Cauvery basin this species is abundant in the lower most part of the G. trilobus assemblage. Rare forms are also recorded from core samples belonging to the upper part of G. kugleri/G. primordius zone and also from the cutting samples belonging to the G. angulisuturalis zone.

> Globigerina ciperoensis BOLLI (Plate I, figs. 5 a, b)

Globigerina ciperoensis BOLLI, 1954, Contr. Cushman Found. Foram. Res., vol. 5, pt. 1, p. 1, text-figs. 3-6.

Globigerina ciperoensis ciperoensis BOLLI, BOLLI, 1957, p. 109, pl. 22, figs. 10 a-b.

Globigerina ouachitaensis ciperoensis Bolli, Blow & Banner, 1962, pp. 90–91, pl. IX E–G, fig. 9 (i–iii).

Stratigraphic range: BOLLI (1957) recorded this species from the G. opima opima zone to the top of G. ciperoensis zone in Trinidad. BLOW & BANNER (1962) recorded this species from the G. oligocaenina zone to G. ciperoensis zone and also mentioned its occurrence in the Aquitanian of South-west France. BLOW (1969) gave the range of G. ouachitaensis ciperoensis forma typica from Zone P 19 to Zone N 4 and forma atypica is shown to range up to Zone N 5. In Cauvery basin this species is known from G. angulisuturalis zone to G. kugleri/G. primordius zone.



Text-figures 5, 6, 7, 8. Globigerina binaiensis KOCH, from core sample at 874 m. in Madanam-1.

## Globigerina corpulenta SUBBOTINA (Plate V, figs. 1 a—c)

Globigerina corpulenta SUBBOTINA, 1953, p. 76, pl. 9, figs. 5-7, pl. 10, figs. 1-4.

Stratigraphic range: SUBBOTINA (1953) has originally described this species from the Upper Eocene, series F 3 (middle), zone of *Globigerinoides* conglobatus and large Globigerinas, from Northern Caucasus, U. S. S. R. In Cauvery basin this species is very common in the *G. mexicana* zone and a few specimens are also recorded from the *G. cerroazulensis* zone.

## Globigerina galavisi BERMUDEZ

(Plate V, figs. 2 a-c, 3)

Globigerina galavisi BERMUDEZ, 1961, Contribucion al Estudio de las Globigerinidea de la Region Caribe-Antillana, 3rd Congress Geol. Venezolano, Boletin de Geologia, Caracas, Venezuela, p. 1.183, pl. 4, fig. 3.

Globigerina galavisi BERMUDEZ, BLOW, 1969, p. 319, pl. 5, figs. 1-3, pl. 16, figs. 4, 5.

Remarks: The specimens from Cauvery basin are well comparable with the illustrations of holotype and metatype given by BLOW (1969).

Stratigraphic range: BLOW (1969) gave the range of this species from the Zone P 13 to P 21 (= N 2). This species is recorded so far from the G. mexicana zone to G. sastrii zone in Cauvery basin.

Globigerina gortanii gortanii (BORSETTI) (Plate II, figs. 1 a, b, 2)

Catapsydrax gortanii Borsetti, 1959, pp. 205–207, pl. 1, figs. 1 a–d. Globigerina turritilina turritilina BLOW & BANNER, 1962, pp. 98–99, pl. XIII, figs. D–G.

Remarks: The forms from Cauvery basin, in general, are well comparable with the holotype and paratypes of *Globigerina turritilina turritilina*. In several of the specimens from Cauvery basin, the umbilicus is either filled with matrix or covered by a compressed aberrant (bulla like) final chamber. The present specimens are also well comparable with the type description and illustrations of *Catapsydrax gortanii* BORSETTI.

Stratigraphic range: BORSETTI (1959) has originally described this species from the Lower Oligocene of Northern Italy. BLOW & BANNER (1962) recorded this species from *G. turritilina turrilitina* zone to *G. oligocaenica* zone in East Africa. BLOW (1969) gave its range from the base of Zone P 18 to the lower part of Zone P 22. In Cauvery basin this species is common in *G. gortanii* zone and *G. sastrii* zone.

> Globigerina gortanii praeturritilina BLOW & BANNER (Plate II, figs. 4 a, b, 5 a, b)

Globigerina turritilina praeturritilina BLOW & BANNER, 1962, p. 99, pl. XIII, figs. A-C.

Stratigraphic range: BLOW & BANNER (1962) gave the range of this species from G. semiinvoluta zone to the lower part of the G. turritilina turritilina zone. BLOW (1969) gave its range from the Zone P 14 to the middle part of Zone P 17. In Cauvery basin this species is recorded from the G. mexicana zone to G. gortanii zone.

Globigerina prasaepis BLOW (Text-figures 9, 10, 11)

Globigerina ampliapertura euapertura (JENKINS), BLOW & BANNER, 1962, p. 84, pl. XI, figs. E-G. (= holotype of G. prasaepis BLOW), not Globigerina euapertura JENKINS, 1960, p. 351, pl. 1, figs. 8 a-c.

Globigerina ampliapertura euapertura (JENKINS), RAJU, 1968, pl. 4, fig. 8.

Globigerina prasaepis BLOW, 1969, pp. 382-383, pl. 10, fig. 13, pl. 18, figs. 3-7.



Text-figures 9, 10, 11. Globigerina prasaepis BLOW, from G. gortanii zone, core sample at 1800 m. in NGT-1.

Remarks: The present author has compared the specimens from Cauvery basin with the hypotype of *Globigerina ampliapertura euapertura* of BLOW & BANNER, 1962 (= holotype of *G. prasaepis* BLOW, 1969) deposited in British Museum (Natural History), London. They are well comparable, except that the sutures in specimens from Cauvery basin are slightly more incised than in the type.

Stratigraphic range: BLOW (1969) gave the range of this species from upper part of the Zone P 17 to Zone P 21. In Cauvery basin this species is common in G. gortanii zone and G. sastrii zone.

## Globigerina pseudocorpulenta CHALILOV (Plate II, figs. 3 a, b)

Globigerina pseudocorpulenta CHALILOV, 1956 (on the pelagic foraminiferal fauna of the Paleogene of Azerbaidzhan; in Russian), Akad. Nauk. Azerb. S. S. R., Inst. Geol., Baku, Trudy vol. 17, p. 245, pl. 4, figs. 3 a-c (in ELLIS & MESSINA, 1940 et seq.).

Remarks: Specimens from Cauvery basin are well comparable with the illustrations and description of the type. However, the presently described forms seem to have less spinosity and the aperture often extends more towards extraumbilical.

Stratigraphic range: CHALILOV (1956) has originally described this species from lower part of the Upper Eocene, from the northeast foothills of the Maly Caucasus, Azerbaidzhan, S. S. R. In Cauvery basin this species is comon in *G. mexicana* zone. A few forms from *O. beckmanni* zone are comparable to this species.

Globigerina cf. rubriformis (SUBBOTINA) (Plate II, figs. 6 a, b)

Globigerinoides rubriformis SUBBOTINA, 1953, (part), p. 92, pl. 14, figs. 6-9 (not pl. 13, figs. 19 a-b).

Remarks: SUBBOTINA (1953) has placed probably the forms of both Globigerina and Globigerapsis under this species. Only one form (pl. 13, figs. 19 a—b) has been shown in her illustrations to contain definite supplementary apertures. However, the holotype (pl. 14, figs. 6 a—c) does not seem to have any supplementary apertures and she also commented that supplementary apertures are missing in some forms.

Stratigraphic range: SUBBOTINA (1953) has originally described this species from the *Globigerinoides conglobatus* zone, Upper Eocene and from the *Bolivina* zone, in Northern Caucasus, U. S. S. R. In Cauvery basin rare forms are recorded from *G. gortanii* zone.

Globigerina sastrii n. sp. (Plate IV, figs. 1 a-c, 2 a-c)

Description: Test low trochospiral, the equatorial profile is subcircular, equatorial periphery is broadly lobate; the axial periphery is broadly rounded with dorso-peripheral shoulder. The dorsal surface is flat to very slightly convex, the ventral surface is strongly vaulted. Wall calcareous, coarsely hispid, rugosities or short thick spines are found throughout, but more prominent around the umbilicus. Chambers, earlier ones spherical, later ones subconical, arranged in 2–3 whorls; the last 2 or 3 chambers increase very rapidly in size and strongly embrace the earlier ones; in umbilical view the last two chambers are prominently visible. Sutures on spiral side curved, less distinct in early stage, radial or oblique later, slightly depressed; umbilical side radial, depressed. Umbilicus small, narrow slit like, often covered by the last chamber, elongate, depressed; not sharply delineated. Aperture is often covered, narrow slit like.

Largest diameter of the holotype is 0.64 mm.

Depository: The types are deposited in the Palaeontology Laboratory, O. N. G. Commission, Baroda, India.

Type sample: A core sample, WLC-3, at a depth of 1580 m. in Karaikal well No. 4.

Age and distribution: Presence of Globigerina gortanii gortanii, G. ampliapertura, G. sellii, Globorotalia increbescens and other forms in the type sample indicates an Oligocene age. This species is recorded from TPD-1, NGT-1, Karaikal wells and Madanam well in Cauvery basin.

Remarks: This is a distinctive, short ranged species. This species is named after V. V. Sastri, O. N. G. Commission, India in recognition of his contributions to biostratigraphy in India.

Globigerina sellii (BORSETTI)

Plate III, figs. 5 a, b, 6)

Globoquadrina sellii Borsetti, 1959, p. 209, pl. 13, figs. 3 a–d. Globigerina oligocaenica BLOW & BANNER, 1962, p. 88, pl. X, figs. G, L–N. Remarks: The specimens from Cauvery basin are compared with the types of G. oligocaenica. In the presently described forms the overlap, on the ventral side, of the last chamber is of lesser degree than in the types. In general, the apertural face of the forms is broader and slightly more flattened than in the holotype of G. oligocaenica. The specimens from Cauvery basin are well comparable with the illustrations and type description of G. sellii.

Stratigraphic range: BORSETTI (1959) has originally described this species from the Oligocene of Northern Italy. BLOW & BANNER (1962) gave its range as restricted to *G. oligocaenica* Zone, Lindi area, East Africa. BLOW (1969) gave its range from Zone P 19 to Zone P 22 (= N 3). In Cauvery basin rare forms are recorded from *G. sastrii* zone.

## Globigerina tripartita tripartita Koch (Plate III, figs. 3 a, b)

Globigerina bulloides d'Orbigny var. tripartita Koch, 1926, Eclogae Geol. Helv., vol. 19 (1925–1926), no. 3, p. 746, text-figs. 21 a-b.

Globigerina tripartita tripartita Koch, Blow & BANNER, 1962, pp. 96–97, pl. X, figs. A-F.

Stratigraphic range: KOCH (1926) described this species from the lower beds of the "Globigerinenmergel" of East Borneo. BLOW (1969) gave the range of this species from the Zone P 14 to Zone P 22. In Cauvery basin typical forms are recorded from G. sastrii zone. Forms comparable to this species are also recorded from G. mexicana and G. cerroazulensis zones.

> Globigerina tripartita rohri Bolli (Plate III, figs. 2, 4 a, b)

Globigerina robri BOLLI, 1957, p. 109, pl. 23, figs. 1 a-4 b.

Remarks: Only forms well comparable to the types are assigned to this subspecies. Some forms from upper part of the G. kugleri/G. primordius zone and lower part of G. trilobus assemblage in Cauvery basin are comparable to this subspecies except in that they developed a definite umbilical teeth, on the basis of which they are referred here to Globoquadrina praedebiscens BLOW & BANNER.

Stratigraphic range: BOLLI (1957) gave its range from the G. ampliapertura zone to C. dissimilis zone in Trinidad. In Cauvery basin a few forms are recorded from the G. sastrii zone.

Globigerina venezuelana Hebberg

(Plate V, fig. 4, plate VI, figs. 1 a-c)

Globigerina venezuelana HEDBERG, 1937, Jour. Paleont., vol. 11, no. 8, p. 681, pl. 92, figs. 7 a-b.

Globigerina venezuelana HEDBERG, BOLLI, 1957, p. 110, pl. 23, figs. 6 a-8 b.

Globigerina venezuelana Hedberg, BLOW & BANNER, 1962, text fig. 11.

Stratigraphic range: BOLLI (1957) recorded this species from the G. ampliapertura Zone to G. menardii Zone in Trinidad. BLOW (1969) gave its range from Zone P 22 to Zone N 9. In Cauvery basin typical forms are recorded so far from the G. kugleri/G. primordius zone and lower part of G. trilobus assemblage.

Globigerina woodi woodi Jenkins

(Plate V, figs. 5 a-c)

Globigerina woodi JENKINS, 1960, p. 352, pl. 2, figs. 2 a–c. Globigerina woodi JENKINS, REISS & GVITZMANN, 1966, pl. 91, figs. 1 a–c. Globigerina woodi woodi JENKINS, JENKINS, 1966, p. 6, pl. 1, figs. 18 a–c.

Stratigraphic range: JENKINS (1960) gave its range from base of the G. woodi zone to G. menardii miotumida zone in Australia. This species has also been recorded from Newzealand (JENKINS, 1965), Israel (REISS & GVIRTZMANN, 1966) and from the type Aquitanian of France (JENKINS, 1966). In Cauvery basin this species is recorded from G. angulisuturalis zone and G. kugleri/G. primordius zone.

Genus Globigerinoides Cushman, 1927

Globigerinoides primordius BLOW & BANNER (Plate VII, figs. 1 a, b, 2 a, b)

Globigerinoides quadrilobatus primordius BLOW & BANNER, 1962, p. 115, pl. IX, figs. D d—F f, fig. 14 (iii—viii).

Remarks: The specimens from Cauvery basin are well comparable to the holotype. The primary aperture of the Cauvery basin forms is more wider than in the holotype. However, these and other minor differences fall within the variation limits of paratypes designated by BLOW & BANNER (1962).

Stratigraphic range: BLOW & BANNER (1962) gave the range of this species as confined to the *G. kugleri* zone in Trinidad. JENKINS (1966) recorded forms comparable to this species from the type Aquitanian of France. BLOW (1969) gave its range from base of the Zone N 4 to lower part of the Zone N 5. In Cauvery basin it is common throughout the *G. kugleri/G. primordius* zone.

Globigerinoides trilobus trilobus (REUSS) (Plate VII, figs. 4 a, b)

Globigerina triloba REUSS, 1850 Denkschr. k. k. Akad. Wiss., Wien, mathem.-naturwiss. Cl., vol. 1, p. 374, pl. 47, figs. 11 a-d (in ELLIS & MESSINA, 1940 et seq.).

Globigerinoides triloba triloba (REUSS), BOLLI, 1957, p. 112, pl. 25, figs. 2 a-c.

Stratigraphic range: BOLLI (1957) gave the range of this species from the C. dissimilis zone to G. menardii zone in Trinidad. In Cauvery basin it appears in uppermost part of the G. kugleri/G. primordius zone and is common throughout the Lower-Middle Miocene.

## Globigerinoides trilobus altiaperturus BOLLI (Plate VII, figs. 4 a, b)

Globigerinoides triloba altiapertura BOLLI, 1957, p. 113, pl. 25, figs. 7 a-8, text fig. 21, no. 3.

Stratigraphic range: BOLLI (1957) gave its range from base of the C. dissimils zone to C. stanforthi zone in Trinidad. This species has also been recorded from the type Burdigalian of France (JENKINS, 1966), Newzealand (JENKINS, 1965), Australia (WADE, 1964), Israel (REISS & GVITTZMANN, 1966). In Cauvery basin it is recorded from uppermost part of the G. kugleri/G. primordius zone and G. trilobus assemblage.

Genus Globigerinita BRÖNNIMANN, 1951 emended BLOW & BANNER, 1962

Globigerinita dissimilis (Cushman & Bermudez) (Plate VI, figs. 2 a, b, 3 a, b)

Globigerina dissimilis Cushman & Bermudez, 1937, p. 25, pl. 3, figs. 4---6. Catapsydrax dissimilis (Cushman & Bermudez), Bolli, 1957, p. 36, pl. 7, figs. 6 a-8 c. Globigerinita dissimilis (Cushman & Bermudez), BLOW & BANNER, 1962, p. 106, pl. 14, figs. A-D.

Remarks: The illustrated specimen from G. kugleri/G. primordius zone in Cauvery basin has a slightly more convex spiral side than in the holotype.

Stratigraphic range: BOLLI (1957) gave its range from T. rohri zone to C. stainforthi zone in Trinidad. In Cauvery basin rare forms of this species are recorded from G. mexicana zone to lower part of G. trilobus assemblage. Very rare forms comparable to G. dissimilis are also recorded from sequence as low as Hantkenina aragonensis zone in Cauvery basin.

Globigerinita stainforthi (BOLLI, LOEBLICH & TAPPAN) (Plate VI, figs. 4 a-c)

Catapsydrax stainforthi Bolli, LOEBLICH & TAPPAN, 1957, p. 36, pl. 7, figs. 11 a-c.

Remarks: The specimens from Cauvery basin are well comparable with the type description and illustrations.

Stratigraphic range: BOLLI (1957) gave the range of this species from the C. dissimilis zone to C. stainforthi zone in Trinidad. BLOW (1969) gave its range from the latest part of Zone N 4 to Zone N 7. In Cauvery basin this species is known from the uppermost part of G. kugleri/G. primordius zone and in the sequence immediately overlying this zone.

Genus Globigerapsis Bolli LOEBLICH & TAPPAN, 1957

Globigerapsis mexicana (CUSHMAN) (Plate XI, figs. 1, 2 a, b, 3 a, b, 4, 5, 6)

Globigerina mexicana Cushman, 1925, Cushman Lab. Foram. Res., Contr., vol. 1, p. 6, pl. 1, fig. 8 a, b.

Globigerinoides semiinvoluta KEIJZER, 1945, Univ. Utrecht, Geogr. Geol. Med. Phys-Geol. Reeks; ser. 2, no. 6, p. 206, pl. 4, figs. 58 a-e.

Globigerapsis semiinvoluta (KEIJZER), BOLLI, 1957, p. 165, pl. 36, figs. 19-20.

Globigerapsis semiinvoluta (KEIJZER), BLOW & BANNER, 1962, p. 125, pl. XV, figs. J–L. Globigerapsis semiinvoluta (KEIJZER), RAJU, 1968, pl. 2, figs. 4, 5.

Globigerapsis mexicana (CUSHMAN), BLOW & SAITO, 1968, Micropaleontology, vol. 14, no. 3, pp. 357–360, text-figs. 1–4.

Remarks: The specimens from Cauvery basin are in general smaller in size than the type. The sutures are not distinct and often the initial chambers are not visible. The sutures in the hypotype of *G. semiinvoluta* of BLOW & BANNER, 1962, are more prominently visible than in these forms.

Stratigraphic range: BOLLI (1957) and BLOW & BANNER (1962) have given the range of this species as restricted to the limits of its zone within the lower part of the Upper Eocene.

## Globigerapsis cf. tropicalis BLOW & BANNER (Plate XI, figs. 7 a, b)

Globigerapsis tropicalis BLOW & BANNER, 1962, pp. 124-125, pl. XV, figs. D-F.

Remarks: The presently illustrated forms from Cauvery basin differ slightly from the holotype of G. tropicalis in being larger in size, in having more depressed sutures, more elongate and lower supplementary apertures. The wall structure is comparable.

Stratigraphic range: BLOW (1969) gave the range of this species from Zone P 10 (= H. aragonensis Zone) to Zone P 16.

Genus Globoquadrina FINLAY, 1947

Globoquadrina altispira globosa BOLLI (Plate VII, figs. 6 a, b)

Globoquadrina altispira globosa BOLLI, 1957, pp. 111-112, pl. 24, figs. 9 a-10 c.

Remarks: The specimens from Cauvery basin are often badly preserved and the umbilicus is filled with pyritized matrix. The last one or two chambers in the last whorl increase rapidly in size than in the forms described by BOLLI (1957).

Stratigraphic range: BOLLI (1957) gave the range of this species from base of the C. dissimilis zone to G. insueta zone and again from G. fohsi robusta zone to G. menardii zone in Trinidad. The presently illustrated forms are from uppermost part of the G. kugleri/G. primordius zone. It is common in the G. trilobus assemblage. BLOW (1969) gave its range from Zone P 22 (= N 3) to Zone N 20.

> Globoquadrina debiscens (CHAPMAN, PARR & COLLINS) (Plate IX, figs. 3, 4 a, b)

Globorotalia debiscens CHAPMAN, PARR & COLLINS, 1934, Linn. Soc. London, Jour. Zool., vol. 38, no. 262, p. 569, pl. 11, figs. 36 a-c (in ELLIS & MESSINA, 1940 et seq.).

Globoquadrina debiscens (CHAPMAN, PARR & COLLINS), BOLLI, 1957, p. 111, pl. 24, figs. 3 a-4 c.

Remarks: The subspecies of G. dehiscens are not recognized in the present study.

Stratigraphic range: BOLLI (1957) gave its range from G. stainforthi zone to G. menardii zone in Trinidad. In Cauvery basin this species is common in the G. trilobus assemblage.

> Globoquadrina cf. larmeui Akers (Plate IX, figs. 5 a, b)

Globoquadrina larmeui Akers, 1955, p. 661, pl. 65, figs. 4 a-c. Globoquadrina larmeui Akers, Jenkins, 1960, p. 355, pl. 3, figs. 1 a-2 c.

Remarks: The specimens from Cauvery basin differ slightly from the type illustrations and description in being more convex dorsally and larger in size.

Stratigraphic range: JENKINS (1960) recorded this species from the Pre-Globoquadrina dehiscens dehiscens zone to the O. universa zone in Australia. In Cauvery basin this species is recorded so far from the G. angulisuturalis zone and G. kugleri/G. primordius zone.

Globoquadrina praedehiscens BLOW & BANNER (Plate VIII, figs. 4 a-c; Plate IX, fig. 12)

Globoquadrina dehiscens praedehiscens BLOW & BANNER, 1962, pp. 116-117, pl. XV, figs. Q-S.

Globoquadrina praedebiscens BLOW & BANNER, SAITO, 1963, p. 193, pl. 55, figs. 7 a-b. Globoquadrina debiscens praedebiscens BLOW & BANNER, REISS & GVIRTZMANN, 1966, pl. 96,

figs. 3—7.

Remarks: The specimens from Cauvery basin are compared with the holotype. In most of the present specimens the last chamber is relatively smaller, the umbilical teeth is slightly the and wider than in the holotype.

Stratigraphic range: BLOW & BANNER (1962) gave its range from middle part of the G. ciperoensis zone to the lower part of G. stainforthi zone in Trinidad. They have also reported its occurrence from Venezuela, East Africa, and Newzealand. In Cauvery basin typical forms are recorded from G. kugleri/G. primordius zone and lower part of the G. trilobus assemblage.

## Genus Globorotalia CUSHMAN, 1927

Globorotalia centralis Cushman & Bermudez

Globorotalia centralis Cushman & Bermudez, 1937, p. 26, pl. 2, figs. 62–65.

Globorotalia centralis Cushman & Bermudez, Bolli, 1957, p. 162, pl. 39, figs. 1-4.

Globorotalia (Turborotalia) centralis Cushman & Bermudez, Blow & Banner, 1962, p. 117, pl. XII, figs. K-M, Fig. 12 c, d.

Turborotalia centralis Cushman & Bermudez, Raju, 1968, pl. 4, figs. 3 a-c.

Remarks: This species was earlier illustrated by the present author (RAJU, 1968) and given as *Turborotalia centralis*.

Stratigraphic range: BOLLI (1957) gave its range from the Globigerapsis kugleri zone to G. cocoaensis zone in Trinidad. BLOW (1969) noted that it ranges from Zone P 11 to Zone P 17. In Cauvery basin it has been recorded from the H. aragonensis zone to top of the G. cerroazulensis zone.

Globorotalia cerroazulensis (COLE)

Globigerina cerroazulensis COLE, 1928, Bull. Amer. Pal., vol. 14, no. 53, p. 217, pl. 32, figs. 11-13.

Globorotalia cocoaensis Cushman, 1928, Contr. Cushman Lab. Foram. Res., vol. 4, pt. 3, p. 75, pl. 10, figs. 3 a-c.

Globorotalia cocoaensis Cushman, Bolli, 1957, p. 169, pl. 39, figs. 5 a-7 b.

Globorotalia (Turborotalia) cerroazulensis (COLE), BLOW & BANNER, 1962, p. 118, pl. XII, figs. D-F, Fig. 12 d, e.

Turborotalia cerroazulensis (COLE), RAJU, 1968, pl. 3, figs. 7 a-c.

Remarks: This species was earlier illustrated by the present author (RAJU, 1968) and given as *Turborotalia cerroazulensis*.

Stratigraphic range: BOLLI (1957) gave its range from the base of the G. semiinvoluta zone to top of the G. cocoaensis zone in Trinidad. BLOW (1969) gave its range from upper part of Zone P 14 (T. rohri/G. howeii zone) to the lower part of Zone P 17. In Cauvery basin it is known from the G. mexicana zone to top of the G. cerroazulensis zone. Some rare forms from O. beckmanni zone appear to be comparable to this species.

> Globorotalia increbescens (BANDY) (Plate IX, figs. 7 a-c)

Globigerina increbescens BANDY,1949, p. 120, pl. 23, figs. 3 a—c. Globorotalia (Turborotalia) increbescens (BANDY), BLOW & BANNER, 1962, pp. 118—119, pl. XIII, figs. T—V, pl. XVII, figs. D, K, Fig. 9 (13—15).

Remarks: The specimens from Cauvery basin are compared with a metatype, from Upper Jackson, Little Stave Creek, Clarks county, Alabama, deposited in British Museum (Natural History), London. The illustrated specimen from Cauvery basin is slightly bigger in size and has slightly higher aperture, otherwise well comparable with the metatype.

Stratigraphic range: This species has been originally described by BANDY (1949) from Jacksonian Stage, Alabama, U. S. A. BLOW (1969) gave the range of this species from Zone P 15 to Zone P 19. In Cauvery basin typical forms are recorded from G. gortanii zone and G. sastrii zone.

Globorotalia kugleri Bolli

(Plate X, figs. 1 a, b, 2 a, b)

Globorotalia kugleri BOLLI, 1957, p. 118, pl. 28, figs. 5 a-6. Globorotalia (Turborotalia) kugleri BOLLI, BLOW, 1969, p. 350, pl. 38, figs. 1-4. Remarks: The specimens from Cauvery basin are well comparable with the type description and illustrations given by BOLLI (1957) and the illustrations of topotypes given by BLOW (1969).

Stratigraphic range: BOLLI (1957) gave the range of this species as restricted to the G. kugleri zone in Trinidad. BLOW (1969) gave its range from Zone P 22 (= N 3) to the top of Zone N 4. In Cauvery basin it is recorded so far only from upper part of the G. kugleri/G. primordius zone.

Globorotalia obesa BOLLI (Plate X, figs. 6 a—c)

Globorotalia obesa BOLLI, 1957, p. 119, pl. 29, figs. 2 a-c. Globigerina obesa BOLLI, VERVLOET, 1966, p. 53, pl. II, figs. 9 a-c, pl. V, figs. 2 a-c.

Stratigraphic range: BOLLI (1957) gave its range from C. dissimilis zone to G. menardii zone in Trinidad. In Cauvery basin this species is common in the upper part of G. kugleri/G. primordius zone and G. trilobus assemblage.

Globorotalia opima nana Bolli (Plate X, figs. 3 a, b)

Globorotalia opima nana BOLLI, 1957, p. 118, pl. 28, figs. 3 a—c. Globorotalia (Turborotalia) opima nana BOLLI, BLOW & BANNER, 1962, pp. 119—120, pl. XIII, figs. Q—S.

Remarks: The specimen illustrated here is slightly larger in size than the holotype. Most of the unillustrated forms are better comparable with the type.

Stratigraphic range: BLOW (1969) gave the range of this species from Zone P 15 to Zone P 22 (= N 3). In Cauvery basin it is recorded from G. mexicana zone to the lower part of G. kugleri/G. primordius zone.

Globorotalia peripheroronda BLOW & BANNER (Plate IX, figs. 6 a—c)

Globorotalia fobsi barisanensis LEROY, BOLLI, 1957, p. 119, pl. 28, figs. 8 a-c.

Globorotalia (Turborotalia) peripheroronda BLOW & BANNER, 1966, Micropaleontology, vol. 12, no. 3, p. 294, pl. 1, figs. 1-3.

Stratigraphic range: BOLLI (1957) gave its range from the C. dissimilis Zone to G. fohsi fohsi Zone in Trinidad. This species has been also reported from several parts of the world including Japan (SAITO, 1963), Newzealand (JENKINS, 1965), Australia (JENKINS, 1960) and Israel (REISS & GVIRTZMANN, 1966). In Cauvery basin a few forms are recorded from G. trilobus assemblage.

Globorotalia siakensis (LEROY)

(Plate X, figs. 4, 5 a-c)

Globigerina siakensis LEROY, 1939, Natuurk. Tijdschr. Nederl.-Indie, vol. 99, no. 6, pp. 39–40, pl. 3, figs. 30–31.

Jahrb. Geol. B. A., 1971, Sonderband 17

Globorotalia mayeri Cushman & Ellisor, Bolli, 1957, p. 118, pl. 28, figs. 4 a-c (not Globorotalia mayeri Cushman & Ellisor, 1939).

Globorotalia siakensis (LEROY), JENKINS, 1960, p. 366, pl. 5, figs. 7 a-c.

Globorotalia (Turborotalia) siakensis LEROY, BLOW (1969), p. 356, pl. 10, figs. 7-9, holotype refigured, pl. 34, figs. 4, 5.

Remarks: This species is very common throughout its range in Cauvery basin.

Stratigraphic range: BOLLI (1957) gave its range from the G. opima opima zone to G. mayeri zone in Trinidad. BLOW (1969) gave its range from Zone P 22 to N 14. In Cauvery basin this species is known from G. angulisuturalis zone to Middle-Upper Miocene.

Genus Globorotaloides BOLLI, 1957

Globorotaloides suteri BOLLI

(Plate VI, figs. 5 a, b)

Globorotaloides suteri Bolli, 1957, p. 116, 166, pl. 27, figs. 9–13 b, pl. 37, figs. 10 a–12. Globorotaloides suteri Bolli, Blow & BANNER, 1962, pp. 122–123, pl. XII, figs. N–P, Fig. 11 (V, IX).

Remarks: The specimens from Cauvery basin are well comparable with the type illustrations and description. These specimens differ slightly from the hypotypes of BLOW & BANNER (1962) in having a low lying bulla and the rate of increase in chamber size is slower.

Stratigraphic range: BLOW & BANNER (1962) recorded this species from the upper Lutetian to about the middle part of the Aquitanian in East Africa and Trinidad. BLOW (1969) gave its range from Zone P 13 to within Zone N 8. In Cauvery basin this species is recorded from *G. mexicana* zone to the *G. kugleri/G. primordius* zone.

Genus Globigerinopsis Bolli, 1962

Globigerinopsis guhai n. sp. (Plate VIII, figs. 1 a-c, 2 a-c, 3 a-c)

Description: Test low trochospiral, equatorial profile of the test subquadrate, equatorial periphery strongly lobate, axial periphery rounded. Wall calcareous, porous, finely pitted, well preserved specimens with fine small spines. Chambers inflated, spherical except the last one or two; last two in spiral view ovate, slightly elongate; the last chamber varies in size, in general equal or larger than the penultimate. Some 16 chambers arranged in 3—4 whorls; the 4—4½, rarely 5 chambers in last whorl increase rapidly in size. Sutures on spiral side radial to slightly oblique, strongly depressed; on umbilical side radial, strongly depressed. Umbilicus wide, deep, filled often with pyritized matrix. Primary aperture a high arch, interiomarginal umbilical to slightly spiroumbilical, without lip or rim. One secondary aperture, a medium arch with small rim. Largest diameter of the holotype is 0.624 mm. Depository: The holotype and paratypes are deposited in the Palaeontology Laboratory, O. N. G. Commission, Baroda, India.

Type sample: Holotype and illustrated paratypes are from a core sample (CC I, 1126.3—1128.8 m., 15—26 cm.) from Karaikal well no. 2, Cauvery basin. Rare specimens are also recorded from cutting sample at a depth of 1000 m. in KKL-1.

Age of the type sample: Uppermost part of Globorotalia kugleri/ Globigerinoides primordius zone, Lower Miocene (Aquitanian age).

Remarks: This species has been placed under the genus Globigerinopsis BOLLI, 1962. However, the present species has a supplementary aperture, whereas the type of Globigerinopsis do not have and also the primary aperture is more spiroumbilical in the types. SCHMID (1967) described a n. sp., Globigerinopsis grilli SCHMID, which has a supplementary aperture, but its wall surface is coarselly pitted and cancellate.

This species is named after Mr. D. K. GUHA, O. N. G. Commission, India, in recognition of his contributions to microfauna and biostratigraphy in India.

Genus Pseudobastigerina BANNER & BBLOW, 1959

Pseudohastigerina barbadoensis BLOW, 1969 (Text-figures 12, 13, 14)

Pseudohastigerina barbadoensis BLOW, 1969, pp. 409-410, pl. 53, figs. 7-9, pl. 54, figs. 1-3.

Remarks: The specimens from Cauvery basin are very well comparable with the illustrations and description of the types.

Stratigraphic range: BLOW (1969) gave the range of this species from Zone P 16 to the top of Zone P 19. In Cauvery basin this species is very common in G. gortanii zone, and very rare in the G. cerroazulensis zone.



Text-figures 12, 13, 14. Pseudohastigerina barbadoensis BLOW, from G. gortanii zone, core sample at 1800 m. in NGT-1.

Nonion micrus COLE, 1927, Bull. Amer. Paleont., vol. 14, no. 51, p. 22, pl. 5, fig. 12. Hastigerina micra (COLE), BOLLI, 1957, p. 161, pl. 35, figs. 1 a-2 b.

Pseudohastigerina micra (COLE), BANNER & BLOW, 1959, pp. 19-20, pl. 3, figs. 6 a-b, text-figs. 4 g-i.

Pseudohastigerina micra (COLE), BLOW, 1969, pl. 53, figs. 1, 4, 5, 6.

Stratigraphic range: BOLLI (1957) gave the range of this species from the *H. aragonensis* zone to *G. cocoaensis* zone in Trinidad. BLOW (1969) gave its range from Zone P 10 to within Zone P 19. In Cauvery basin basin typical forms are recorded from *H. aragonensis* to top of the *G. cerroazulensis* zone. Only rare forms from *G. gortanii* zone are comparable to this species.

Genus Hantkenina Cushman, 1925

Hantkenina alabamensis Cushman

(Plate XI, figs. 8, 9 a, b; Plate XII, fig. 2)

Hantkenina alabamensis CUSHMAN, 1925, Proc. U. S. Nat. Mus., vol. 66, no. 2567, art. 30, pp. 3-4, pl. 1, figs. 1-6, pl. 2, fig. 5, text-fig. 1.

Hantkenina alabamensis CUSHMAN, BANDY, 1949, p. 76, pl. 11, figs. 9 a-b.

Hantkenina (Hantkenina) alabamensis Cushman, Brönnimann, 1950, p. 414, pl. 56, figs. 10, 14–16.

Hantkenina alabamensis Cushman, Bolli, LOEBLICH & TAPPAN, 1957, pp. 26-28, pl. 2, fig. 8.

Hantkenina alabamensis CUSHMAN, NAGAPPA, 1959, pl. 11, figs. 14 a-b.

Hantkenina alabamensis Cushman, Blow & Banner, 1962, pp. 126–127, pl. XVI, figs. C-D, J, K.

Hantkenina alabamensis Cushman, Raju, 1968, pl. 1, fig. 11.

Remarks: The specimens from Cauvery basin show considerable variation in shape of the chambers and sutures. A few forms have hispid wall surface, particularly in the early chambers of the last whorl. A solitary form with twin spines (see Plate XI, figs. 9 a, b) is also found.

Stratigraphic range: BLOW (1969) gave the range of this species from within Zone P 13 (= O. *beckmanni* zone) to the top of Zone P 16. In Cauvery basin typical forms of this species are recorded so far from the G. mexicana zone and G. cerroazulensis zone.

Hantkenina suprasuturalis BRÖNNIMANN (Plate XII, figs. 9 a, b, 10; Plate XIII, fig. 1)

Hantkenina (Hantkenina) suprasuturalis BRÖNNIMANN, 1950, p. 416, pl. 56, figs. 12, 13, text-fig. 2.

Remarks: The specimens from Cauvery basin are well comparable with the illustrations and description of the type. Several small sized specimens appear to grade morphologically to *Hantkenina alabamensis*.

Stratigraphic range: BRÖNNIMANN (1950) has originally reported this

species from the Upper Eocene of Trinidad and Barbados. In Cauvery basin typical forms of this species are common in the *G. mexicana* zone and rare in *G. cerroazulensis* zone.

Hantkenina cf. thalmanni Brönnimann (Plate XII, figs. 1 a-b)

Hantkenina thalmanni BRÖNNIMANN, 1950, pp. 415-416, pl. 55, figs. 19-21, pl. 56, figs. 3, 11.

Remarks: Only three specimens which are comparable with the illustrations and description of the types are found from the *G. mexicana* zone in Cauvery basin. It is not clear from the available material whether they are juvenile forms of *H. suprasuturalis*.

Hantkenina trinitatensis BRÖNNIMANN (Plate XII, figs. 3, 4, 5)

Hantkenina (Applinella) trinitatensis BRÖNNIMANN, 1950, pp. 410-411, pl. 56, fig. 17.

Remarks: Only a few specimens which are well comparable with the illustrations and description of the types are found from the *G. mexicana* zone in Cauvery basin.

Genus Cribrohantkenina THALMANN, 1942

Cribrohantkenina inflata (Howe) (Platte XIII, figs. 2--8)

Hantkenina inflata Howe, 1928, Jour. Paleont., vol. 2, no. 1, p. 14, fig. 2.

Hantkenina mccordi Howe & WALLACE, 1932, Louisiana Dept. Conserv., Geol. Bull., no. 2, p. 55-56, tab. 10, figs. 1 a, b.

Hantkenina danvillensis Howe & WALLACE, 1934, Jour. Paleont., vol. 8, pp. 35-37, pl. 5, figs. 14-17.

Hantkenina (Cribrobantkenina) bermudezi THALMANN, 1942, Amer. Jour. Sci., vol. 240, no. 11, pp. 812, 815, pl. 1, figs. 5-6.

Cribrohantkenina mccordi Howe & WALLACE, BANDY, 1949, p. 76, pl. 11, figs. 10 a-b.

Hantkenina (Cribrohantkenina) bermudezi THALMANN, BRÖNNIMANN, 1950, p. 417, pl. 56, figs. 6–9.

Cribrohantkenina bermudezi THALMANN, BOLLI, LOEBLICH & TAPPAN, 1957, pp. 28-29, pl. 2, figs. 9 a-11 b.

Hantkenina (Cribrohantkenina) bermudezi THALMANN, NAGAPPA, 1959, pl. 11, figs. 15 a-b.

Cribrobantkenina danvillensis Howe & WALLACE, BLOW & BANNER, 1962, p. 128, pl. XVI, figs. G-H, Fig. 19 (i-vii).

Cribrohantkenina inflata (Howe), SPRAUL, 1963, Jour. Pleont., vol. 37, no. 2, p. 367, pl. 14, figs. 1 a-4 b.

Cribrohantkenina inflata (Howe), RAJU, 1968, pl. 1, fig. 7.

Remarks: BLOW & BANNER (1962) discussed in detail the apertural characters of *Cribohantkenina*. SPRAUL (1963, cited above) has reillustrated the type of *Hantkenina inflata* HOWE, 1928 and shown that it is a true *Cribrohantkenina*, thus clarified the taxonomic status of several of the species.

In Cauvery basin both juvenile and adult specimens of this species are recorded from G. cerroazulensis zone. A study of these forms indicate that C. inflata has maintained the club shaped final chambers almost throughout its ontogeny. Similar data was earlier presented by DIENE & PROTO DECIMA (1964).

Stratigraphic range: This species is known to be an important worldwide index fossil of Upper Eocene. BLOW & BANNER (1962) recorded this species from about middle part of *G. semiinvoluta* zone to the top of *C. danvillensis* zone in Lindi area, East Africa. This species is common throughout the *G. cerroazulensis* zone in Cauvery basin. Very rare and poorly preserved specimens with doubtful cribrate aperture are found in *G. mexicana* zone.

#### Genus CASSIGERINELLA POKORNY, 1955

Cassigerinella chipolensis (CUSHMAN & PONTON)

- Cassidulina chipolensis CUSHMAN & PONTON, 1932, Florida Geol. Survey, Bull., no. 9, p. 98, pl. 15, figs. 2 a-c.
- Cassigerinella boudecensis POKORNY, 1955, Vestnik, Ustred. Geol., 30, pp. 136-140, textfigs. 1-3.

Cassigerinella globolocula IVANOVA, 1958, in BYKOVA, Mikrofauna S. S. S. R., sb. 9, vip. 115; Trudi. Vses. Neft. Nauchno-Issled. Geol. Inst. (VNIGRI), p. 57, pl. 11, figs. 1—3.

Cassigerinella chipolensis (Cushman & Ponton), Blow & Banner, 1962, pp. 81–82, pl. XV, figs. M, N.

Cassigerinella chipolensis (CUSHMAN & PONTON), RAJU, 1968, pl. 1, figs. 9, 10.

Remarks: This species from Cauvery basin was illustrated earlier by the present author (RAJU, 1968). In general the size of the forms increase, slightly, from G. gortanii zone to G. sastrii zone.

Stratigraphic range: BOLLI (1957) gave the range of this species from the G. ampliapertura zone to G. fohsi robusta zone in Trinidad. BLOW (1969) gave its range from Zone P 18 to Zone N 13. In India this species is known from Kutch, Cambay basin, Assam, Cauvery basin, Quilon area and Andaman Islands. In Cauvery basin this species first appears in the basal part of the G. gortanii zone and is very common in the sequence from the G. sastrii zone to Middle Miocene. A few forms are found in the section above the extinction level of Miogypsina antellia.

Genus Chiloguembelina LOEBLICH & TAPPAN, 1956

Chiloguembelina cubensis (PALMER)

Guembelina cubensis PALMER, 1934, Mern. Soc. Cubana Hist. Nat., vol. 8, p. 74, text-figs. 1-6.

Chiloguembelina cubensis (PALMER), BECKMANN, 1957, p. 89, pl. 21, fig. 21, text-fig. 14, nos. 5-8.

Chiloguembelina cubensis (PALMER), RAJU, 1968, pl. 1, figs. 4 a-b.

Remarks: This species from Cauvery basin was illustrated earlier by the present author (RAJU, 1968). This small and distinct species is abundant in the fine fractions of the samples, within its range, in Cauvery basin. This species is also common in shallow marine facies of Oligocene age in Cambay basin, western India and found together with Nummulites fichteliintermedius.

Stratigraphic range: BECKMANN (1957) gave the range of this species from *Porticulasphaera mexicana* zone to *G. opima opima* zone in Trinidad. The species has also been reported from various parts of the world in sections of Upper Eocene to Oligocene. BLOW (1969) gave the range of *Chiloguembelina* ex group *cubensis* from Zone P 13 to very close to the Zone N 4/Zone N 3 (= P 22) boundary. In Cauvery basin this species is common from the *G. mexicana* zone to *G. ampliapertura* zone.

#### Chiloguembelina martini (PIJPERS)

Textularia martini PIJPERS, 1933, Univ. Utrecht Geogr. Geol. Med. Phys.-Geol. Reeks, no. 8, p. 57, figs. 6-10.

Chiloguembelina martini (PIJPERS), BECKMANN, 1957, p. 89, pl. 21, fig. 14, text-fig. 14, nos. 9-11, 14-18, 20-23.

Chiloguembelina martini (PIJPERS), RAJU, 1968, pl. 1, fig. 12.

Remarks: This species from Cauvery basin was earlier illustrated by the present author (RAJU, 1968).

Stratigraphic range: BECKMANN (1957) gave the range of this species from the Globorotalia aragonensis zone o the top of G. cocoaensis zone (=G. cerroazulensis zone) in Trinidad. In Cauvery basin this species is recorded from the Globorotalia spinuloinflata subzone (Lower Eocene) to the top of G. cerroazulensis zone.

## References

- AKERS, W. H. (1955): Some planktonic foraminifera of the American Gulf Coast and suggested correlations with the Caribbean Tertiary. — J. Paleontol., 29, pp. 647—664. Menasha.
- BANDY, O. L. (1949): Eocene and Oligocene foraminifera from Little Stave Creek, Clarke County, Alabama. — Bull. Amer. Paleontol., 32, no. 131, pp. 1—211. Ithaca.
- BANDY, O. L. (1964): Cenozoic planktonic foraminiferal zonation. Micropaleontology, 10, pp. 1—17. New York.
- BANNER, F. T., & BLOW, W. H. (1959): The classification and stratigraphical distribution of the Globigerinaceae. — Palaeontology, 2, pp. 1–27. London.
- BANNER, F. T., & BLOW, W. H. (1965): Progress in the planktonic foraminiferal biostratigraphy of the Neogene. — Nature, 208, no. 5016, pp. 1164—1166. London.
- BAUMANN, P., & ROTH, P. H. (1969): Zonierung des Obereozans und Oligozans des Monte Cagnero (Zentralapennin) mit planktonischen Foraminiferen und Nannoplankton. — Eclogae Geol. Helv., 62, pp. 303—323. Basel.
- BECKMANN, J. P. (1957): Chiloguembelina LOEBLICH & TAPPAN, and related Foraminifera from the Lower Tertiary of Trinidad, B. W. I. — U. S. Nat. Mus. Bull., 215, pp. 83—96. Washington.

- BERGGREN, W. A. (1969): Paleogene biostratigraphy and planktonic foraminifera from Northern Europe. -- Proc. I. Int. Conference on Planktonic Microfossils, Geneva, 1967, 1, pp. 121-160. E. J. Brill, Leiden.
- BLOW, W. H. (1969): Late Middle Eocene to Recent planktonic foraminiferal biostratigraphy. — Proc. I. Int. Conference on Planktonic Microfossils, Geneva, 1967, 1, pp. 199–422, pls. 1–54. E. J. Brill, Leiden.
- BLOW, W. H., & BANNER, F. T. (1962) in: EAMES et al.: Fundamentals of Mid-Tertiary Stratigraphical Correlations. — Pp. 61—151, pls. 8—17. Cambridge Univ. Press, Cambridge.
- BLOW, W. H., & SAITO, T. (1968): The morphology and taxonomy of Globigerina mexicana CUSHMAN, 1925. — Micropaleontology, 14, pp. 357-360. New York.
- BOLLI, H. M. (1957 a): Planktonic Foraminifera from the Oligocene Miocene Cipero and Lengua Formations of Trinidad, B. W. I. — U. S. Nat. Mus. Bull., 215, pp. 97—127. Washington.
- BOLLI, H. M. (1957b): Planktonic Foraminifera from the Eocene Navet and San Fernando Formations of Trinidad, B. W. I. — U. S. Nat. Mus. Bull., 215, pp. 155—172. Washington.
- BOLLI, H. M. (1962): Globigerinopsis, a new genus of the foraminiferal family Globigerinidae. Eclogae Geol. Helv., 55, pp. 280–284. Basel.
- BOLLI, H. M. (1966): Zonation of the Cretaceous to Pliocene Marine Sediments based on
- planktonic Foraminifera. Bol. Inform. Asoc. Venezolana Geol. Min. Petr., 9, pp. 3-32. Caracas.
- BOLLI, H. M., LOEBLICH, A. R. jr., & TAPPAN, H. (1957): Planktonic foraminiferal families Hantkeninidae, Orbulinidae, Globorotaliidae and Globotruncanidae. — U. S. Nat. Mus. Bull., 215, pp. 3–50. Washington.
- BORSETTI, M. E. (1959): Tre nuovi foraminiferi planctonici dell'Oligocene Piacentino. Ann. Mus. Geol. Bologna, Giorn. Geol., 27, pp. 205–212, pl. 13. Bologna.
- BRABB, E. E. (1969): Eocene Colloquium. Geotimes, 14, pp. 19-20. Washington.
- BRÖNNIMANN, P. (1950) The genus *Hantkenina* Cushman in Trinidad and Barbados, B. W. I. — J. Paleontol., 24, pp. 397—420. Menasha.
- CATI, F., and others (1968): Biostratigrafia del Neogen mediterraneo basata sui foraminiferi planctonoci. -- Boll. Soc. Geol. Ital., 87, pp. 491-503.
- CLARKE, W. J., & BLOW, W. H. (1969): The inter-relationships of some Late Eocene, Oligocene and Miocene larger Foraminifera and planktonic biostratigraphic indices. — Proc. I. Int. Conference on Planktonic Microfossils, Geneva, 1967, 2, pp. 82—97. E. J. Brill, Leiden.
- CUSHMAN, J. A., & BERMUDEZ, J. (1937): Further New species of Foraminifera from the Eocene of Cuba. — Contr. Cushman Lab. Foram. Res., 13, pp. 1–29. Ithaca.
- DIENI, I., & PROTO DECIMA, F. (1964) Cribrohantkenina ed altri Hantkeninidae nell'Eocene superiore di Castelnuovo (Colli Euganei). — Riv. Ital. Paleontol., 70, pp. 555—592. Milano.
- DROOGER, C. W. (1964): Problems of Mid-Tertiary stratigraphic interpretation. Micropaleontology, 10, pp. 369—374. New York.
- EAMES, F. E., BANNER, F. T., BLOW, W. H., & CLARKE, W. J. (1962): Fundamentals of mid-Tertiary stratigraphical correlations. -- Cambridge Univ. Press, pp. 1–162. Cambridge.
- ELLIS, B. F., & MESSINA, A. R. (1940) et seq.): A Catalogue of Foraminifera. Amer. Mus. Nat. Hist., Spec. Publ. New York.
- GUHA, D. K., MADAN MOHAN, KUMAR, P., IYENGAR, R., & RAJU, D. S. N. (1965): Marine Neogene microfauna from Karaikal, South India. — Bull. Geol. M. M. Soc. India, 34, pp. 1---3. Calcutta.
- JENKINS, D. J. (1960): Planktonic Foraminifera from the Lake Entrance Oil Shaft, Victoria, Australia. — Micropaleontology, 6, pp. 345—371. New York.
- JENKINS, D. J. (1965): Planktonic foraminiferal zones and new taxa from the Danian to Lower Miocene of New Zealand. — N. Z. J. Geol. Geophys., 8, pp. 1088—1126. Wellington.

- JENKINS, D. J. (1966): Planktonic foraminifera from the type Aquitanian—Burdigalian of France. — Contr. Cushman Found. Foram. Res., 17, pp. 1--15. Ithaca.
- MARKS, P., & WEBB, P. (1966): Discussion to the Symposium on micropaleontological lineages and zones. — Proc. III. Session CMNS, Berne, 1964, pp. 140—145. E. J. Brill, Leiden.
- NAGAPPA, Y. (1959): Foraminiferal biostratigraphy of the Cretaceous—Eocene succession in India-Pakistan-Burma region. — Micropaleontology, 5, pp. 145—192. New York.
- PAPP, A., and others (1968): Zur Nomenklatur des Neogens Österreich (mit einer stratigraphischen Tabelle). With English text by E. J. Tynan. — Verh. Geol. B.-A., 1968, pp. 1–27.
- RAJU, D. S. N. (1966): Note on Cenozoic planktonic foraminifera from Cauvery basin, S. India. – Bull. O. N. G. Commission, India, 3, pp. 14–19. Dehradun.
- RAJU, D. S. N. (1967): Late Cretaceous to early Miocene planktonic foraminifera from Cauvery basin, India and some intercontinental correlations. — Abstract, Planktonic Conference, Geneva, 1967, pp. 120—121. Geneva.
- RAJU, D. S. N. (1968): Eocene—Oligocene planktonic foraminiferal biostratigraphy of Cauvery basin, S. India. — Mem. Geol. Soc. India, 2, pp. 286—299, pls. 1—4. Bangalore.
- RAMANATHAN, S. (1968): Stratigraphy of Cauvery basin with reference to its Oil Prospects. — Mem. Geol. Soc. India. 2, pp. 153–167. Bangalore.
- REISS, Z., & GVIRTZMANN, G. (1966): Subsurface Neogene stratigraphy of Israel. Proc. III Session, CMNS, Berne, 1964, pp. 312—324, pls. 88—99. E. J. Brill, Leiden.
- SAITO, T. (1963): Miocene planktonic foraminifera from Honshu, Japan. Tohoku Univ., Sci. Rept., Ser. 2 (Geol.), 35, pp. 123-209. Sendai.
- SCHMID, M. E. (1967): Zwei neue planktonische Foraminiferen aus dem Badener Tegel von Sooß, NO. — Ann. Naturhist. Mus. Wien, 71, pp. 347—352. Wien.
- SPRAUL, G. L. (1963): Current status of the Upper Eocene foraminiferal guide fossil, Cribrobantkenina. — J. Paleontol., 37, pp. 336—370. Tulsa.
- SUBBOTINA, N. N. (1953): Fossil foraminifera of the U.S.S.R. (Globigerinidae, Hantkeninidae and Globorothaliidae). Vses. Neft. Nauchno-Issled. Geol.-Razved. Inst., Trudy, n. ser., 76, pp. 1–296. Moskau.
- THALMANN, H. E. (1942): Foraminiferal genus Hantkenina and its subgenera. Amer. J. Sci., 240, pp. 809–820. New Haven.
- VERVLOET, C. C. (1966): Stratigraphical and micropaleontological data on the Tertiary of Southern Piemont (Northern Italy). — Schotanus & Jens, Utrecht NV, pp. 1—88. Utrecht.
- WADE, M. (1964): Application of lineage concept to biostratigraphic zoning based on planktonic foraminifera. — Micropaleontology, 10, pp. 273—290. New York.

(All figures: a = ventral view, b = dorsal view, c = side view)

#### PLATE I

- Figures 1 a—c. Globigerina ampliapertura BOLLI, from G. gortanii zone, core sample at 1800 m. in NGT-1.
- Figures 2, 3 a, b, 4. Globigerina angulisuturalis BOLLI, from G. kugleri/G.primordius zone, core sample at 1421.5 m. in KKL-2.
- Figures 5 a, b. Globigerina ciperoensis BOLLI, from G. kugleri/G. primordius zone, core sample at 1421.5 m. in KKL-2.
- Figure 6. Globigerina angustiumbilicata BOLLI, from G. kugleri/G. primordius zone, core sample at 1126.5 m. in KKL-2.
- Figures 7 a, b. Globigerina cf. pseudoampliapertura BLOW & BANNER, from G. gortanii zone, core sample at 1800 m. in NGT-1.
- Figure 8. Globigerina angiporoides HORNIBROOK, from G. gortanii zone, core sample at 1800 m. in NGT-1.





















#### PLATE II

- Figures 1 a, b, 2. Globigerina gortanii gortanii (BORSETTI), from G. sastrii zone, core sample at 1579 m. in KKL-4.
- Figures 3 a, b. Globigerina pseudocorpulenta CHALILOV, from G. mexicana zone, core sample at 1904.2 m. in NGT-1.
- Figures 4 a, b, 5 a, b. Globigerina gortanii praeturritilina BLOW & BANNER, from G. mexicana zone, core sample at 1904.2 m. in NGT-1.
- Figures 6 a, b. Globigerina cf. rubriformis (SUBBOTINA), from G. gortanii zone, core sample at 1800 m. in NGT-1.

PLATE II



























#### PLATE III

Figures 1 a, b. Globigerina sp., from G. mexicana zone, core sample at 1904.2 m. in NGT-1.

Figures 3 a, b. Globigerina tripartita tripartita KOCH, from G. sastrii zone, core sample at 1579 m. in KKL-4.

- Figures 2, 4 a, b. Globigerina tripartita robri BOLLI, from G. sastrii zone, core sample at 1582 m. in KKL-4.
- Figures 5 a,b, 6. Globigerina sellii BORSETTI, 5 a, b from G. sastrii zone, core sample at 1582 m. in KKL-4; 6 from cutting sample at 1515 m. in KKL-2.

2

3a

47





5b

lmm.



4 a



#### PLATE IV

Figures 1 a—c. Globigerina sastrii n. sp., holotype from core sample at 1579 m. in KKL-4, from G. sastrii zone.

Figures 2 a-c. Globigerina sastrii n. sp., paratype from core sample at 932 m. in TPD-1.













Jahrb. Geol. B. A., 1971, Sonderband 17

#### PLATE V

- Figures 1 a-c. Globigerina corpulenta SUBBOTINA, from G. mexicana zone, core sample at 1904.2 m. in NGT-1.
- Figures 2 a-c, 3. Globigerina galavisi BERMUDEZ, from G. mexicana zone, core sample at 1904.2 m. in NGT-1.
- Figure 4. Globigerina venezuelana HEDBERG, from G. kugleri/G. primordius zone, core sample at 1126.5 m. in KKL-2. Figures 5 a-c. Globigerina woodi woodi JENKINS, from G. kugleri/G. primordius zone,
- core sample at 1421.5 m. in KKL-2.























%

#### PLATE VI

Figures 1 a-c. Globigerina venezuelana HEDBERG, from G. kugleri/G. primordius zone, core sample at 1126.5 m. in KKL-2.

Figures 2 a, b, 3 a, b. Globigerinita dissimilis (CUSHMAN & BERMUDEZ), from G. kugleri/G. primordius zone, core sample at 1126.5 m. in KKL-2.

Figures 4 a—c. Globigerinita stainforthi (BOLLI, LOEBLICH & TAPPAN), from G. kngleri/G. primordius zone, core sample at 1126.5 m. in KKL-2.

Figures 5 a, b. Globorotaloides suteri BOLLI, from a cutting sample at 1463 m. in KKL-1.

PLATE VI















4c







#### PLATE VII

- Figures 1 a, b, 2 a, b. Globigerinoides primordius BLOW & BANNER, from G. kugleri/G. primordius zone, core sample at 1421.5 m. in KKL-2.
- Figures 3 a, b. Globigerinoides trilobus altiaperturus BOLLI, from core sample at 639 m. in KKL-4.
- Figures 4 a, b. Globigerinoides trilobus trilobus (REUSS), from core sample at 495 m. in TPD-1.
- Figures 5 a, b. Globigerinoides cf. subquadratus BRÖNNIMANN, from core sample at 639 m. in KKL-4.
- Figures 6 a, b. Globoquadrina altispira globosa BOLLI, from the upper part of G. kugleri/G. primordius zone, core sample at 1126.5 m. in KKL-2.

.

PLATE VII





١a

















0,5



lmm.





#### PLATE VIII

- Figures 1 a-c. Globigerinopsis gubai n. sp., holotype from core sample at 1126.5 m. in KKL-2, upper part of the G. kugleri/G. primordius zone.
- Figures 2 a-c, 3 a-c Globigerinopsis guhai n. sp., paratypes from the same level as the holotype.
- Figures 4 a .-- c. Globoquadrina praedehiscens BLOW & BANNER, from G. kugleri/G. primordius zone, core sample at 1126.5 m. in KKL-2.

PLATE VIII

























#### PLATE IX

Figures 1, 2. Globoquadrina praedehiscens BLOW & BANNER, from G. kugleri/G. primordius zone, core sample at 1126.5 m. in KKL-2.

- Figures 3, 4 a, b. Globoquadrina debiscens (CHAPMAN, PARR & COLLINS), from core sample at 495 m. in TPD-1.
- Figures 5 a, b. Globoquadrina cf. larmeui AKERS, from G. angulisuturalis zone, core sample at 1495.5 m. in KKL-2.
- Figures 6 a.-c. Globorotalia peripheroronda BLOW & BANNER, from core sample at 495 m. in TPD-1.
- Figures 7 a—c. Globorotalia increbescens (BANDY), from G. gortanii zone, core sample at 1800 m. in NGT-1.

PLATE IX



#### PLATE X

- Figures 1 a, b, 2 a, b. Globorotalia kugleri BOLLI, from G. kugleri/G. primordius zone, core sample at 1126.5 m. in KKL-2.
- Figures 3 a, b. Globorotalia opima nana Bolli, from G. sastrii zone, core sample at 1582 m. in KKL-4.
- Figures 4, 5 a—c. Globorotalia siakensis (LEROY), from G. kugleri/G. primordius zone, core sample at 1126.5 m. in KKL-2.
- Figures 6 a—c. Globorotalia obesa BOLLI, from G. kugleri/G. primordius zone, core sample at 1126.5 m. in KKL-2.

PLATE X







**0**∙2 0-3mm. 0;1

















Q



imm.

0.5





#### PLATE XI

- Figures 1—6. Globigerapsis mexicana (CUSHMAN), all from G. mexicana zone, core sample at 1904.2 m. in NGT-1, 1—4 juvenile forms with one to two supplementary apertures.
- Figures 7 a, b. Globigerapsis cf. tropicalis BLOW & BANNER, from G. mexicana zone, core sample at 1904.2 m. in NGT-1.
- Figure 8. Hantkenina alabamensis CUSHMAN, from G. mexicana zone, core sample at 1904.2 m. in NGT-1.
- Figure 9 a, b. Hantkenina alabamensis CUSHMAN, from a cutting sample at 1580 m. in KKL-1. A from with twin spines.

PLATE XI

















#### PLATE XII

÷,

- Figures 1 a, b. Hantkenina cf. thalmanni BRÖNNIMANN, from G. mexicana zone, core sample at 1904.2 m. in NGT-1.
- Figure 2. Hantkenina alabamensis CUSHMAN, from G. cerroazulensis zone in KKL-4.

.

- Figures 3, 4, 5. Hantkenina trinitatensis BRÖNNIMANN, from G. mexicana zone, core sample at 1904.2 m. in NGT-1.
- Figures 6, 7, 8. Hantkenina sp., from G. cerroazulensis zone, core sample at 1716 m. in KKL-4.
- Figures 9 a, b, 10. Hantkenina suprasuturalis BRÖNNIMANN, from G. mexicana zone, core sample at 1904.2 m. in NGT-1.



Jahrb. Geol. B. A., 1971, Sonderband 17

#### PLATE XIII

- Figure 1. Hantkenina suprasuturalis BRÖNNMANN, from G. mexicana zone, core sample at 1904.2 m. in NGT-1.
- Figures 2, 7. Cribrohantkenina inflata (Howe), from G. cerroazulensis zone, core sample at 1690 m. in KKL-4.
- Figures 3, 4, 5, 6, 8 a, b. Cribrohantkenina inflata (Howe), from G. cerroazulensis zone, all from a core sample at 1716 m. in KKL-4.
- Figures 9 a, b. Pseudohastigerina micra (COLE), from G. mexicana zone, core sample at 1904.2 m. in NGT-1.
- Figures 10, 11. Hantkenina sp., from G. cerroazulensis zone, core sample at 1690 m. in KKL-4, juvenile forms with the spines bending backward.

PLATE XIII

0;5

0;3 mm.

Ó-2

Q

(mm

<u>ộ</u>

# **ZOBODAT - www.zobodat.at**

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: <u>Jahrbuch der Geologischen Bundesanstalt</u> <u>Sonderbände</u>

Jahr/Year: 1970

Band/Volume: 17

Autor(en)/Author(s): Raju D. S. N.

Artikel/Article: <u>Upper Eocene to Early Miocene Planktonic Foraminifera</u> from the Subsurface Sediments in Cauvery Basin, South India 7-67