

Diagnose einer neuen Arca.

Von

W. K o b e l t.

Arca Amaliae n. sp.

Concha magna, irregulariter ovata, antice inflata, postice subattenuata, solida, ponderosa, tumida, aequivalvis, inaequilatera, alba, hic illic fuscescenti obscure tincta et annulata. Margo superior subarcuatus, utrimque declivis, cum anteriore truncato, dein declivi angulum compressum auriculatum formans, inferior primum bene rotundatus, dein subretusus, posterior declivis, ad apicem acute rotundatus. Superficies costis radiantibus crassis 36, marginem versus fasciculatis, quam interstitia profunda plus quam duplo latioribus, hic illic obscure nodosis striisque concentricis in interstitiis obsoletis sculpta. Umbones ante $\frac{1}{3}$ longitudinis positi, pertumidi, apicibus regulariter intortis, sat approximatis; area mediocris, profunde declivis, transversim regulariter sulcata, sulcis dentibus cardinis respondentibus. Latus internum leviter fuscescens, limbo incrassato albo extra lineam pallealem, margine valde crenato-dentato.

Long. 90, alt. 67, crass. 60 mm.

Arca Amaliae Kobelt in Martini-Chemnitz ed. II. (VIII. 2)
t. 8 fig. 1. 2.

Hab. — ? Exstat in Museo Senckenbergiano.



ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Jahrbücher der Deutschen Malakozoologischen Gesellschaft](#)

Jahr/Year: 1887

Band/Volume: [14](#)

Autor(en)/Author(s): Kobelt Wilhelm

Artikel/Article: [Diagnose einer neuen Arca. 374](#)