

Jb. Geol. B.-A.	ISSN 0016-7800	Band 134	Heft 2	S. 257-270	Wien, Oktober 1991
-----------------	----------------	----------	--------	------------	--------------------

Evolution of Middle and Upper Cretaceous Floras in Central and Western Europe

By ERWIN KNOBLOCH & DIETER HANS MAI*)

With 1 Text-Figure, 4 Tables and 4 Plates

Paleobotany
 Cenomanian
 Maestrichtian
 Fruits
 Seeds
 Evolution
 Paleofloristic Periods

Contents

Zusammenfassung	257
Abstract	257
1. Introduction	257
2. Documentation of Modern Genera and Families in the Middle and Upper Cretaceous	258
3. Evolutionary Paleofloristic Periods from the Cenomanian to the Maestrichtian	259
3.1. Cenomanian	259
3.2. Upper Turonian – Santonian	261
3.3. Santonian – Campanian	261
3.4. Maestrichtian	262
4. Conclusions	262
References	270

Evolution von Mittel- und Oberkreideflören in Zentral- und Westeuropa

Zusammenfassung

Die Arbeit gibt eine Zusammenfassung über inkohlte Samen und Früchte aus der mittleren und oberen Kreide von West- und Mitteleuropa. Zahlreiche rezente paläotropische Gattungen und Familien wurden nachgewiesen: manche Gattungen sind vollkommen ausgestorben. In der Mittel- und Oberkreide werden 4 Entwicklungsabschnitte der Flora unterschieden: 1. Cenoman, 2. Ober-Turon-Santon, 3. Santon-Campan, 4. Maastricht.

Abstract

A summary is presented of carbonized seeds and fruits from the Middle and Upper Cretaceous of Western and Middle Europe. Numerous living paleotropical genera and families are mentioned. Some of the genera studied are totally extinct. Four evolutionary intervals could be recognized among the Middle and Upper Cretaceous floras: 1. Cenomanian, 2. Upper Turonian-Santonian, 3. Santonian-Campanian, 4. Maestrichtian.


1. Introduction

During the years 1975–1983, we studied rich collections of carbonized diaspores from the Middle and Upper Cretaceous of Western and Middle Europe. Geographically speaking, we investigated localities in Czechoslovakia, Germany, Austria, Hungary, Poland and the Netherlands. The fruits and seeds were mainly

sieved out from greyish clays and silts. They are known from freshwater and marine sediments.

In many cases, we are able to clarify the systematic position of the plant fossils. In this way, it is possible to document the presence of many genera and families in the Cretaceous for the first time and to provide new information on the origin of the angiosperms.

*) Author's addresses: Dr. ERWIN KNOBLOCH, Geological Survey, Malostranské nám. 19, ČSFR-11821 Praha 1; Doz. Dr. sc. nat. DIETER HANS MAI, Museum für Naturkunde, Bereich Botanik-Arboretum, Späthstraße 80–81, D-1192 Berlin-Baumschulenweg.



Text-Fig. 1:

The most important localities, formations and regions with Cretaceous fruits and seeds in Central and Western Europe.

1 = South-Limburg, Aachen Formation, Upper Santonian-?Lower Campanian, The Netherlands; 2 = Aachen, Aachen Formation, Upper Santonian or ?Lower Campanian, Germany; 3 = Walbeck, Maestrichtian, Germany; 4 = Eisleben, Maastrichtian, Germany; 5 = Cenomanian of West and Central Bohemia, Peruc Member, Czechoslovakia; 6 = Cenomanian of Moravia, Peruc Member, Czechoslovakia; 7 = Slezské Pavlovice, bore-hole OS-5, Upper Turonian-Lower Coniacian, Czechoslovakia; 8 = Klikov Formation, Upper Turonian-Upper Santonian, South Bohemia, Czechoslovakia; 9 = Flysh in the Moravian-Silesian Bezkydy Mts., Campanian-Maestrichtian, Czechoslovakia; 10 = Kössen, Niederndorf, Gosau Formation, Maestrichtian, Austria; 11 = Vienna, Flysh, Sievering Formation, Campanian-Maestrichtian, Austria; 12 = Bakony Mts., Ajka Formation, Upper Santonian - Lower Campanian, Hungary.

The study of angiosperm fruits and seeds from the Middle to Upper Cretaceous (Cenomanian-Maestrichtian) of Europe has resulted in the recognition of 270 species and 24 families and has established the existence in the Cretaceous of

- 1) many modern genera,
- 2) many extinct genera referable to Recent families and
- 3) extinct genera that cannot be referred to any existing family.

2. Documentation of Modern Genera and Families in the Middle and Upper Cretaceous

The Magnoliaceae are represented by fruits with 1-2 seeds (*Liriodendron* - Pl. 2, Fig. 7, *Liriodendroidea*) as well as fruits with more seeds (*Padragkutia* - Pl. 3, Fig. 7; *Manglietia*). *Padragkutia* has up to 17 seeds per locule. *Liriodendron* is documented by seeds for the first time in the Senonian of southern Bohemia and this indicates that many leaf fossils previously assigned to the genus may be correctly identified.

The new genus *Prototinomiscium* (Pl. 3, Fig. 1) demonstrates that extinct members of the Menispermaceae are not confined only to the British Lower Tertiary, but that they also occurred in the Senonian of continental Europe. The same is the case for *Typha* (Typhaceae) and *Spirematospermum* (Zingiberaceae).

The Hamamelidaceae were rather abundant during the Cretaceous and are represented in the Maestrichtian by the modern genera *Disanthus* and *Rhodoleia* (Pl. 4, Fig. 14) and in the Cenomanian to Maestrichtian by the extinct genus *Klikovispermum* (Pl. 1, Figs. 8 and 14, Pl. 2, Fig. 11) which includes 17 species. *Klikovispermum* is closely related to *Rhodoleia* and has 12-16 anatropous and semicircular seeds per locule.

The Platanaceae, which are of widespread occurrence in Cretaceous floras of the northern hemisphere, are mainly represented by leaves, fruiting organs being uncommon.

The Urticaceae are also very abundant and have been demonstrated with certainty in the Cretaceous;

nutlets assigned to the genus *Boehmeria* (Pl. 1, Figs. 13 and 16, Pl. 2, Fig. 9) which today is mainly restricted to tropical areas, occur as early as the Cenomanian. The genus is represented by 9 species among the Mid- to Upper Cretaceous floras of Europe. In addition to *Boehmeria*, the Cretaceous Urticaceae include species assigned to the modern genera *Debregeasia* and *Memorialis* (Pl. 2, Fig. 1, Pl. 3, Fig. 16) and to the extinct genus *Urticoidea* (Pl. 3, Fig. 6).

Three extinct genera possibly related to the Juglandaceae are *Calathiocarpus* (Pl. 2, Fig. 13, Pl. 3, Fig. 11), *Caryanthus* (Pl. 2, Figs. 3 and 10) and *Manningia*. They are mostly characterized by the presence of a distinct hypanthium. *Caryanthus triasseris* (KNOBL.) KNOBL. et MAI is the most common fossil in the Cretaceous of Europe.

The Theaceae, which by the Tertiary are represented by numerous genera and species, appear to have had a maximum evolution in the Cretaceous. Thus, the modern genus *Eurya* (Pl. 3, Fig. 9) is represented by 3 species and *Visnea* by 1 species. The extinct genus *Palaeoschima* (Pl. 4, Fig. 9), which is closely related to the modern genus *Schima*, also exhibits some characteristics related to the Ericaceae - Andromedae, and the extinct genus *Protopvisnea* (Pl. 2, Figs. 4 and 8, Pl. 3, Fig. 4), that is closely related to the modern genus *Visnea*, also exhibits many features characteristic of other families.

In the uppermost Cretaceous (Maestrichtian) the Pentaphylacaceae are documented by fruits assigned to the extinct genus *Allericarpus* and the modern genus *Pentaphylax* (Pl. 4, Fig. 11).

The Ulmaceae family is represented by the genera *Aphananthe* (Pl. 4, Fig. 3) and *Gironniera*, and the Saurauiaaceae family by two species of the genus *Saurauia* (Pl. 3, Fig. 15).

Many of the extinct genera from the Cretaceous that we have recognized show relationships to various genera and families, rather than to a single taxon. This is also the case for the extinct genus *Discoclethra* (Pl. 3, Fig. 3, which shows many characteristics in common with the Clethraceae (i.e. placentation, desiccation, deeply 5-lobed and persistent calyx and seed shape) but also exhibits many characteristics related to other families, e.g. Celastraceae (discus, valves),

Hamamelidaceae (style), Cunoniaceae (discus, style) and Ericaceae (seed wall).

Several kinds of capsular fruits document the presence in the Cretaceous of Ericaceae (*Leucothoe* – Pl. 4, Fig. 5, *Viticocarpum*) and Cyrillaceae (*Diplycosiopsis*) and related taxa (*Epacridicarpum*, *Purdiaeopsis* – Pl. 4, Fig. 2 and *Valvaecarpus* – Pl. 3, Fig. 12, Pl. 4, Fig. 6).

Seeds from the Upper Cretaceous of southern Bohemia show a close relationship with the modern *Aldrovanda* (Pl. 2, Fig. 5) and thus indicate the presence of the Droseraceae in the Cretaceous.

The two well-known extinct Rutaceous genera from the Lower Tertiary of England, *Rutaspermum* and *Shrubsolea*, have also been documented in the Cretaceous, and the Sapindaceae are represented therein by the extinct genus *Sapindospermum* (Pl. 2, Fig. 6).

While the genera *Sabia* (Pl. 3, Fig. 14; Pl. 4, Fig. 8) and *Meliosma* (Sabiaceae) are already known from the Tertiary, we have now described (see KNOBLOCH & MAI, 1984, 1986) a new genus from the Cretaceous, *Insioticarpus*, with 4 species, that extends the fossil record of the family back to the Cenomanian.

The Mastixiaceae, that are well represented by many genera in the Tertiary, are known only from the uppermost Cretaceous (Maestrichtian) by the genera *Beckettia* (Pl. 4, Fig. 12), *Mastixicarpum* (Pl. 4, Fig. 1), *Eomasitia* and *Mastixiopsis*. The same is the case for the Araliaceae (*Aralia*, *Acanthopanax* – Pl. 4, Fig. 7) that have been suggested, based on leaf fossils (Potomac Formation, USA), as early as the Lower Cretaceous, and also the Icacinaceae (*Icacinicarya*, *Iodes* – Pl. 4, Fig. 4) and the Aquifoliaceae (*Ilex*).

In addition to the Cretaceous fossils referable to Recent genera and families and to extinct genera related to Recent families, we have described (KNOBLOCH & MAI, 1984) 26 genera that cannot so far be attributed to any modern family. This could be explained by incomplete knowledge of modern carpology or by the fact that they belonged to families that are now extinct. A large number of species have been accommodated into these extinct genera. Some of them, with their characteristics, are listed below: *Bicameria* (Pl. 3, Fig. 8): fruits bilocular with a median septum (5 species); *Discosemen* (Pl. 1, Fig. 12): seed anatropous, disk-shaped and with several cell-layers of the seed wall (6 species); *Laramisemen* (Pl. 1, Fig. 17, Pl. 2, Fig. 2, Pl. 3, Fig. 13): seed anatropous, raphe forming an edge starting below hilum (30 species); *Spirella* (Pl. 1, Figs. 1–4 and 9): seed with longitudinal ridges (18 species); *Walbeckia* (Pl. 3, Figs. 5 and 10): one-loculed, oval two-valved fruit (10 species); *Eislebenia* (Pl. 4, Fig. 13): 1 seed with a prominent tegmen (2 species).

3. Evolutionary Paleofloristic Periods from the Cenomanian to the Maestrichtian

Fossil seeds and fruits are not rare in the Cretaceous of Middle and Western Europe. They were recorded at 176 localities (mostly boreholes). The richest layers are known from the Klikov Formation (South Bohemia), Walbeck and Eisleben (Germany) and from the Aachen Formation (Germany, The Netherlands).

A great part of the plant fossils could be assigned to living genera and families. 46 new genera were defined

by the authors (KNOBLOCH & MAI, 1984, 1986). 26 genera are totally extinct and 20 show some affinities to recent families. Some of the extinct genera are also known from the Cretaceous to the Paleocene (i.e. *Barclayopsis*, *Klikovispermum*, *Valvaecarpus*, *Mastixiopsis*, *Spirella*, *Normacarpus*, *Laramisemen*, *Rusavia*, *Walbeckia* – MAI, 1987). The number of extinct genera decreases from the Cenomanian to the Paleocene.

Up to now, four periods could be recognized in the evolution of Cretaceous floras:

- 1) Cenomanian
- 2) Upper Turonian–Santonian
- 3) Santonian–Campanian
- 4) Maestrichtian.

These 4 intervals include only some basic evolutionary tendencies. Even though the number of species studied is sufficient, our knowledge is still limited. We do not know of any rich plant collections from the Turonian, Campanian or Lower Maestrichtian; we do not know precisely if the Klikov Formation belongs to the whole stratigraphic range Upper Turonian–Upper Santonian (KNOBLOCH, 1985) or whether the Aachen Formation belongs only to the Upper Santonian or to the Lower Campanian (ALBERS & FELDER, 1979). The stratigraphic positions of some localities from the flysch of Moravia are not clear. The stratigraphic position of the Eisleben locality in Germany was ranged to the Maestrichtian, but probably belongs to the Paleocene. The stratigraphic range of some species during the Cretaceous is not known up to now.

3.1. Cenomanian

(Table 1, Plate 1)

In the Cenomanian the angiosperms show a significant worldwide „explosion“ in their evolution. Leaves of angiosperms are predominant in most Bohemian fossil sites from the Cenomanian. Only 24 species of fossil fruits and seeds have been recorded in the Cenomanian sediments of Bohemia and Moravia.

Contrary to the younger floras, the flora from the Cenomanian documented by seeds and fruits shows a

Table 1.
Angiospermous seeds and fruits from the Cenomanian
(Peruc Member, Czechoslovakia)

Hamamelidaceae	<i>Klikovispermum micromalechii</i> KNOBL. et MAI <i>Klikovispermum pragense</i> KNOBL. et MAI <i>Klikovispermum semicirculare</i> KNOBL. et MAI <i>Klikovispermum verrucosum</i> KNOBL. et MAI
Platanaceae	<i>Platanus laevis</i> (VEL) VEL
Urticaceae	<i>Boehmeria cenomanica</i> KNOBL. et MAI
?Juglandaceae	<i>Caryanthus triasseri</i> (KNOBL.) KNOBL. et MAI
Sabiaceae	<i>Insioticarpus moravicus</i> KNOBL. et MAI
Magnoliophyta fam. indet.	<i>Brnikia ovalis</i> KNOBL. et MAI <i>Carpolithus nyssiformis</i> KNOBL. et MAI <i>Carpolithus ovalis</i> KNOBL. et MAI <i>Carpolithus pecinovensis</i> KNOBL. et MAI <i>Curvospermum marketense</i> KNOBL. et MAI <i>Discosemen faveolatum</i> KNOBL. et MAI <i>Laramisemen hypericoides</i> KNOBL. et MAI <i>Laramisemen minor</i> KNOBL. et MAI
Stemonaceae	<i>Spirella margaritifera</i> KNOBL. et MAI <i>Spirella minutissima</i> KNOBL. et MAI <i>Spirella pragensis</i> KNOBL. et MAI <i>Viniklaria bohemica</i> KNOBL. et MAI

Table 2.

Important species from the Upper Turonian – Santonian and their distribution in younger formations.

OS-5 = Slezské Pavlovice borehole in southern Silesia, Upper Turonian – Lower Coniacian, Czechoslovakia; Klikov Formation = Upper Turonian – Santonian, Czechoslovakia; Aachen Formation = Upper Santonian or ?Lower Campanian; Flysch in northern Moravia and Vienna = Campanian, Maestrichtian, Austria, Czechoslovakia; Eisleben = Maestrichtian, Germany. Specimens in one sample: A = 1, B = 2–15, C = 16–100, D = more than 100.

		OS-5	Klikov Fm.	Aachen Fm.	Flysh	Eisleben
Magnoliaceae	<i>Liriodendron papilliformis</i> KNOBL. et MAI		A			
Sapindaceae	<i>Sapindospermum nitidum</i> KNOBL. et MAI		A			
?Juglandaceae	<i>Caryanthus deltoides</i> (KNOBL.) KNOBL. et MAI	A	A-C	B	?A	
	<i>Caryanthus trebecensis</i> KNOBL. et MAI	A-B	A-D	B		
Hamamelidaceae	<i>Klikovispermum malechii</i> (KNOBL.) KNOBL. et MAI	A	A-C	B		
	<i>Klikovispermum bohemicum</i> KNOBL. et MAI	A	A			
Urticaceae	<i>Boehmeria cyrokyi</i> (KNOBL.) KNOBL. et MAI	A	A-D	?A	B	B
?Juglandaceae	<i>Calathiocarpus minimus</i> KNOBL. et MAI		A-C	A-B		
Magnoliophyta fam. indet.	<i>Laramisemen microcanaliculatum</i> KNOBL. et MAI	A	A-B			
	<i>Laramisemen moravicum</i> KNOBL. et MAI	A				
Stemonaceae	<i>Spirella multistriata</i> (VANG) KNOBL. et MAI		A	B		C
	<i>Spirella multistriatiformis</i> KNOBL. et MAI	A		B		
Theaceae	<i>Protovisnea zahajensis</i> KNOBL. et MAI		A-B			
	<i>Protovisnea reticulata</i> KNOBL. et MAI		A-C			
Urticaceae	<i>Memorialis verrucosa</i> KNOBL. et MAI		A-B			
?Juglandaceae	<i>Caryanthus communis</i> KNOBL. et MAI		A-D			
Cyrillaceae	<i>Epacridicarpum cretaceum</i> (JUNG) KNOBL. et MAI		A-C		B	
Droseraceae	<i>Palaeoaldrovanda splendens</i> KNOBL. et MAI		A-B			
Magnoliophyta fam. indet.	<i>Bicameria holyi</i> KNOBL. et MAI		A-D	? A		
	<i>Budvaricarpus serialis</i> KNOBL. et MAI		A-D			
	<i>Vangerovia tricornuta</i> (KNOBL.). KNOBL. et MAI		A-B			B

most isolated and archaic character. A number of extinct genera dominate. Only the genera *Platanus* and *Boehmeria* are still living.

The gymnosperms (Taxodiaceae, Cupressaceae and Pinaceae) are characteristic for the Cenomanian of the Bohemian Massif. They are lacking

Table 3.

The most important species from the Santonian – Campanian and their distribution in older and younger formations. Klikov Formation = Upper Turonian – Santonian, Czechoslovakia; Aachen Formation = Upper Santonian – ?Lower Campanian, Germany, The Netherlands; Ajka Formation = Upper Santonian – Lower Campanian, Hungary; Flysch of Czechoslovakia and Austria = Campanian, Maestrichtian; Walbeck, Eisleben = Maestrichtian in Germany. Specimens in one sample: A = 1, B = 2–15, C = 16–100, D = more than 100.

		Klikov Fm.	Aachen Fm.	Ajka Fm.	Flysh	Walbeck Eisleben
Menispermaceae	<i>Prototinomiscium vangerowii</i> KNOBL. et MAI	A	C			
Magnoliaceae	<i>Padragkulia haasii</i> KNOBL. et MAI	B		C		
Urticaceae	<i>Memorialis rhenana</i> KNOBL. et MAI		B			
	<i>Urticoidea cucurbitoides</i> (VANG.) KNOBL. et MAI	A	B			
	<i>Urticoidea pedicellata</i> KNOBL. et MAI	B	A			
?Juglandaceae	<i>Calathiocarpus hollandicus</i> KNOBL. et MAI		B			
Theaceae	<i>Protovisnea cancellata</i> (VANG.) KNOBL. et MAI		B		?B	
Saurauiaceae	<i>Saurauia alenae</i> KNOBL. et MAI	A	A			A
Clethraceae	<i>Discoelethra maxima</i> KNOBL. et MAI		C			
	<i>Discoelethra polysperma</i> KNOBL. et MAI	A	C			
Cyrillaceae	<i>Epacridicarpum cretaceum</i> (JUNG) KNOBL. et MAI	C			B	B
	<i>Valvaecarpus debeyi</i> KNOBL. et MAI	B	C			
Magnoliophyta fam. indet.	<i>Bicameria orbicularis</i> (KNOBL.) KNOBL. et MAI	C	B			
	<i>Eislebenia goczani</i> KNOBL. et MAI		?B			
	<i>Laramisemen faboides</i> (VANG.) KNOBL. et MAI	A	B			
	<i>Operculispermum padragkulense</i> KNOBL. et MAI				C	
	<i>Walbeckia ajoensis</i> KNOBL. et MAI				B	
	<i>Quedlinburgia hungarica</i> KNOBL. et MAI				A	
Sabiaceae	<i>Sabia menispermioides</i> KNOBL. et MAI		A			
Theaceae	<i>Eurya crassitesta</i> KNOBL.				A-B	C
Magnoliophyta fam. indet.	<i>Walbeckia aquisgranensis</i> KNOBL. et MAI		B-C			

in the younger sediments (Turonian–Senonian). Many angiosperms represent monotypic and endemic genera.

The floristic characterization of this interval is made after the finds from the Peruc Member in Bohemia and Moravia (Czechoslovakia).

3.2. Upper Turonian–Santonian

(Table 2, Plate 2)

An increasing number of new extinct genera, but also a number of living genera are characteristic for the Upper Turonian and Santonian. As opposed to the Cenomanian, different species are known. Numerous species were hitherto endemic, as they were known only from the Klikov Formation.

The fruits and seeds from this interval are represented by many genera with ancient features. They belong to the following systematic units:

- **Juglandales**
?Budvaricarpus, Calathiocarpus, Caryanthus, Manningia
- **Hamamelidales**
Klikovispermum
- **Urticales**
Moroidea, Urticoidea
- **Magnoliales**
Padragkutia
- **Theales – Ericales**
Epacridicarpum, Discoclethra, Protovisena, Valvaecaropus

- **Celastrales**
Icacinicarya
- **Droserales**
Palaeoaldrovanda

The number of modern genera increases: Eurya, Lriodendron, Memorialis, Rhoiptelea, Sabia, Saurauia.

Some of the fossil genera became extinct between the Santonian and the Campanian.

The floristic characterization of this intervals was made mainly after the finds in the Klikov Formation in southern Bohemia (Upper Turonian–Santonian), the Merboltice Formation in northern Bohemia (Coniacian–Lower Santonian), the floristic remnants from Quedlingburg (Germany, Santonian) and the Slezské Pavlovice OS-5 borehole in southern Silesia (Upper Turonian–Lower Coniacian).

3.3. Santonian–Campanian

(Table 3, Plate 3)

This interval remains problematic. We know of no rich collections from Campanian sediments. This interval is defined accordig to the finds in the Aachen Formation (Upper Santonian or ?Lower Campanian), The Ajka Formation in the Bakony Mts. in Hungary (Upper Santonian–Lower Campanian) and some finds from the flysh of Moravia (Czechoslovakia) and Vienna (Austria). There are some common features between the Santonian–Campanian and the Uper Turonian–Santonian

Table 4.

Important species from the Maestrichtian (Walbeck, Eisleben) and their distribution in the flysh and Gosau Formation of Moravia and Austria (Campanian, Maestrichtian). The species mentioned are lacking in the Klikov and Aachen Formations. Specimens in one sample: A = 1, B = 2–15, C = 16–100, D = more than 100.

		Walbeck, Eisleben	Flysh, Gosau
Zingiberaceae	<i>Spirematospermum friedrichii</i> KNOBL. et MAI	B	
Magnoliaceae	<i>Manglietia hercynica</i> MAI <i>Prototinomiscium testudinarum</i> KNOBL. et MAI	B B	
Hamamelidaceae	<i>Rhodoleia cretacea</i> KNOBL. et MAI	B	
Ulmaceae	<i>Aphananthe cretacea</i> KNOBL. et MAI	B	
Urticaceae	<i>Boehmeria krutzschii</i> KNOBL. et MAI	C	
Theaceae	<i>Palaeoschima becvensis</i> (KNOBL.) KNOBL. et MAI	C	A–C
Cyrillaceae	<i>Valvaecarpus pterocaryaeformis</i> (KNOBL.) KNOBL. et MAI	C	A–D
Sabiaceae	<i>Sabia praeovalis</i> KNOBL. et MAI	C	
Magnoliophyta fam. indet.	<i>Eislebenia naviculaeformis</i> KNOBL. et MAI <i>Normacarpus calyciferus</i> KNOBL. et MAI <i>Rusavia clavata</i> (KNOBL.) KNOBL. et MAI	B–C C	A–B
Stenomaceae	<i>Spirella cerebrata</i> KNOBL. et MAI <i>Spirella ellipsoidea</i> KNOBL. et MAI	B B	
Mastixiaceae	<i>Mastixicarpum cretaceum</i> KNOBL. et MAI <i>Becketia samuelis</i> KNOBL. et MAI <i>Becketia pyriformis</i> KNOBL. et MAI <i>Eomastixia rostrata</i> KNOBL. et MAI	C C C B	
Araliaceae	<i>Acanthopanax gigantocarpus</i> KNOBL. et MAI	D	
Cyrillaceae	<i>Purdiaeopsis campanulatus</i> KNOBL. et MAI	B	
Pentaphylacaceae	<i>Pentaphylax protogaea</i> KNOBL. et MAI <i>Allericarpus pentaphylacoides</i> KNOBL. et MAI	B C	
Theaceae	<i>Visnea minima</i> KNOBL. et MAI	B	
Sabiaceae	<i>Insitiocarpus compressus</i> KNOBL. et MAI	B	
Ericaceae	<i>Leucothoe praecox</i> KNOBL. et MAI	C	
Icacinaceae	<i>Iodes germanica</i> KNOBL. et MAI	B	

and some differences to the Maestrichtian. The features common with the Upper Turonian–Santonian are connected with the coincident species from the Aachen and Klikov Formations, the differences from the floras of the Maestrichtian are connected with the first appearances of numerous living genera during the Maestrichtian.

3.4. Maestrichtian

(Table 4, Plate 4)

Contrary to the Santonian–Campanian in the Maestrichtian many living genera appeared and a great number of extinct genera are missing. 33 genera appear for the first time and 15 of them represent living plants. Among them numerous evergreen elements of the paleotropic geoflora are present:

- **Mastixiaceae**
Becketia, Eomastixia, Mastixicarpum, Mastixiopsis
- **Theaceae**
Palaeoschima, Visnea
- **Ericaceae**
Diplycosiopsis, Leucothoe, Viticocarpum
- **Pentaphylacaceae**
Allericarpus, Pentaphylax
- **Magnoliaceae**
Liriodendroidea, Manglietia
- **Hamamelidaceae**
Disanthus, Rhodoleia

- **Ulmaceae**
Aphananthe, Gironniera
- **Aquifoliaceae**
Ilex
- **Icacinaceae**
Iodes
- **Zingiberaceae**
Spirematospermum

The essential features of the Maestrichtian floras are quite different with respect to the older floras.

The floristic characterization of this interval is given by the rich flora from Walbeck in Germany. Some species from Walbeck are known also from the flysch and the Gosau Formaiton in Czechoslovakia and Austria.

4. Conclusions

Fossil fruits and seeds contribute significantly to our understanding of evolutionary trends in Upper Cretaceous angiosperms. The morphological features of the Upper Cretaceous fruits and seeds demonstrate that the angiosperms had then reached the organization level of those of the present day. The majority of families from the Central-European Cretaceous are of Laurasian origin. The assemblage corresponds to evergreen, subtropical and temperate vegetations in the northern hemisphere today. The origin of some families is questionable since they show affinity with both Laurasian and west Gondwana regions.

Plate 1

Seeds and fruits from the Cenomanian

- Figs. 1,2: ***Spirella pragensis* KNOBL. et MAI (Stenomaceae).**
Praha-Břevnov, Radimova street, Peruc Member, Central Bohemia.
Fig. 1: ×36.
Fig. 2: ×17.
- Figs. 3,4: ***Spirella minutissima* KNOBL. et MAI (Stenomaceae).**
Praha-Břevnov, St. Margaret convent, Peruc Member, Central Bohemia.
Fig. 3: ×60.
Fig. 4: ×80.
- Fig. 5: ***Carpolithus pecinovensis* KNOBL. et MAI (Magnoliophyta fam. indet.).**
Pecínov, Peruc Member, West Bohemia, ×14.
- Fig. 6: ***Viniklaria bohemica* KNOBL. et MAI (Magnoliophyta fam. indet.).**
Vyšehořovice, Peruc Member, Central Bohemia, ×67.
- Fig. 7: ***Brnikia ovalis* KNOBL. et MAI (Magnoliophyt fam. indet.).**
Brník, Peruc Member, Central Bohemia, ×10.
- Fig. 8: ***Klikovispernum pragense* KNOBL. et MAI (Hamamelidaceae).**
Praha-Břevnov, V-26, 23.5 m, Peruc Member, Central Bohemia, ×44.
- Fig. 9: ***Spirella margaritifera* KNOBL. et MAI (Stemonaceae).**
Kněževes PV-2, 25.4 m, Peruc Member, Central Bohemia, ×40.
- Figs. 10-11: ***Carpolithus nyssiformis* KNOBL. et MAI (Magnoliophyta fam. indet.).**
Fig. 10: Pecínov, Peruc Member, West Bohemia, ×10.
Fig. 11: Vyšehořovice, Peruc Member, Central Bohemia, ×16.
- Fig. 12: ***Discosemen faveolatum* KNOBL. et MAI (Magnoliophyta fam. indet.).**
Březinka, Peruc Member, Moravia, ×10.
- Fig. 13: ***Boehmeria cenomanica* KNOBL. et MAI (Urticaceae).**
Zeměchy, 201-53, 19.5 m, Peruc Member, West Bohemia, ×17.
- Fig. 14: ***Klikovispernum verrucosum* KNOBL. et MAI (Hamamelidaceae).**
Slaný SA-15, 49.5 m, Peruc Member, West Bohemia, ×70.
- Fig. 15: ***Carpolithus ovalis* KNOBL. et MAI (Magnoliophyta fam. indet.).**
Pecínov, Peruc Member, West Bohemia, ×42.
- Fig. 16: ***Boehmeria cenomanica* KNOBL. et MAI (Urticaceae).**
Zeměchy, 201-53, 19.5 m, Peruc Member, West Bohemia, ×36.
- Fig. 17: ***Laramisemen hypericoides* KNOBL. et MAI (Magnoliophyta fam. indet.).**
Praha-Břevnov, St. Margaret convent, V-26, 23.5 m, Peruc Member, Central Bohemia, ×40.




Plate 2

Seeds and fruits from the Upper Turonian–Santonian

- Fig. 1: *Memorialis verrucosus* (KNOBL.) KNOBL. et MAI (Urticaceae).
Branišov GB-3, 32.6–32.7 m, Klikov Formation, South Bohemia, ×100.
- Fig. 2: *Laramisemen microcanaliculatum* KNOBL. et MAI (Magnoliophyta fam. indet.).
Branišov, Br-21, 16.7–17.0 m, Klikov Formation, South Bohemia, ×100.
- Fig. 3: *Caryanthus trebecensis* KNOBL. et MAI (?Juglandaceae).
Zahájí, Za-2, 17.6–18.9 m, Klikov Formation, South Bohemia, ×50.
- Fig. 4: *Protovisnea reticulata* KNOBL. et MAI (Theaceae).
Zahájí – Na Blanech, Klikov Formation, South Bohemia, ×54.
- Fig. 5: *Palaeoaldrovanda splendens* KNOBL. et MAI (Droseraceae).
Opatovice Hl-1, 175.9 m, Klikov Formation, South Bohemia, ×22.
- Fig. 6: *Sapindospermum nitidum* KNOBL. et MAI (Sapindaceae).
Třebeč Tj-4a, 76.0–77.0 m, Klikov Formation, South Bohemia, ×25.
- Fig. 7: *Liriodendron papilliformis* KNOBL. et MAI (Magnoliaceae).
Opatovice Hl-1, 66.7 m, Klikov Formation, South Bohemia, ×25.
- Fig. 8: *Protovisnea zahajensis* KNOBL. et MAI (Theaceae).
České Budějovice–Rožnov ČB-3, 62.8–63.1 m, Klikov Formation, South Bohemia, ×70.
- Fig. 9: *Boehmeria ctyrokyi* (KNOBL.) KNOBL. et MAI (Urticaceae).
Branišov Br 21, 23.0–23.5 m, Klikov Formation, South Bohemia, ×50.
- Fig. 10: *Caryanthus deltoides* (KNOBL.) KNOBL. et MAI (?Juglandaceae).
Opatovice, Hl-1, 214.7–215.8 m, Klikov Formation, South Bohemia, ×38.
- Fig. 11: *Klikovispermum malechii* (KNOBL.) KNOBL. et MAI (Hamamelidaceae).
Opatovice, Hl-1, 237.8–238.5 m, Klikov Formation, South Bohemia, ×21.
- Fig. 12: *Vangerovia tricornuta* (KNOBL.) KNOBL. et MAI (Magnoliophyta fam. indet.).
České Budějovice–Vrato, TSv-7, 40.0–41.1 m, Klikov Formation, South Bohemia, ×50.
- Fig. 13: *Calathiocarpus minimus* KNOBL. et MAI (?Juglandaceae).
Záblatí 65/23, 43.2 m, Klikov Formation, South Bohemia, ×70.
- Fig. 14: *Budvaricarpus serialis* KNOBL. et MAI (Magnoliophyta fam. indet.).
Branišov Br 21, 60.5–60.8 m, Klikov Formation, South Bohemia, ×22.




Plate 3

Seeds and fruits from the Santonian–Campanian

- Fig. 1: *Prototinomiscium vangerowii* KNOBL. et MAI (Menispermaceae).
Aachen, Aachen Formation, Germany, $\times 45$.
- Fig. 2: *Operculispermum padragkutense* KNOBL. et MAI (Magnoliophyta fam. indet.).
Jokai-bánya, Ajka Formation, Hungary, $\times 16$.
- Fig. 3: *Discoclethra polysperma* KNOBL. et MAI (Clethraceae).
Aachen, Aachen Formation, Germany, $\times 14$.
- Fig. 4: *Protovisnea cancellata* (VANG.) KNOBL. et MAI (Theaceae).
Aachen, Aachen Formation, Germany, $\times 50$.
- Fig. 5: *Walbeckia aquisgranensis* KNOBL. et MAI (Magnoliophyta fam. indet.).
Aachen, Aachen Formation, Germany, $\times 25$.
- Fig. 6: *Urticoidea cucurbitoides* (VANG.) KNOBL. et MAI (Urticaceae).
Aachen, Aachen Formation, Germany, $\times 40$.
- Fig. 7: *Padragkutia haasii* KNOBL. et MAI (Magnoliaceae).
Jokai bánya, Ajka Formation, Hungary, $\times 30$.
- Fig. 8: *Bicameria orbicularis* (KNOBL.) KNOBL. et MAI (Magnoliophyta fam. indet.).
České Budějovice-Vráto TSv-7, 80.3–80.8 m, Klikov Formation, Czechoslovakia, South Bohemia, $\times 50$.
- Fig. 9: *Eurya crassitesta* KNOBL. (Theaceae).
Kössen, Gosau Formation, Austria, $\times 30$.
- Fig. 10: *Walbeckia ajkoensis* KNOBL. et MAI (Magnoliophyta fam. indet.).
Ajka Aj-167, 103.0 m, Ajka Formation, Hungary, $\times 37$.
- Fig. 11: *Calathiocarpus hollandicus* KNOBL. et MAI (?Juglandaceae).
Staatsmijn Maurits 554, Aachen Formation, The Netherlands, $\times 22$.
- Fig. 12: *Valvaecarpus debeyi* KNOBL. et MAI (Cyrillaceae).
České Budějovice-Budvar, 80.5 m, Klikov Formation, South Bohemia, $\times 15$.
- Fig. 13: *Laramisemen faboides* (VANG.) KNOBL. et MAI (Magnoliophyta fam. indet.).
Epen 44, 3.5–6.0 m, Aachen Formation, The Netherlands, $\times 120$.
- Fig. 14: *Sabia menispermioides* KNOBL. et MAI (Sabiaceae).
České Budějovice-Budvar, 148.5 m, Klikov Formation, Czechoslovakia, $\times 15$.
- Fig. 15: *Sauraia alenae* KNOBL. et MAI (Saurauiaceae).
Branná Sch-2, 116.1 m, Klikov Formation, Czechoslovakia, $\times 38$.
- Fig. 16: *Memorialis rhenana* KNOBL. et MAI (Urticaceae).
Aachen, Aachen Formation, Germany, $\times 40$.





Plate 4

Seeds and fruits from the Maestrichtian

- Fig. 1: *Mastixicarpum cretaceum* KNOBL. et MAI (Mastixiaceae).
Walbeck, Maestrichtian, Germany, $\times 6.5$.
- Fig. 2: *Purdiaeopsis campanulatus* KNOBL. et MAI (Cyrillaceae).
Walbeck, Maestrichtian, Germany, $\times 20$.
- Fig. 3: *Aphananthe cretacea* KNOBL. et MAI (Ulmaceae).
Walbeck, Maestrichtian, Germany, $\times 10$.
- Fig. 4: *Iodes germanica* KNOBL. et MAI (Icacinaceae).
Eisleben, Maestrichtian, Germany, $\times 6.5$.
- Fig. 5: *Leucothoe praecox* KNOBL. et MAI (Ericaceae).
Eisleben, Maestrichtian, Germany, $\times 12$.
- Fig. 6: *Valvaecarpus pterocaryaeformis* (KNOBL.) KNOBL. et MAI (Cyrillaceae).
Horní Bečva, Flysh, Maestrichtian, Czechoslovakia, $\times 35$.
- Fig. 7: *Acanthopanax gigantocarpus* KNOBL. et MAI (Araliaceae).
Walbeck, Maestrichtian, Germany, $\times 6.5$.
- Fig. 8: *Sabia praevalis* KNOBL. et MAI (Sabiaceae).
Walbeck, Maestrichtian, Germany, $\times 6.5$.
- Fig. 9: *Palaeoschima becvensis* (KNOBL.) KNOBL. et MAI (Theaceae).
Wien-Sievering, Sievering Formation, Flysh, Maestrichtian, Austria, $\times 30$.
- Fig. 10: *Rusavia clavata* (KNOBL.) KNOBL. et MAI (Magnoliophyta fam. indet.).
Rusava, Rusava Member, Flysh, Maestrichtian, Czechoslovakia, $\times 30$.
- Fig. 11: *Pentaphylax protogaea* KNOBL. et MAI (Pentaphylacaceae).
Walbeck, Maestrichtian, Germany, $\times 30$.
- Fig. 12: *Beckettie samuelis* KNOBL. et MAI (Mastixiaceae).
Eisleben, Maestrichtian, Germany, $\times 6.5$.
- Fig. 13: *Eislebenia naviculaeformis* KNOBL. et MAI (Magnoliophyta fam. indet.).
Eisleben, Maestrichtian, Germany, $\times 10$.
- Fig. 14: *Rhodoleia cretacea* KNOBL. et MAI (Hamamelidaceae).
Walbeck, Maestrichtian, Germany, $\times 25$.



References

- ALBERS, H.J. & FELDER, W.M.: Litho-, Biostratigraphie und Palökologie der Oberkreide und des Alttertiärs Obersanton-Dan (Paläozän) von Aachen-Südlimburg (Niederlande, Deutschland, Belgien). – Aspekte der Kreide Europas, IUGS Series A, **6**, 47–84, Stuttgart 1979.
- KNOBLOCH, E.: Paläobotanisch-biostratigraphische Charakteristik der Klikov-Schichtenfolge (Oberturon–Santon) in Südböhmen. – Sbor. geol. Věd, Ř. G, **40**, 101–145, Prag 1985.
- KNOBLOCH, E. & MAI, D.H.: Neue Gattungen nach Früchten und Samen aus dem Cenoman bis Maastricht (Kreide) von Mitteleuropa. – Feddes Repert., **95**, 3–41, Berlin 1984.
- KNOBLOCH, E. & MAI, D.H.: Monographie der Früchte und Samen in der Kreide von Mitteleuropa. – Rozpr. Ústř. Úst. geol., **47**, 1–219, Prag 1986.
- MAI, D.H.: Neue Früchte und Samen aus paläozänen Ablagerungen Mitteleuropas. – Feddes Repert., **98** (3–4), 197–229, Berlin 1987.

Manuskript bei der Schriftleitung eingelangt am 20. März 1991.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Jahrbuch der Geologischen Bundesanstalt](#)

Jahr/Year: 1991

Band/Volume: [134](#)

Autor(en)/Author(s): Knobloch Ervin, Mai Dieter Hans

Artikel/Article: [Evolution of Middle and Upper Cretaceous Floras in Central and Western Europe 257-270](#)