

Dinoflagellate Cysts at the Karpatian/Badenian Boundary of Wagna (Styrian Basin, Austria)

ALI SOLIMAN*) & WERNER E. PILLER**)

5 Text-Figures, 2 Plates

Österreichische Karte 1 : 50.000
Blatt 190

Steiermark
Steirisches Becken
Paratethys
Karpatium
Badenium
Dinoflagellaten

Contents

Zusammenfassung	403
Abstract	403
1. Introduction	404
2. Geological Setting	404
3. Material and Methods	404
4. Results and Discussion	404
4.1. Biostratigraphy	406
4.2. Paleoenvironmental Interpretation	408
5. Taxonomic Note	409
Plates 1–2	410
References	414

Dinoflagellaten-Zysten von der Karpatium/Badenium-Grenze bei Wagna (Steirisches Becken, Österreich)

Zusammenfassung

Von der Lokalität Wagna (Karpatium/Badenium, Unter/Mittel-Miozän) werden erstmals Dinoflagellaten-Zysten beschrieben. Die erfassten Vergesellschaftungen beinhalten 38 Taxa und liefern neue biostratigraphische und paläoökologische Informationen. Die Karpatium/Badenium-Grenze wird in allen drei studierten Profilen durch das Erstauftreten von *Operculodinium? borgerholtense* und *Batiacasphaera sphaerica* markiert. Generell ist die Dinozysten-Diversität in allen Proben relativ gering, trotzdem ist ein deutlicher Rückgang direkt unterhalb der Karpatium/Badenium-Grenze auffällig. Das stimmt mit den Foraminiferen-Daten überein und ebenso mit dem Meeresspiegel-Rückgang 3. Ordnung an dieser Grenze. Im Gegensatz zu kalkigen planktischen Foraminiferen scheinen Dinozysten mit ihren organischen Kammerwänden von höheren Nährstoffgehalten, die durch verstärkte vulkanische Aktivität im Karpatium hervorgerufen wurden, nicht beeinflusst worden zu sein.

Abstract

From the locality Wagna (Karpatian/Badenian, Early/Middle Miocene) dinoflagellate cysts are described for the first time. The detected assemblages include 38 taxa and provide new biostratigraphic as well as paleoenvironmental information. The Karpatian/Badenian boundary is clearly marked in all three studied sections by the first occurrences of *Operculodinium? borgerholtense* and *Batiacasphaera sphaerica* with the onset of the Badenian. Generally, dinocyst diversity is relatively low in all studied samples but a distinctive decline just below the Karpatian/Badenian boundary is recorded. This is in accordance with foraminiferal data and coincides with the 3rd order sea-level fall at the Karpatian/Badenian (Early/Middle Miocene) boundary. In contrast to calcareous planktic foraminifers, organic-walled dinocysts seem not to be affected by higher nutrient levels, which may have been induced by increased volcanic activities during the Karpatian.

*) Dr. ALI SOLIMAN, Department of Geology, Faculty of Science, Tanta University, Tanta 31527, Egypt.
Karl-Franzens Universität Graz, Institut für Erdwissenschaften, Bereich Geologie und Paläontologie, Heinrichstraße 26, A 8010 Graz, Austria.
ali.soliman@uni-graz.at

**) o. Univ.-Prof. Dr. WERNER E. PILLER, Karl-Franzens Universität Graz, Institut für Erdwissenschaften, Bereich Geologie und Paläontologie, Heinrichstraße 26, A 8010 Graz, Austria.
werner.piller@uni-graz.at

1. Introduction

The brickyard of Wagna is the best available surface outcrop in the Central Paratethys for the Karpatian/Badenian (Early/Middle Miocene) boundary. This outcrop is well studied in terms of foraminifers and calcareous nannoplankton (RÖGL et al., 2002; SPEZZAFERRI et al., 2002, 2004) as well as for magnetostratigraphy (STINGL & SCHOLGER, 2005) and stable isotope distribution (LATAL & PILLER, 2003).


Dinoflagellate cysts (dinocysts) have not been described in detail so far, neither from Wagna in particular nor from Miocene strata of the Styrian Basin in general. Only two short reports have been published (SOLIMAN & PILLER, 2004, 2005). However, from the entire Central Paratethys Miocene dinocyst assemblages are known from few studies only, e.g., BALTES (1967, 1969), HOCHULI (1978) and JIMÉNEZ-MORENO et al. (in print) for the Early and Middle Miocene and SÜTO-SZENTAI (1994, 1995, 2002, 2003) for the Late Miocene.

This is in clear contrast to the knowledge of time equivalent assemblages from the Atlantic Ocean, northern Europe, and the Mediterranean Sea, from where dinocysts have been extensively documented (e.g., for the North Atlantic by HEAD et al. (1989a,b), MANUM et al. (1989), DE VERTEUIL & NORRIS (1996); for northern Europe by PIASECKI (1980), STRAUSS et al. (2001), DYBKJÆR & RASMUSSEN (2000), DYBKJÆR (2004), MUNSTERMAN & BRINKHUIS (2004); for the Mediterranean by POWELL (1986), BIFFI & MANUM (1988), EL BEIALY (1988), ZEVENBOOM (1995), TORRICELLI & BIFFI (2001), EL BEIALY & ALI (2002), SOLIMAN (2006)).

The present study examines the upper Lower Miocene and lower Middle Miocene dinocysts in the brickyard Wagna (Text-Fig. 1) and provides new information on their stratigraphic distribution at the Karpatian/Badenian boundary as well as on their potential for paleoenvironmental reconstruction.

2. Geological Setting

The outcrop Wagna is located approximately 20 km south of Graz near Leibnitz (Text-Fig. 1). Geologically, it belongs to the Styrian Basin which is part of the Neogene



Text-Fig. 1.
Simplified geologic overview map of the Styrian Basin and location of the studied outcrop.

Pannonian Basin system. The evolution of the Styrian Basin was described in detail by EBNER & SACHSENHOFER (1991, 1995) and SACHSENHOFER (1996).

The sediments in the abandoned brickyard of Wagna represent two units. The lower part belongs to the Kreuzkrumpel Formation and is made up mainly of grey shales with intercalated siltstones, which are informally termed "Steirischer Schlier". The upper part is dominated by marls and sands with interlayered coral and red algal limestones belonging to the Weissenegg Formation. Between these lithostratigraphic units a major unconformity occurs, known as "Styrian unconformity", which is related to the "Styrian Tectonic Phase" of STILLE (1924). In interaction with a global sea level change this tectonic activity caused a major erosional hiatus between the two lithological units, which is coeval with the Karpatian/Badenian (= Early/Middle Miocene) boundary. These tectonic movements, accompanied by volcanic activity, were not a single event but a long term process represented by an angular unconformity already below the Karpatian/Badenian boundary and another unconformity at the boundary (RÖGL et al., 2002; LATAL & PILLER, 2003). The erosional phase is indicated by reworked pebbles in a silty fine sand recognized as "Geröllmergel" (KOLLMANN, 1965; FRIEBE, 1990; LATAL & PILLER, 2003). Biostratigraphically, the major part of the outcrop belongs to calcareous nannoplankton zone NN4, only in the upper part of section 3 NN5 is also represented (RÖGL et al., 2002; SPEZZAFERRI et al., 2002, 2004; LATAL & PILLER, 2003).

3. Materials and Methods

The outcrop was logged in three sections which overlap laterally. In all 3 sections the Karpatian/Badenian boundary is represented (Text-Figs. 2–4). The sections were already described in detail by LATAL & PILLER (2003) and SPEZZAFERRI et al. (2004). In the current study a total of 37 samples were examined in detail for organic-walled dinoflagellate cysts and acritarchs. These samples and the sample numbers are identical with those studied by SPEZZAFERRI et al. (2002, 2004), RÖGL et al. (2002), and LATAL & PILLER (2003).

For studying dinocysts, 20 to 30 grams of dry rock sample were processed following the standard palynological procedure (WOOD et al., 1996) including 30 % hydrochloric acid (HCl), 40 % hydrofluoric acid (HF), ultrasonic treatment for 10–15 seconds, and sieving over a 20 µm nylon mesh. Neither oxidation nor alkali treatment were applied to the residues. Two microscope slides were made from each sample using glycerine jelly as a mounting medium. Dinocysts were counted and photographed from both slides using a Zeiss Axioplan 2 microscope fitted with a Leica DFC 320 digital camera. Dinocyst nomenclature generally follows that of FENSOME & WILLIAMS (2004).

The rock samples, palynological residues and slides are stored in the collection of the Institute of Earth Sciences, Geology and Paleontology, University of Graz, Austria.


4. Results and Discussion

Dinoflagellate cysts are recorded from all studied samples, however, preservation varies from poor to good and diversity is generally low with a total number of 38 in situ taxa (in alphabetical order):

1. *Apteodinium tectatum* PIASECKI, 1980.
2. *Batiacasphaera sphaerica* STOVER, 1977.
3. *Cleistosphaeridium ancyreum* (COOKSON & EISENACK) EATON et al., 2001.
4. *Cleistosphaeridium placacanthum* (DEFLANDRE & COOKSON) EATON et al., 2001.

5. *Cousteaudinium aubryae* DE VERTEUIL & NORRIS, 1996.
6. *Cribroperidinium giuseppei* (MORGENROTH) HELENES, 1984.
7. *Cribroperidinium tenuitubulatum* (GREAT) HELENES, 1984.
8. *Dapsilidinium pastielsii* (DAVEY & WILLIAMS) BUJAK et al., 1980.
9. *Distalidinium pseudocolligerum* (STOVER) BUJAK et al., 1980.
10. *Distatodinium paradoxum* (BROSIOUS) EATON 1976.
11. *Hystrichokolpoma denticulatum* MATSUOKA, 1974.
12. *Hystrichokolpoma rigaudiae* DEFLANDRE & COOKSON, 1955.
13. *Hystrichosphaeropsis obscura* HABIB, 1972.
14. *Lejeunecysta communis* BIFFI & GRIGNANI, 1983.
15. *Lejeunecysta convexa* MATSUOKA & BUJAK, 1988.
16. *Lejeunecysta diversiforma* (BRADFORD) ARTZNER & DÖRHÖFER, 1978.
17. *Lejeunecysta fallax* (MORGENROTH) ARTZNER & DÖRHÖFER, 1978 emend. BIFFI & GRIGNANI, 1983.
18. *Lejeunecysta globosa* BIFFI & GRIGNANI, 1983.
19. *Lingulodinium machaerophorum* (DEFLANDRE & COOKSON) WALL, 1967.
20. *Melitasphaeridium choanophorum* (DEFLANDRE & COOKSON) HARLAND & HILL, 1979.
21. *Nematosphaeropsis labyrinthus* (OSTENFELD) REID, 1974.
22. *Operculodinium ? borgerholtense* LOUWYE, 2001, emend. herein.
23. *Palaeocystodinium powellii* STRAUSS et al., 2001.
24. *Pentadinium laticinctum* GERLACH, 1961 emend. BENEDEK et al., 1982.
25. *Polysphaeridium zoharyi* (ROSSIGNOL) BUJAK, DOWNIE, EATON & WILLIAMS, 1980.
26. *Reticulatosphaera actinocoronata* (BENEDEK) BENEDEK & MATSUOKA, 1986.
27. *Selenopemphix bothrion* HARLAND & PUDSEY, 2002.
28. *Selenopemphix brevispinosa* HEAD, NORRIS & MUDIE, 1989.
29. *Selenopemphix nephroides* BENEDEK emend. BUJAK in BUJAK et al., 1980.
30. *Selenopemphix quanta* (BRADFORD) MATSUOKA, 1985.
31. *Spiniferites pseudofurcatus* (KLUMPP) SARJEANT, 1970 emend. SARJEANT, 1981.
32. *Spiniferites solidago* DE VERTEUIL & NORRIS, 1996.
33. *Sumatradinium druggii* LENTIN, FENSOME & WILLIAMS, 1994.
34. *Sumatradinium soucouyantiae* DE VERTEUIL & NORRIS, 1992.
35. *Trinovantedinium* sp. cf. *T. applanatum* (BRADFORD) BUJAK & DAVIES, 1983.
36. *Tuberculodinium vancampoe* (ROSSIGNOL) WALL, 1967.
37. *Xandarodinium* sp.
38. Dinocyst VI of MANUM (1976), sensu SCHIÖLER, 2005.

The distribution of the most abundant taxa is shown in Text-Figures 2–4. Dinocyst microphotographs are shown in Plates 1 and 2. Pollen and spores were observed in all three sections but not studied.



Text-Fig. 3.

Stratigraphic distribution of selected dinoflagellate cysts and associated palynomorphs in section 2.

For explanation of lithology compare LATAL & PILLER (2003) and SPEZZAFERRI et al. (2004).

4.1. Biostratigraphy


The studied sections are well dated by means of planktic and benthic foraminifers and calcareous nannoplankton (e.g., RÖGL et al., 2002; SPEZZAFERRI et al., 2002, 2004; CORIC, 2002; Text-Fig. 5). Dinoflagellate cyst distribution provides additional biostratigraphic information.

Karpatian

The Karpatian part of the studied sections is characterized by the common occurrence of *Operculodinium centrocarpum*, *Lingulodinium machaerophorum*, *Reticulatosphaera actinocoronata*, and *Spiniferites* spp. undiff. These taxa are present in almost all samples. Sporadic occurrences are detected for *Melitasphaeridium choanophorum*, *Spiniferites pseudofurcatus*, *Cribroperidinium tenuitubulatum*, *Hystrichokolpoma rigaudiae*, *Dapsilodinium* spp., and *Pentadinium laticinctum*. In addition to the above mentioned species a group of characteristic Early Miocene species, as *Sumatradinium soucouyantiae*, *Cousteaudinium aubryae*, *Hystrichosphaeropsis obscura*, *Cleistosphaeridium placacanthum*, *Tuberculodinium vancampoe*, *Palaeocystodinium powellii*, and *Hystrichokolpoma denticulatum*, was also recognized. The acritarch species *Cyclopsiella lusatica* occurs in some samples (Text-Figs. 2–4).

Outside the Central Paratethys *Hystrichosphaeropsis obscura* (Pl. 1, Figs. 14, 18) indicates Burdigalian age according to many authors (e.g., HAQ et al., 1987; MANUM et al., 1989; POWELL, 1992; STOVER et al., 1996; HARDENBOL et al., 1998; DYBKJÆR, 2004). *S. soucouyantiae* (Pl. 1, Fig. 16), which occurs in all studied sections of Wagna (section 1: sample 10; section 2: sample 1; section 3: sample 6), is restricted to the Early Miocene (DE VERTEUIL & NORRIS, 1996; HARDENBOL et al., 1998; WILLIAMS et al., 2004; GRADSTEIN et al., 2004).

Cousteaudinium aubryae (Pl. 2, Figs. 20, 21) is persistently recorded from section 2 and sporadically from section 3; no record comes from section 1. In Cortemilia section (Italy) the lowest occurrence (LO) of *C. aubryae* (described as *Thalassiphora gonoperforata*) defines the Tgo Interval Zone (uppermost Burdigalian) of ZEVENBOOM (1995). In the eastern United States, however, the LO of this species is dated as mid-Aquitania (base of dinocyst zone DN2) and is correlated with the lower part of nannoplankton zone NN2 (DE VERTEUIL & NORRIS, 1996). In the southern North Sea Basin its LO is also detected at the base of dinocyst zone M2 and dated as mid-Aquitania (MUNSTERMAN & BRINKHUIS, 2004). WILLIAMS et al. (2004) argue for a heterochronous occurrence of this species with a mid-Aquitania LO



Text-Fig. 4.
Stratigraphic distribution of selected dinoflagellate cysts and associated palynomorphs in section 3.
For explanation of lithology compare LATAL & PILLER (2003) and SPEZZAFERRI et al. (2004).

for mid latitudes in the northern hemisphere and an upper Burdigalian LO for equatorial regions.


The absence of some of the early to middle Early Miocene marker taxa (e.g., *Exochosphaeridium insigne*, *Cordosphaeridium cantharellus*) from the studied sections is in accordance with the Karpatian age (late Burdigalian). In the eastern Alpine Foreland Basin of Austria (section Strass-Eberschwang) the highest occurrence (HO) of these species defines the top of the *E. insigne* Assemblage Biozone (Ein), which is middle Ottnangian in age and correlated to calcareous nannoplankton Zone NN3 (JIMÉNEZ-MORENO et al., in press).

Badenian

The Badenian sediments are represented by 1 sample in section 1, 6 samples in section 2 and 6 samples in section 3. The dinocyst assemblage of the Karpatian part of the

sections is generally also recorded in the Lower Badenian part. In addition, *Operculodinium ? borgerholtense* (Pl. 1, Figs. 6–12) and *Batiacasphaera sphaerica* (Pl. 1, Figs. 1–5) have their LO immediately above the Karpatian/Badenian boundary. The uppermost occurrences of the two lower Middle Miocene marker species, *Distatodinium paradoxum* (Pl. 1, Figs. 13, 17) and *C. aubryae*, are recorded in sections 2 and 3 (Text-Figs. 3, 4).

The HO of *D. paradoxum* and *C. aubryae* is reported from the top of dinocyst zone DN4 in the eastern USA correlated with lowermost NN5 (lower Langhian) (DE VERTEUIL & NORRIS, 1996). The highest occurrence of both species was documented in the lowermost Middle Miocene at the top of Cpl dinocyst zone of well Nieder Ochtenhausen (STRAUSS et al., 2001) and at the top of dinocyst zone DN4 of northern Germany (KÖTHE, 2003). In the southern North Sea area the HO of *C. aubryae* defined the dinocyst zone SNSM6 (MUNSTERMAN & BRINKHUIS, 2004).



Text-Fig. 5.
Summarized stratigraphic ranges of selected dinoflagellate cysts and acritarchs in the Wagna outcrop.

The most characteristic bioevent of the Badenian in the studied samples is the first occurrence of *O. ? borgerholtense*. This species is not abundant but persistently recorded. In the Retznei outcrop (Badenian), few kilometers south of Wagna, this species is very abundant and shows high intraspecific variability in its morphologic features (personal observation). The exact stratigraphic range of this species is unknown yet, but it is well documented from the Middle Miocene of the North Sea, Gulf of Suez, and Central Paratethys basins (LOUWYE, 2001; SOLIMAN, 2006; JIMÉNEZ-MORENO et al., in print; personal observations).

The LO of *B. sphaerica* is also recorded in the Badenian of sections 2 and 3. This species was first described from the Lower Miocene *Globorotalia kugleri* Zone of the Blake Plateau, offshore North Carolina (STOVER, 1977). It is, however, also recorded from the Aquitanian (Ebu Zone) of the Santa

Croce di Arcevia Section in Italy (ZEVENBOOM, 1995), the Burdigalian stratotype area, Bordeaux, France (LONDEIX & JAN DU CHÊNE, 1998), the uppermost Upper Burdigalian – Lower Langhian of the Zonderschot Sands, northern Belgium (LOUWYE, 2000), and the middle Ottnangian (Burdigalian) of the Strass-Eberschwang section, Austria (JIMÉNEZ-MORENO et al., in print).

Thus, the LO of *B. sphaerica* in Wagna is younger than all other known records.

Karpatian/Badenian Boundary

In the Styrian Basin the Karpatian/Badenian boundary is represented by the “Styrian unconformity” (Text-Fig. 5), which is related to the “Styrian Tectonic Phase” (STILLE, 1924; LATAL & PILLER, 2003). In the brickyard of Wagna SPEZZAFERRI et al. (2002, 2004) attributed the “Steirischer Schlier”, which represents the sediments below the unconformity, to the middle Karpatian based on the presence of *Uvigerina graciliformis*, *Pappina primiformis*, a typical small sized planktic foraminiferal assemblage, and the rare occurrence of *Globigerinoides bisphericus* (planktic foraminiferal zone M4b). The portion above the unconformity is assigned to the Lower Badenian based on foraminifers (*Globigerinoides trilobus*, *Praeorbulina glomerata circularis*) and correlated with planktic foraminiferal zone M5a (Text-Fig. 5). Sedimentological features and an incomplete evolutionary lineage from *Globigerinoides bisphericus* to *Praeorbulina* species, however, clearly indicate a gap in sedimentation. In terms of calcareous nannoplankton all samples of the 3 sections studied belong to zone NN4 based on the co-occurrence of *Sphenolithus heteromorphus* and *Helicosphaera ampliapertura*. The latter increases in abundance just below the unconformity (RÖGL et al., 2002).

With regard to the dinocyst assemblages the boundary is marked by the lowest occurrences (LO) of *O. ? borgerholtense* and *B. sphaerica*. Both species are not recorded below the unconformity but occur persistently above, especially *O. ? borgerholtense* (Text-Fig. 5).

4.2. Paleoenvironmental Interpretation

Generally, the recorded dinoflagellate cyst assemblages (Text-Figs. 2–4) reflect a neritic environment for the studied area. The sporadic occurrences of *N. labyrinthus* in the “Steirischer Schlier” (Karpatian) of sections 1 and 2 indicate that these beds were deposited in a deep water environment. DALE (1996) pointed out that the presence of even a few specimens of these cysts is a reliable indication of deep waters. In the Badenian parts of the sections *N. labyrinthus* is completely missing. For the same part of section 1 SPEZZAFERRI et al. (2004) suggested a paleodepth from 250 m to 300 m based on benthic agglutinated and calcareous foraminifers.

A significant number of thermophile species, as *T. vancampoe*, *L. machaerophorum*, *S. nephroides*, *P. zoharyi*, and *M. choanophorum*, indicates subtropical conditions for all studied samples (e.g., EDWARDS & ANDRLE, 1992; HEAD & WESTPHAL, 1999; MARRET & ZONNEVELD, 2003). Temperate to cool-water species, as *Impagidinium pallidum* BUJAK, 1984 and *Spiniferites elongatus* REID, 1974, were not recognized. No distinct change between the Karpatian and Badenian parts of the sections was detected. For the Karpatian this contrasts with the results of SPEZZAFERRI et al. (2002), in which the foraminiferal ratios point to cool climatic conditions. This interpretation is, however, already in contrast to the coeval sediments of the Molasse Basin (Laa an der Thaya, Göllersdorf) where warm climatic conditions are indicated by foraminiferal assemblages. Since the two basins (Molasse Basin and Styrian Basin) are only a few hundred kilometers apart one plausible explanation could be the increased volcanic activity in the Styrian Basin that caused

higher nutrient concentrations. The latter may have provoked a small and poorly diversified planktic foraminiferal assemblage veiling the original climate signal (SPEZZAFERRI et al., 2004). The organic walled dinoflagellate cyst assemblages seem to be less affected by this volcanically induced nutrification. The only hint to a nutrient enrichment in the Karpatian part of the sections is the higher abundance and diversification of heterotrophic protoperidinioid dinoflagellate cysts, as *Lejeunecysta* spp., *Selenopemphix* spp., and *Trinovantedinium* spp. (e.g., WALL et al., 1977; BUJAK, 1984; POWELL et al., 1990; LEWIS et al., 1990; SOLIMAN, 2006).

Generally, dinocyst diversity is relatively low in all the studied samples, however, a decline is noted just below the Karpatian/Badenian boundary (Text-Figs. 3, 4). This decline clearly coincides with a drastic change in foraminiferal assemblages with an increasing number of shallow-water taxa, as *Ammonia*, *Nonion*, *Elphidiella*, *Elphidium*, and small globigerinids as referred by RÖGL et al. (2002) and SPEZZAFERRI et al. (2002, 2004). On the base of this foraminiferal change a decreasing water depth from ca. 240 m to 50 m was calculated (SPEZZAFERRI et al., 2004). This shallowing of the depositional environment may be also the reason for the decreasing dinocyst diversity and is in accordance with the sea-level low stand at the Karpatian/Badenian boundary (FRIEBE, 1993; EBNER & SACHSENHOFER, 1995), which can be correlated with the TB 2.3 sequence boundary of HAQ et al. (1988) and Lan-1 by VAKARCS et al. (1998) (LATAL & PILLER, 2003).

5. Taxonomic Note

Operculodinium ? borgerholtense LOUWYE, 2001

Plate 1, Figs. 6–12

2001 *Operculodinium ? borgerholtense* LOUWYE, p. 126–127, Figs. 4.1–4.12.

Discussion: In their emendation of the genus *Operculodinium* MATSUOKA et al. (1997) did not state the presence of granules and/or small rods as a surface ornamentation as well as spinules and/or small granules on the process shaft. *O. ? borgerholtense* is distinguished from “typical” species of this genus by its prismatic luxuria (low granules and/or small rods) and in that shafts of the processes have spinules or granules. However, the latter can also be smooth in some specimens. The antapical plate with fused low and high ornamentation is clearly seen in many specimens but may depend on the specimen’s orientation (Pl. 1, Fig. 12). These characteristic morphologic features distinguish this species from other *Operculodinium* species and require its emendation.

Previous records: Middle Miocene (Antwerp Sands, Berchem Formation) of northern Belgium (LOUWYE, 2001); uppermost Lower Miocene–Middle Miocene, Gulf of Suez, Egypt (SOLIMAN, 2006); Middle Miocene (Badenian-Sarmatian) of the Central Paratethys (JIMÉNEZ-MORENO et al., in print); Middle Miocene of the Styrian Basin, Retznei section (personal observation); Middle Miocene of the North Sea and North Atlantic basins (personal communications by M.J. HEAD and S. LOUWYE).

Acknowledgements

This study was financially supported by the Commission for the Palaeontological and Stratigraphical Research of Austria (Austrian Academy of Sciences) and project P-14366-Bio of the Austrian Science Fund. Field work was assisted by A. KROH (Vienna), C. LATAL (Graz), and C. ERHART (Graz).

Plate 1

Dinoflagellate cysts from the Wagna outcrop.

- Figs. 1, 5: *Batiacasphaera sphaerica*.
Sample W01-22, E. F. Ref. B8, low and high focus of indet. orientation.
- Figs. 2–4: *Batiacasphaera sphaerica*.
Sample W01-19, slide A, E. F. Ref. X15-1, high focus showing surface ornamentation, mid-focus and low focus showing apical archeopyle.
- Figs. 6–8: *Operculodinium ? borgerholtense*.
Sample W01-19, slide A, E. F. Ref. M56, different foci.
- Figs. 9–10: *Operculodinium ? borgerholtense*.
Sample W01-19, slide A, E. F. Ref. K17, dorsal view showing 2P archeopyle.
- Figs. 11–12: *Operculodinium ? borgerholtense*.
Sample W02-18, slide B, E. F. Ref. A28-3, oblique antapical view showing antapical plate.
- Figs. 13, 17: *Distatodinium paradoxum*.
Sample W01-17, slide A, E. F. Ref. R22, note characteristic distal ends of processes.
- Figs. 14, 18: *Hystriosphæropsis obscura*.
Sample W01-17, slide A, E. F. Ref. Y25-2, ventral view of ventral surface.
- Fig. 15: *Sumatradinium druggii*.
Sample W01-16, slide A, E. F. Ref. U34-4, note annular thickenings at the processes shafts.
- Fig. 16: *Sumatradinium soucouyantiae*.
Sample Wa2-10, E. F. Ref. X54-1, specimen in dorsal view.
- Fig. 19: *Palaeocystodinium powellii*.
Sample W02-6, slide A, E. F. Ref. H46.
- Fig. 20: *Palaeocystodinium powellii*.
Sample W02-6, slide A, E. F. Ref. F66.
- Figs. 21–22: Dinocyst VI of MANUM (1976), sensu SCHIÖLER, 2005.
Sample W01-22, slide A, E. F. Ref. P6, different foci of uncertain position.

Scale bar 20 μm .

E. F. Ref. = England Finder Reference.




Plate 2

Dinoflagellate cysts from the Wagna outcrop.

- Figs. 1–2: *Cribroperidinium tenuitubulatum*.
Sample W02-11, slide A, E. F. Ref. N45-1, mid and high foci of a lateral view showing the apical horn and surface ornamentation.
- Figs. 3, 7: *Cleistosphaeridium placacanthum*.
Sample W02-10, slide A, E. F. Ref. N63, indet. orientation, different foci showing the developed base sutures and distal ends of processes.
- Figs. 4, 8: *Lingulodinium machaerophorum*.
Sample W01-13, slide A, E. F. Ref. T23, low and high foci, note granular surface ornamentation.
- Figs. 5, 6: *Melitasphaeridium choanophorum*.
Sample W02-11, slide A, E. F. Ref. F53-3, different foci of uncertain orientation.
- Figs. 9–10: *Pentadinium laticinctum*.
Sample W02-11, slide A, E. F. Ref. H65-3, different foci.
- Figs. 11: *Reticulosphaera actinocoronata*.
Sample W02-14, slide A, E. F. Ref. V48-1, mid-focus.
- Figs. 12–13: *Cyclopsiella lusatica*.
Sample W01-19, slide A, E. F. Ref. S18-3, low and high focus, note the pylome is near the periphery.
- Figs. 14, 19: *Pyxidinospis* sp.
Sample W01-22, slide A, E. F. Ref. Y28-4.
- Figs. 15–16: *Nematosphaeropsis labyrinthus*.
Sample W01-13, slide A, E. F. Ref. K38-3, different foci.
- Figs. 17–18: *Spiniferites solidago*.
Sample W02-11, slide A, E. F. Ref. T59, dorsal view, different foci, note mid-shaft vacuoles and apical boss.
- Figs. 20–21: *Cousteaudinium aubryae*.
Sample W02-9, slide A, E. F. Ref. F34, dorsal view of dorsal surface showing archeopyle sutures.
- Fig. 22: *Lejeunecysta diversiforma*.
Sample Wa2-23, E. F. Ref. J50, mid-focus.
- Figs. 23–24: *Selenopemphix bothrion*.
Sample Wa3-23, E. F. Ref. L31, low and high foci.
- Fig. 25: *Lejeunecysta fallax*.
Sample W01-20, slide A, E. F. Ref. D8-3.
- Fig. 26: *Trinovantedinium* sp. cf. *T. applanatum*.
Sample W01-15, slide A, E. F. Ref. S36-4, lateral view.

Scale bar 20 μ m.

E. F. Ref. = England Finder Reference.



References

- BALTES, N. (1967): Microflora from Miocene salt-bearing formation of the pre-Carpathian depression (Rumania). – *Rev. Palaeobot. Palynol.*, **2**, 183–194.
- BALTES, N. (1969): Distribution stratigraphique des dinoflagellés et des acritarches tertiaires en Roumanie. – In: BRÖNNIMANN, P. & RENZ, H.H. (Eds.): 1st International Conference on Planktonic Microfossils, Geneva, 1967, Proceedings, vol. 1, 26–45, Leiden (E.J. Brill).
- BIFFI, U. & MANUM, S.B. (1988): Late Eocene–Early Miocene dinoflagellate cyst stratigraphy from the Marche Region (Central Italy). – *Bolletino della Società Paleontologica Italiana*, **27**(2), 163–212.
- BUJAK, J.P. (1984): Cenozoic dinoflagellate cysts and acritarchs from the Bering Sea and northern North Pacific, D.S.D.P. Leg 19. – *Micropaleontology*, **30**, 180–212.
- CORIC, S. (2002): Calcareous nannoplankton biostratigraphy from the Early/Middle Miocene boundary (Karpatian-Badenian boundary) in the Styrian and Vienna Basin. – 7. Österreichischer Sedimentologenworkshop, 9. 11. 2002, Seewalchen am Attersee, Programm und Kurzfassungen, p. 3.
- DALE, B. (1996): Dinoflagellate cyst ecology: modeling and geological applications. – In: JANSONIUS, J. & MCGREGOR, D.C. (Eds.): *Palynology: Principles and Applications*. – Vol. 3, 1249–1275, American Association of Stratigraphic Palynologists Foundation.
- DE VERTEUIL, L. & NORRIS, G. (1996): Part 1. Dinoflagellate cyst zonation and allostratigraphy of the Chesapeake Group. – In: DE VERTEUIL, L. & NORRIS, G. (Eds.): *Miocene dinoflagellate stratigraphy and systematics of Maryland and Virginia*. – *Micropaleontology*, **42** (supplement), 1–172.
- DYBKJÆR, K. (2004): Dinocyst stratigraphy and palynofacies studies used for refining a sequence stratigraphic model – uppermost Oligocene to lower Miocene, Jylland, Denmark. – *Rev. Palaeobot. Palynol.*, **131**, 201–249.
- DYBKJÆR, K. & RASMUSSEN, E.S. (2000): Palynological dating of the Oligocene–Miocene successions in the Lille Bælt area, Denmark. – *Bulletin of the Geological Society of Denmark*, **47**, 87–103.
- EBNER, F. & SACHSENHOFER, R.F. (1991): Die Entwicklungsgeschichte des Steirischen Tertiärbeckens. – *Mitt. Abt. Geologie Paläontologie Landesmuseum Joanneum*, **49**, 1–96.
- EBNER, F. & SACHSENHOFER, R.F. (1995): Palaeogeography, subsidence and thermal history of the Neogene Styrian Basin (Pannonian basin system, Austria). – *Tectonophysics*, **242**, 133–150.
- EDWARDS, L.E. & ANDRLE, A.S. (1992): Distribution of selected dinoflagellate cysts in modern marine sediments. – In: HEAD, M.J. & WRENN, L.H. (Eds.): *Neogene and Quaternary Dinoflagellate Cysts and Acritarchs*, American Association of Stratigraphic Palynologists Foundation, 259–288.
- EL BEIALY, S.Y. (1988): Palynostratigraphy of late Tertiary sediments in Kafr El-Dawar well No. 1, Nile Delta, Egypt. – *Revue de Micropaléontologie*, **30**(4), 249–260.
- EL BEIALY, S.Y. & ALI, A.S. (2002): Dinoflagellates from the Miocene Rudeis and Kareem formations borehole GS-78-1, Gulf of Suez, Egypt. – *Journal of African Earth Sciences*, **35**, 235–245.
- FENSOME, R.A. & WILLIAMS, G.L. (2004): The Lentin and Williams index of fossil dinoflagellates: 2004 Edition. – *American Association of Stratigraphic Palynologists, Contributions Series*, **42**, 909 pp.
- FRIEBE, J.G. (1990): Lithostratigraphische Neugliederung und Sedimentologie der Ablagerungen des Badenians (Miozän) um die Mittelsteirische Schwelle (Steirisches Becken, Österreich). – *Jb. Geol. B.-A.*, **133**/2, 223–257.
- FRIEBE, J.G. (1993): Sequence stratigraphy in a mixed carbonate-siliciclastic depositional system (Middle Miocene; Styrian Basin, Austria). – *Geol. Rdsch.*, **82**, 281–294.
- GRADSTEIN, F.M., OGG, J.G., SMITH, A.G., BLEEKER, W. & LOURENS, L.J. (2004): A new geologic time scale with special reference to Precambrian and Neogene. – *Episodes*, **27**, 83–100.
- HAQ, B.U., HARDENBOL, J. & VAIL, P.R. (1987): Chronology of fluctuating sea levels since the Triassic. – *Science*, **235**, 1156–1167.
- HAQ, B.U., HARDENBOL, J. & VAIL, P.R. (1988): Mesozoic and Cenozoic chronostratigraphy and cycles of sea-level changes. – In: WILCUS, C.K., HASTINGS, B.S., KENDALL, C.G.S.C., POSAMANTIER, H.W., ROSS, C.A. & VAN WAGONER, J.C. (Eds.): *Sea-level changes: an integrated approach*, Society of Economic Paleontologists and Mineralogists Special Publication, **42**, 71–108.
- HARDENBOL, J., THIERRY, J., FARLEY, M. B., JACQUIN, T., DE GRACIANSKY, P.-C. & VAIL, P. (1998): Cenozoic Biochronostratigraphy. – In: DE GRACIANSKY, P.-C., HARDENBOL, J., THIERRY, J., FARLEY, M. B. & VAIL, P. (Eds.): *Mesozoic and Cenozoic Sequence Stratigraphy of European Basins*, SEPM, Special Publication, **60**, Chart 3.
- HEAD, M.J., NORRIS, G. & MUDIE, P.J. (1989a): Palynology and dinocyst stratigraphy of the Upper Miocene and lowermost Pliocene, ODP Leg 105, site 646, Labrador Sea. – In: SRIVASTAVA, S.P., ARTHUR, M. & CLEMENT, B. (Eds.): *Proceedings of the Ocean Drilling Program, Scientific Results*, **105**, 423–451.
- HEAD, M.J., NORRIS, G. & MUDIE, P.J. (1989b): New species of dinocysts and a new species of acritarch from the Upper Miocene and lowermost Pliocene, ODP Leg 105, Site 646, Labrador Sea. – In: SRIVASTAVA, S.P., ARTHUR, M. & CLEMENT, B. (Eds.): *Proceedings of the Ocean Drilling Program, Scientific Results*, **105**, 453–466.
- HEAD, M.J. & WESTPHAL, H. (1999): Palynology and paleoenvironments of a Pliocene carbonate platform: the Clino Core, Bahamas. – *Journal of Paleontology*, **73**, 1–25.
- HOCHULI, P.A. (1978): Palynologische Untersuchungen im Oligozän und Untermiozän der Zentralen und Westlichen Paratethys. – *Beiträge zur Paläontologie von Österreich*, **4**, 1–132.
- JIMÉNEZ-MORENO, G., HEAD, M.J. & HARZHAUSER, M. (in print): Early and Middle Miocene dinoflagellate cyst stratigraphy of the central Paratethys, central Europe. – *Journal of Micropaleontology*.
- KOLLMANN, K. (1965): Jungtertiär im Steirischen Becken. – *Mitt. Geol. Ges. Wien*, **57**/2, 479–632.
- KÖTHE, A. (2003): Dinozysten-Zonierung im Tertiär Norddeutschlands. – *Revue de Paléobiologie*, **22**(2), 895–923.
- LATAL, C. & PILLER, W.E. (2003): Stable Isotope Signatures at the Karpatian/Badenian Boundary in the Styrian Basin. – In: BRZOBHATY, R., CÍCHA, I., KOVAC, M. & RÖGL, F. (Eds.): *The Karpatian – a Lower Miocene Stage of the Central Paratethys*, 37–48, Brno (Masaryk University Brno).
- LEWIS, J., DODGE, J.D. & POWELL, A.J. (1990): Quaternary dinoflagellate cysts from the upwelling system offshore Peru, hole 686B, ODP Leg 112. – *Proceedings of the Ocean Drilling Program, Scientific Results*, **112**, 323–327.
- LONDEIX, L. & JAN DU CHÊNE, R. (1998): Burdigalian dinocyst stratigraphy of the stratotypic area (Bordeaux, France). – *Geobios*, **30**, 283–294.
- LOUWYE, S. (2000): Dinoflagellate cysts and acritarchs from the Miocene Zonderschot Sands (northern Belgium): stratigraphic significance and correlation with contiguous areas. – *Geologica Belgica*, **3**, 55–65.
- LOUWYE, S. (2001): New dinoflagellate cysts from the Berchem Formation, Miocene, northern Belgium (southern North Sea Basin). – *Geobios*, **34**, 121–130.
- MANUM, S.B., BOULTER, M.C., GUNNARSDOTTIR, H., RANGNES, K. & SCHOLZE, A. (1989): Eocene to Miocene palynology of the Norwegian Sea (ODP Leg 104). – In: ELDHOLM, O., THIEDE, J., TAYLOR, E. et al. (Eds.): *Proceedings of the Ocean Drilling Program, Scientific Results*, **104**, 611–662.
- MARRET, F. & ZONNEVELD, K.A.F. (2003): Atlas of modern organic-walled dinoflagellate cyst distribution. – *Rev. Palaeobot. Palynol.*, **125**, 1–200.
- MATSUOKA, K., MCMINN, A. & WRENN, J.H. (1997): Restudy of the holotype of *Operculodinium centrocarpum* (Deflandre and Cookson) Wall (Dinophyceae) from the Miocene of Australia, and the taxonomy of related species. – *Palynology*, **21**, 19–33.
- MUNSTERMANN, D.K. & BRINKHUIS, H. (2004): A southern North Sea Miocene dinoflagellate cyst zonation. – *Geologie en Mijnbouw*, **83** (4), 267–285.
- PIASECKI, S. (1980): Dinoflagellate cyst stratigraphy of the Miocene Hodde and Gram Formations, Denmark. – *Geological Society of Denmark, Bulletin*, **29**, 53–76.
- POWELL, A.J. (1986): A dinoflagellate cyst biozonation for the Late Oligocene to Middle Miocene succession of the Langhe region, Northwest Italy. – In: WRENN, J.H., DUFFIELD, S.L. & STEIN, J.A. (Eds.): *Papers from the first symposium on Neogene dinoflagellate cyst biostratigraphy*. – *American Association of Stratigraphic Palynologists, Contributions Series*, **17**, 105–127.
- POWELL, A.J. (1992): Dinoflagellate cysts of the tertiary system. – In: POWELL, A.J. (Ed.): *A stratigraphic index of dinoflagellate cysts*,

- British Micropalaeontological Society Publication Series, London (Chapman and Hall), 155–251.
- POWELL, A.J., DODGE, J.D. & LEWIS, J. (1990): Late Neogene to Pleistocene palynological facies of the Peruvian continental margin upwelling, Leg 112. – Proceedings Ocean Drilling Program, Scientific Results, **112**, 297–321.
- RÖGL, F., SPEZZAFERRI, S. & CORIC, S. (2002): Micropaleontology and biostratigraphy of the Karpatian–Badenian transition (Early–Middle Miocene boundary) in Austria (Central Paratethys). – Courier Forschungsinstitut Senckenberg, **237**, 47–67.
- SACHSENHOFER, R.F. (1996): The Neogene Styrian Basin: An overview. – Mitt. Ges. Geol. Bergbaustud. Österr., **41**, 19–32.
- SOLIMAN, A. (2006): Lower and Middle Miocene dinoflagellate cysts, Gulf of Suez, Egypt. – Unpublished Ph.D. Thesis, Graz University, 327 p.
- SOLIMAN, A. & PILLER, W.E. (2004): Miocene Dinoflagellate cysts of the Styrian Basin, Austria. – Ber. Inst. Erdwiss., K.-F.-Universität. Graz, **9**, 379.
- SOLIMAN, A. & PILLER, W.E. (2005): Karpatian–Badenian (Middle Miocene) dinoflagellate cysts of the Styrian Basin, Austria. – Ber. Inst. Erdwiss., K.-F.-Universität Graz, **10**, 129.
- SPEZZAFERRI, S., CORIC, S., HOHENEGGER, J. & RÖGL, F. (2002): Basin-scale paleobiogeography and paleoecology: an example from Karpatian (Latest Burdigalian) benthic and planktonic foraminifera and calcareous nannoplankton from the Central Paratethys. – Geobios, Mémoire spécial, **24**, 241–256.
- SPEZZAFERRI, S., RÖGL, F., CORIC, S. & HOHENEGGER, J. (2004): Paleoenvironmental changes and agglutinated foraminifera across the Karpatian/Badenian (Early/Middle Miocene) boundary in the Styrian Basin (Austria, Central Paratethys). – In: BÜLK, M. & KAMINISKI, M.A. (Eds.): Proceedings of the Sixth International Workshop on Agglutinated Foraminifera. – Grzybowski Foundation Special publication, **8**, 423–459.
- STILLE, H. (1924): Grundfragen der vergleichenden Tektonik. – 443 pp., Berlin (Bornträger).
- STINGL, K. & SCHOLGER, R. (2005): Detailed magnetic investigations in the Karpatian to Badenian sections of the Retznei and Wagna quarries (Styrian Basin, Austria). – Patterns and Processes in the Neogene of the Mediterranean Region, 12th Congress R.C.M.N.S. – 6.–11. September 2005, Vienna, Abstracts, p. 213.
- STOVER, L.E. (1977): Oligocene and Early Miocene dinoflagellates from Atlantic Corehole 5/5B, Blake Plateau, 66–90. – In: ELSIK, W.C. (Ed.): Contributions of stratigraphic palynology with emphasis on North America. Vol. 1 Cenozoic palynology. – American Association Stratigraphic Palynologists, Contribution Series, **5A**, 66–89.
- STOVER, L.E., BRINKHUIS, H., DAMASSA, S.P., DE VERTEUIL, L., HELBY, R.J., MONTEIL, E., PARTRIDGE, A.D., POWELL, A.J., RIDING, J.B., SMELROR, M. & WILLIAMS, G.L. (1996): Mesozoic – Tertiary dinoflagellates, acritarchs and prasinophytes. – In: JANSONIUS, J. & MCGREGOR, D.C. (Eds.): Palynology: Principles and Applications. – American Association of Stratigraphic Palynologists Foundation, **2**, 641–750.
- STRAUSS, C., LUND, J.J. & LUND-CHRISTENSEN, J. (2001): Miocene dinoflagellate cyst biostratigraphy of the Nieder Ochtenhausen Research Borehole (NW Germany). – Geol. Jb., **A152**, 395–447.
- SÜTÖ-SZENTAI, M. (1994): Microplankton associations of organic skeleton in the surroundings of Villány Mts. – Földtani Közlöny, **124**(4), 451–478.
- SÜTÖ-SZENTAI, M. (1995): Délkelet-Dunántúl ősföldrajzi képe a Panonniai emelet idején. – Folia Comloensis, **6**, 35–55.
- SÜTÖ-SZENTAI, M. (2002): Analysis of microplankton of organic skeleton from borehole Nagykozár-2 (S-Hungary). – Folia Comloensis, **11**, 93–110.
- SÜTÖ-SZENTAI, M. (2003): The organic-walled microplankton in borehole Máriakéménd-3 (Southern Hungary). – Folia Comloensis, **12**, 129–142.
- TORRICELLI, S. & BIFFI, U. (2001): Palynostratigraphy of the Numidian Flysch of northern Tunisia (Oligocene–Early Miocene). – Palynology, **25**, 29–56.
- VAKARCS, G., HARDENBOL, J., ABREU, V.S., VAIL, P.R., VÁRNAI, P. & TARI, G. (1998): Oligocene–Middle Miocene depositional sequences of the Central Paratethys and their correlation with regional stages. – In: GRACIANSKY, P.-C., HARDENBOL, J., JACQUIN, T. & VAIL, P.R. (Eds.): Mesozoic and Cenozoic sequence stratigraphy of European basins. – SEPM, Special Publication, **60**, 209–231.
- WALL, D., DALE, B., LOHMAN, G.P. & SMITH, W.K. (1977): The environmental and climatic distribution of dinoflagellate cysts in modern marine sediments from regions in the North and South Atlantic oceans and adjacent seas. – Marine Micropaleontology, **2**, 121–200.
- WILLIAMS, G.L., BRINKHUIS, H., PEARCE, M.A., FENSOME, R.A. & WEEGINK, J.W. (2004): Southern Ocean and global dinoflagellate cyst events compared: Index events for the Late Cretaceous – Neogene. – In: EXON, N.F., KENNETT, J.P. & MALONE, M.J. (Eds.): Proceedings of the Ocean Drilling Program, Scientific Results, **189**, 1–98.
- WOOD, G.D., GABRIEL, A.M. & LAWSON, J.C. (1996): Palynological techniques - processing and microscopy. – In: JANSONIUS, J. & MCGREGOR, D.C. (Eds.): Palynology: Principles and Applications, American Association of Stratigraphic Palynologists Foundation, **1**, 29–50.
- ZEVENBOOM, D. (1995): Dinoflagellate cysts from the Mediterranean Late Oligocene and Miocene. – CIP-Gegevens Koninklijke Bibliotheek, Den Haag, 221 pp. (Published Ph. D. thesis, State University of Utrecht).

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Jahrbuch der Geologischen Bundesanstalt](#)

Jahr/Year: 2007

Band/Volume: [147](#)

Autor(en)/Author(s): Soliman Ali, Piller Werner E.

Artikel/Article: [Dinoflagellate Cysts at the Karpatian/Badenian Boundary of Wagna \(Styrian Basin, Austria\) 405-417](#)