

Palaeoclimate reconstructions of the last 4000 years in the Southern Northwest territories using ostracods, and implications for the long-term viability of the Tibbitt to Contwoyto Winter Road

Ginny BENARDOUT & R. Timothy PATTERSON

Analysis of ostracod assemblages from cored lacustrine sediments will provide proxy data as a contribution to the project: Palaeoclimatological Assessment of the southern Northwest Territories and implications for the long-term viability of the Tibbitt to Contwoyto Winter Road.




Fig. 1: Location map of the TCWR in Northern Canada showing the TCWR and sample locations. Source: MACUMBER et al. (in prep.).

Funded by the Natural Sciences and Research Council of Canada (NSERC), the multi-proxy project is aimed at providing data from biological and chemical analysis of lake sediment cores and tree rings to track climate change over the last 4000 years in the southern Northwest Territories. With the region predicted to warm at a rate twice as fast as the rest of the world (IPCC 2001, 2007), there are concerns over maintaining the Tibbitt to Contwoyto Winter Road (87% of which runs over frozen lakes; Fig. 1), where projected future warming trends of 2–3°C over the next 50 years (SCHLESINGER & MITCHELL 1987; ACIA 2004) are likely to decrease the annual ice cover by 100 to 130 days. If this predicted warming occurs, changes in ice cover and stability will have drastic implications for the feasibility of mining operations, so there is a crucial need for planners, developers and policy makers to understand the likely impact of future climate variability on the Tibbitt to Contwoyto Winter Road (TCWR).




Fig. 2: Freeze core ROAD10-47-1, face 1 of 1, section 1 of 2: (left) light photograph; (middle) positive X-ray; (right) negative X-ray. The larger bars on the X-ray scale are separated by 5 cm and the smaller bars are separated by 1 cm. Photograph from MACUMBER et al. (in prep.).

Systematic temperature records for the region are available only from the 1940s, and so there is a need for more detailed long-term palaeoclimate information to help understand current short-term climate oscillations in context of the Holocene climate. Three lakes were freeze-cored (Fig. 2) for samples in 2009.

Analysis of ostracod assemblages from the lake core sediments will provide additional palaeoenvironmental and palaeotemperature information for the multi-proxy project (to include diatoms, chironomids, thecamoebians, pollen and dendrochronology), thus helping to develop a more complete understanding of how medium and long-term climate oscillations are correlated to short-term variations.

The project will employ the Mutual Ostracod Temperature Range method (HORNE 2007) to reconstruct palaeotemperatures for the region. Current mean daily January and July temperature ranges are from -17.5°C to -27.5°C and +7.5°C to +17.5°C respectively. Preliminary results and analysis from core material will be presented and discussed.

Acknowledgements

My thanks go to Dr. Dave HORNE for his part in arranging my participation in the project. Thanks also to the School of Geography at Queen Mary, University of London, for providing research funding which has enabled my attendance at this conference.

References

- ACIA (Arctic Climate Impact Assessment, 2004): Impacts of a warming Arctic: Arctic climate impact assessment. – 144 p., Cambridge University Press, Cambridge, UK.
- HORNE, D. J. (2007): A Mutual Temperature Range method for Quaternary palaeoclimatic analysis using European nonmarine Ostracoda. – *Quaternary Science Reviews*, 26: 1398-1415, Oxford.
- IPCC (2001): Intergovernmental Panel on Climate Change Third Assessment Report.
- IPCC (2007): Intergovernmental Panel on Climate Change Fourth Assessment Report.
- MACUMBER, A.L., NEVILLE, L.A., GALLOWAY, J.M., PATTERSON, R.T., FALCK, H., SWINDLES, G., SENKOWSKI, C. & MADSEN, E. (in prep.): Paleoclimatological Assessment of the Southern Northwest Territories and Implications for the Long-Term Viability of the Tibbitt to Contwoyto Winter Road, Part II: March 2010 Field Season Results; Northwest Territories Geoscience Office, NWT Open Report.
- SCHLESINGER, M.E. & MITCHELL, J.F.B. (1987): Climate model simulations of the equilibrium climatic response to increased carbon dioxide. – *Reviews of Geophysics*, 25: 760-798, Washington DC.

Authors addresses:

Ginny Benardout

School of Geography, Queen Mary University of London, Mile End Road, London E1
4NS, UK

g.benardout@qmul.ac.uk

R. Tim Patterson

Department of Earth Sciences, College of Natural Sciences, Carleton University, Ottawa,
Ontario K1S 5B6, Canada

tpatters@earthsci.carleton.ca