

Linzer biol. Beitr.	42/1	617-620	30.7.2010
---------------------	------	---------	-----------

## A new species of *Tetartopeus* from Israel (Coleoptera: Staphylinidae: Paederinae)

B. FELDMANN

**A b s t r a c t :** *Tetartopeus hamulifer* nov.sp., the first representative of the genus from Israel, is described and illustrated.

**K e y w o r d s :** Coleoptera, Staphylinidae, Paederinae, *Tetartopeus*, Palaearctic region, Israel, new species.

### Introduction

The paederine genus *Tetartopeus* CZWALINA 1888 is represented in the Palaearctic region by 32 species (SMETANA 2004; ANLAŞ 2009; ASSING 2004, 2009b). The fact that three of the six known species from Turkey were described only in the last six years, shows that the knowledge of the diversity of the genus in this region is far from complete. Therefore, it does not seem surprising that during an examination of staphylinid material from Israel two specimens of an undescribed species of *Tetartopeus* were found.

### Species description

#### *Tetartopeus hamulifer* nov.sp. (Figs 1-8)

**T y p e m a t e r i a l :** Holotype ♂: "Israel: Ma'agar Bental, 33°08N 35°47'E, 7.v.2007, L. Friedman / Holotypus ♂ *Tetartopeus hamulifer* sp.nov., det. B. Feldmann 2009" (National Collection of Insects, Zoological Museum, Tel Aviv University); Paratype ♂: "Israel (Haifa distr.), Ein Afek Reserve, W Kiryat Motskin (Haifa), (loamy foreshore of brackish ponds), 29.III.2008, D.W. Wrase (37)" (private collection M. Schülke, Berlin).

**D e s c r i p t i o n :** Total length 7.5 mm, length from anterior margin of clypeus to posterior apex of suture 3.5 mm. Habitus as in Fig. 1. Coloration: body blackish, elytra reddish, with the basal and scutellar area more or less infusate; abdomen black with the posterior half of segment VII and the segments VIII-X reddish; legs pale yellowish; antennae reddish-brown, with antennomeres I, II, and basal part of III reddish.

Head 1.13 times as long as wide; punctuation moderately coarse and moderately dense, interstices in lateral and posterior dorsal portions on average approximately as wide as, or slightly wider than diameter of punctures, in median dorsal area distinctly wider than

diameter of punctures; microsculpture in central dorsal area absent, in posterior and lateral area very shallow, interstices shiny. Eyes about half the length of postocular region from posterior margin of eye to neck (Fig. 2). Antenna slender; antennomeres IV-X longer than wide (Fig. 3).

Pronotum 1.26 times as long as wide and 1.09 times as wide as head; punctation similar to that of head; impunctate midline relatively narrow; interstices without microsculpture and shiny (Fig. 2).

Elytra 1.23 times as wide as pronotum and as long as pronotum (Fig. 2); punctation similar to that of forebody or slightly denser; interstices without distinct microsculpture.

Abdomen approximately as wide as elytra; punctation fine and dense; interstices with distinct fine microsculpture and matt; posterior margin of tergite VII with palisade fringe.

♂: posterior margin of tergite VIII weakly convex; posterior margin of sternite VII weakly concave (Fig. 4); sternite VIII with extensive cluster of relatively long black setae, posterior margin in the middle and on either side of middle weakly concave (Fig. 5); aedeagus of distinctive shape (Figs 6-8).

♀: unknown.

**Comparative notes:** As is suggested by the similar morphology of the male primary and secondary sexual characters, *Tetartopeus hamulifer* seems to be closely related to the geographically close *T. persicus* COIFFAIT 1972 and *T. tezcani* ANLAŞ 2009 (for illustrations see ASSING 2008 and ANLAŞ 2009). From both species, *T. hamulifer* is readily separated by the morphology of the aedeagus.


From *T. persicus* (Iran, Iraq, southeastern Anatolia), *T. hamulifer* is additionally distinguished by the slightly different punctation of the head and the elytra (*T. persicus*: punctation of head also present in central dorsal area; punctation of elytra denser).

From *T. tezcani* (central southern Anatolia) the new species is also distinguished by its shorter antennae and its relatively longer elytra.

From other geographically close species with reddish elytra (*T. adanensis* ASSING 2004, *T. czwalinai* (JAKOBSON 1909), *T. inexcisus* ASSING 2009, *T. stylifer* (REITTER 1909)), *T. hamulifer* is distinguished by the distinctive morphology of the aedeagus. For illustrations and information on the distributions of these species see ANLAŞ (2009) and ASSING (2004, 2007, 2008, 2009a, 2009b).

**Etymology:** This specific epithet (Latin, adjective) refers to the hook-like shape of the apical part of the ventral process (lateral view).

**Distribution and bionomics:** *Tetartopeus hamulifer* is the first representative of the genus to become known from Israel. The species was collected in two localities in northern Israel (Golan Heights, Haifa district). The paratype was collected on the loamy foreshore of brackish ponds.



**Figs 1-8:** *Tetartopeus hamulifer* nov.sp. (1-2, 5-8: holotype): (1) habitus; (2) forebody; (3) antenna; (4) male sternite VII; (5) male sternite VIII; (6) aedeagus in lateral view; (7) internal sac of aedeagus; (8) apical portion of aedeagus in lateral view. Scale bars: 1-2: 1.0 mm; 3-6: 0.5 mm; 7-8: 0.1 mm.

### Acknowledgements

I am indebted to Ariel Leib Leonid Friedman (National Collection of Insects, Zoological Museum, Tel Aviv University) and Michael Schülke (Berlin) for the loan of the material under their care. Thanks are due to Volker Assing (Hannover) for the loan of *Tetartopeus tezcani* from his collection, for photographs of the holotype of *Tetartopeus persicus* and for critically proof-reading the manuscript.

### Zusammenfassung

*Tetartopeus hamulifer* nov.sp., der erste Vertreter der Gattung aus Israel, wird beschrieben und abgebildet.



# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Linzer biologische Beiträge](#)

Jahr/Year: 2010

Band/Volume: [0042\\_1](#)

Autor(en)/Author(s): Feldmann Benedikt

Artikel/Article: [A new species of Tetartopeus from Israel \(Coleoptera: Staphylinidae: Paederinae\) 617-620](#)