

Linzer biol. Beitr.	43/2	1219-1228	19.12.2011
---------------------	------	-----------	------------

A Faunistic study on Ichneumonidae(Hymenoptera) in Türkmen Mountain, Turkey

F. EROĞLU, A. KIRAÇ & O. BİROL

A b s t r a c t : In this study, Ichneumonidae fauna of Türkmen Mountain in Eskişehir province was investigated between May 2009 and November 2009. Totally 15 species belonging to 15 genera and 9 subfamilies were found; 5 species from Pimplinae, 2 from both Tryphoninae and Cryptinae, 1 from each of Tersilochinae, Cremastinae, Ichneumoninae, Diplozantinae, Banchinae, and Metopiinae. 2 species noted in the text by asterisk are new records for Turkey fauna. The synonyms, hotspots, general distribution of all species and distribution in Turkey were given on the basis of literature.

K e y w o r d s : Ichneumonidae, fauna, Türkmen Mountain (Eskişehir), new records, Turkey.

Introduction

The Ichneumonidae of Turkey has not been studied well, although it is among the most interesting zoogeographical sections of the Western Palearctic region. In the catalogue of KOLAROV (1995) only 383 species were listed. Lately some new records have been added to this fauna (KOLAROV et al. 1997, 1999; YURTCAN et al. 1999; YURTCAN et al. 2002; ÇORUH et al. 2002, etc.), predominantly from Thrace, Western and Northeastern Turkey. There are several studies about Ichneumonidae in Central Anatolia Region. ÖZDEMİR (2001) listed totally 24 species, 13 species of them from Diplazontinae and 11 from Tryphoninae were obtained. 12 species noted new records for Turkey. The biodiversity hotspots hold especially high numbers of endemic species, yet their combined area of remaining habitat covers only 2.3 percent of the Earth's land surface. Each hotspot faces extreme threats and has already lost at least 70 percent of its original natural vegetation. Over 50 percent of the world's plant species and 42 percent of all terrestrial vertebrate species are endemic to the 34 biodiversity hotspots (MYERS et al. 2000). It is also one of the most remarkable regions of the world in terms of the biodiversity hotspots such that three of them have major extensions into Turkey: Caucasus, Irano-Anatolian and Mediterranean Basin (MYERS et al. 2000).

Material and methods

The Ichneumonidae samples were collected from Türkmen Mountain in Eskişehir Province, between May 2009–November 2009. There were 3 stations appointed to investigate of Ichneumonidae fauna depending on different altitudes (Table 1). The samples were obtained from grass-type plants by using a standard sweeping-net and Malaise trap. The samples are preserved in the collection of the Department of Biology, Faculty of Arts and Sciences at Süleyman Demirel University, Isparta, Turkey.

Table 1: Habitats, altitudes and coordinates of appointed stations.

Lokation	Habitat	Altitude	Coordinates
1. Station	<i>Pinus nigra</i> , <i>Triticum sativum</i> , <i>Hordeum vulgare</i>	1000 m	39° 33' 16.00" N 30° 27' 32.00" E
2. Station	<i>Pinus sylvestris</i> , <i>Quercus pubescens</i>	1300 m	39° 27' 12.00" N 30° 24' 22.00" E
3. Station	<i>Pinus nigra</i> , <i>Pinus sylvestris</i>	1500 m	39° 28' 29.00" N 30° 21' 29.00" E

List of the species

Subfamily Pimplinae

Pimplaspuria (GRAVENHORST 1829)

Pimplaspuria GRAVENHORST 1829 – Ichn. eur. 3: 179.

Material examined: 1. Station, 1000 m, 15.08.2009, 1♀.

General distribution: Western Palearctic, Eastern Palearctic, European, Oriental (YU & HORSTMANN 1997b).

Distribution in Turkey: Ankara-Şereflikoçhisar-AyaşveKaradenizBölgesi (ÖZDEMİR & KILINÇER 1990); Adana-Balcalı, Pozantı, Saimbeyli, Adıyaman-Gölbaşı, Edirne-Hadımağa, Gaziantep-Oğuzeli, Hatay-Dörtöyl-Karıncalı, İçel-Namrun, Kırklareli-İğneada-Limanköy, Vize-Kömürköy, Şanlıurfa-Atatürk Barajı, Tekirdağ-Işıklar (KOLAROV & BEYARSLAN 1994); Balıkesir-İvrindi-Güngörmez, Bilecik-Küplü, Bursa- Aksu, Cumalıkızık, İncirli, Yenişehir, Yeşiltarla, Çanakkale-Bayramiç-Evciler (KOLAROV et al. 1997); Çanakkale-Bozcaada-Merkez, Gökçeada-Aydıncık, Kale, Tigem, Ugurlu (KOLAROV et al. 1997); Erzurum-Oltu-Çamlıbel, Uzundere-Şelale (KOLAROV et al. 1999); Afyon-Bayat-Koroğlu, Sincanlı-Akören, Denizli-Tavas-Tekkeköy, Manisa-Gördes-Güneşli, Muğla-Marmaris-Değirmenyani, Uşak-Banaz (KOLAROV et al. 2002); Çanakkale-Gelibolu-Evreşe, Edirne-TavukOrmanı, İstanbul-Sarıyer-Bilezikçi Çiftliği, Kırklareli-Kızılcıkdere, Tekirdağ-Çerkezköy (YURTCAN 2004), Adana- HalepÇamlığı (BUNCUKÇU 2008).

Hots pots: Irano-Anatolian, Caucasus, Mediterranean Basin, Japan, Mountains of Central Asia.

***Itoplectis maculator* (FABRICIUS 1775)**

Ichneumon maculator FABRICIUS 1775 – Systema Entomologiae, sistens Insectorum classes, ordines, genera, species. Flensburgiet Lipsae. 832 pp.

M a t e r i a l e x a m i n e d : 3. Station, 1500 m, 12.09.2009, 1♂.

G e n e r a l d i s t r i b u t i o n : Western Palearctic, Eastern Palearctic, European, (YU & HORSTMANN 1997b).

D i s t r i b u t i o n i n T u r k e y : Ankara, Eskişehir, Konya, Nevşehir, Yozgat (ÖZDEMİR & KILINÇER 1990), Erzurum (KOLAROV et al. 1999), Çanakkale, İstanbul, Edirne, Kırklareli, Tekirdağ (YURTCAN 2004), Artvin, Bayburt, Erzurum, Gümüşhane, Rize, Kars (ÇORUH et al. 2005), Kasnak Meşesi-Isparta (KIRTAY 2008).

H o t s p o t s : Irano-Anatolian, Caucasus, Mediterranean Basin.

***Zaglyptus multicolor* (GRAVENHORST 1829)**

Polysphincta multicolor GRAVENHORST 1829 – Ich. Eur. Pars III. 1097 pp.

M a t e r i a l e x a m i n e d : 1. Station, 1000 m, 10.10.2009, 1♀.

G e n e r a l d i s t r i b u t i o n : Western Palearctic, Eastern Palearctic, European Middle East (YU & HORSTMANN 1997b).

D i s t r i b u t i o n i n T u r k e y : Adıyaman-Besni, Elazığ-Baskıl (KOLAROV & BEYARSLAN 1994), Isparta-Eğirdir-Gönen (GÜRBÜZ 2004), Kırklareli-Kayalıve Erzurum (ÇORUH 2005); Edirne-Meriç-Küplü, Tekirdağ-Uçmakdere (YURTCAN 2004), Adana-Halep Çamlığı (BUNCUKÇU 2008).

H o t s p o t s : Irano-Anatolian, Caucasus, Mediterranean Basin, Mountains of Central Asia.

***Oxyrrherixis carbonator* (GRAVENHORST 1807)**

Cryptus carbonator GRAVENHORST 1807 – Göttingen 476 pp.

M a t e r i a l e x a m i n e d : 1. Station, 1000 m, 07.06.2009, 1♀; 3. Station, 1500 m 07.06.2009, 1♂; 1. Station, 1000 m, 07.06.2009, 1♂; 3. Station, 1500 m, 27.06.2009, 1♂.

G e n e r a l d i s t r i b u t i o n : Western Palearctic, Eastern Palearctic, European, Nearctic (YU & HORSTMANN 1997b).

D i s t r i b u t i o n i n T u r k e y : Ankara, Eskişehir, Konya, Nevşehir, Yozgat (ÖZDEMİR & KILINÇER 1990), Erzurum (KOLAROV et al., 1999), Çanakkale, İstanbul, Edirne, Kırklareli, Tekirdağ (YURTCAN 2004), Artvin, Bayburt, Erzurum, Gümüşhane, Rize, Kars (ÇORUH et al. 2005).

H o t s p o t s : Irano-Anatolian, Caucasus, Mediterranean Basin, California Floristic Province.

****Acrodactyla quadrisculpta* (GRAVENHORST 1820)**

Ichneumon quadrisculptus GRAVENHORST 1820 – Monographia Ichneumonum Pedemontanae Regionis. Memorie della Reale Accademia delle Scienze di Torino 24: 275-388.

M a t e r i a l e x a m i n e d : 3. Station, 1500 m, 12.09.2009, 1♂.

G e n e r a l d i s t r i b u t i o n : Western Palearctic, Eastern Palearctic, European, Nearctic, Australia, Oriental (YU & HORSTMANN 1997b).

H o t s p o t s : Caucasus, Mediterranean Basin, California Floristic Province.

Subfamily T r y p h o n i n a e

***Tryphonatriceps* (STEPHENS 1835)**

Tryphonatriceps STEPHENS 1835 – III. Of. Brit. Ent., Mandibulata 7: 262.

M a t e r i a l e x a m i n e d : 1. Station, 1000 m, 07.06.2009, 1♂.

G e n e r a l d i s t r i b u t i o n : Gagaus, England, Italy, Lithuania, Central Europe, Moldova, Northern Europe, Russia, Spain, Turkey, Ukraine, Western Europe (YURTCAN et al. 2002).

D i s t r i b u t i o n i n T u r k e y : Edirne, İçel (YURTCAN et al. 2002), Erzurum-İstanbul (KOLAROV 1994).

H o t s p o t s : Irano-Anatolian, Caucasus, Mediterranean Basin.

Netelia (Netelia) fuscicornis (HOLMGREN 1860)

Paniscus fuscicornis HOLMGREN 1860 – Svenska Vetensk. Akad. Hendl. (n.f) 2 (8): 32.

M a t e r i a l e x a m i n e d : 3. Station, 1500 m, 10.10.2009, 1♂.

G e n e r a l d i s t r i b u t i o n : Western Palearctic, Eastern Palearctic, European, Oriental (YU & HORSTMANN 1997b).

D i s t r i b u t i o n i n T u r k e y : Hatay (KOLAROV 1987), Nevşehir-Ürgüp-Göreme-Karain (KOLAROV 1994), Edirne, Elazığ, Kahramanmaraş, Kırklareli, Tekirdağ (KOLAROV & BEYARSLAN 1994b), Kayseri-Erciyes, Konya-Merem, Van, Hatay (KOLAROV 1995a), Balıkesir, Bursa (KOLAROV et al. 1997b), Adana, Ankara, Bayburt, Erzurum (KOLAROV et al. 1999), Ankara, Eskişehir, Kırşehir, Nevşehir (ÖZDEMİR 2001), Edirne, Kırklareli (YURTCAN et al. 2002), Kasnak Meşesi-Isparta (KIRTAY 2008).

H o t s p o t s : Irano-Anatolian, Caucasus, Mediterranean Basin, Himalaya, Mountains of Central Asia, Japan, Western Ghats and Sri Lanka.

Subfamily C r e m a s t i n a e

Pristomerus vulnerator (PANZER 1799)

Ichneumon vulnerator G.W.F. PANZER 1799 – Faunae Insectorum Germanicae. Heft 70-72.

M a t e r i a l e x a m i n e d : 3. Station 1500 m, 12.09.2009, 3♀♀.

G e n e r a l d i s t r i b u t i o n : Europe, China, India, Japan, Korea, Siberia, Turkey (KOLAROV 1995b, 1997b).

D i s t r i b u t i o n i n T u r k e y : Ankara, Bursa, Karadeniz region, Samsun, Tekirdağ (KOLAROV 1995b, 1997b).

H o t s p o t s : Irano-Anatolian, Caucasus, Mediterranean Basin, Mountains of Central Asia, Japan, Western Ghats and Sri Lanka.

Subfamily C r y p t i n a e

Dichrogaster longicaudata (THOMSON 1884)

Hemiteles longicaudatus THOMSON 1884 – Opuscula Entomologica. Lund. X: 939-1028.

M a t e r i a l e x a m i n e d : 2. Station, 1300 m, 15.08.2009, 1♀; 1. Station 1000 m, 15.08.2009, 1♂; 2. Station, 1300 m, 24.05.2009, 1♂.

G e n e r a l d i s t r i b u t i o n : Western Palearctic, Eastern Palearctic, European, Nearctic, (YU & HORSTMANN, 1997b).

D i s t r i b u t i o n i n T u r k e y : Isparta, Eğirdir (KOLAROV & GÜRBÜZ 2007), Kasnak Meşesi-Isparta (KIRTAY 2008).

H o t s p o t s : Mediterranean Basin.

***Cryptus spinosus* (GRAVENHORST 1829)**

Cryptus spinosus GRAVENHORST 1829 – Ichneumologia Europaea. Pars II. Vratislaviae. 989pp.

Material examined: 1. Station, 1000 m, 23.05.2009, 1♂: 3. Station, 1500 m, 18.10.2009, 1♂: 3. Station, 1500 m, 15.08.2009, 1♂.

General distribution: European, Western Palearctic (YU & HORSTMANN 1997b).

Distribution in Turkey: Adana (KOLAROV 1995), Isparta-Gökçay (GÜRBÜZ & KOLAROV 2008).

H o t s p o t s : Mediterranean Basin.

Subfamily D i p l a z o n t i n a e

***Diplazon laetatorius* (FABRICIUS 1781)**

Ichneumon laetatorius FABRICIUS 1781 – Spec. Ins. Tom. I. 552 pp.

Material examined: 1. Station, 1000 m, 18.07.2009, 1♀: 1. Station, 1000 m, 10.10.2009, 1♀.

General distribution: Australia, Palearctic, Ethiopia, Europe, Nearctic, Neotropical (YU & HORSTMANN 1997b).

Distribution in Turkey: Ankara (TUATAY et al. 1972), İstanbul (KOLAROV 1989a), Ankara (ÖNCÜER 1991), Edirne, Tekirdağ, Kırklareli (YURTCAN et al. 1999), Isparta-Şarkikaraağaç-Merkez-Sav-Gönen (GÜRBÜZ 2004), Adana-Halep Çamlığı (BUNCUKÇU 2008).

H o t s p o t s : Irano-Anatolian, Mediterranean Basin, Caucasus, Mountains of Central Asia, Western Ghats and Sri Lanka, Mountains of Southwest China, New Zeland, Japan, Philippines, Indo-Burma, California Floristic Province, Madagascar and Indian Ocean Islands, Cape Floristic Region, Maputaland-Pandoland- Albany, Cerrado Atlantic Forest, Chilean Winter Rainfall-Valdivian Forests, Tropical Andes, Mesoamerica, Madrean Pine- Oak Woodlands.

Subfamily B a n c h i n a e

***Exetastes gracilicornis* (GRAVENHORST 1829)**

Exetastes gracilicornis GRAVENHORST 1829 – Ichneumonologia Europaea Pars. III Vratislaviae.

Material examined: 1. Station, 1000 m, 18.07.2009, 1♀: 3. Station 1500 m, 15.08.2009, 3♂♂: 2. Station, 1300 m, 18.10.2009, 2♂♂, 2♀♀.

General distribution: Western Palearctic, Eastern Palearctic, European, (YU & HORSTMANN 1997b).

Distribution in Turkey: Osmaniye-Fenk Plateau (GÜRBÜZ et al. 2008).

H o t s p o t s : Caucasus, Mediterranean Basin, Mountains of Central Asia.

Subfamily M e t o p i i n a e

***Chorinaeus funebris* (GRAVENHORST 1829)**

Exochus funebris GRAVENHORST 1829 – Ichneumologia Europaea. Pars I. Vratislaviae. 827pp.

Material examined: 1. Station, 1000 m, 10.10.2009, 1♀.

General distribution: Western Palearctic, Eastern Palearctic, European, Nearctic (YU & HORSTMANN 1997b).

Distribution in Turkey: Eskişehir Türkmen Dağı, Trakya (KOLAROV 1989a).

H o t s p o t s : Mediterranean Basin, California Floristic Province.

Subfamily Ichneumoninae***Coelichneumon rudis* (FONSCOLOMBE 1847)**

Ichneumon rudis FONSCOLOMBE 1847 – Ichneumologie provençale. Annales de la société Entomologique de France (2) 5: 51-70.

Material examined: 2. Station, 1300 m, 10.10.2009, 2♀♀: 1. Station, 1000 m, 10.10.2009, 2♀♀.

General distribution: Western Palearctic, European (YU & HORSTMANN 1997b).

Distribution in Turkey: Adana-HalepÇamlığı (GÜRBÜZ et al. 2008).

Hotspots: Mediterranean Basin.

Subfamily Tersilochinae****Probles versutus* (HOLMGREN 1860)**

Thersilochus versutus HOLMGREN 1860 – Monographia ophionidum Sueciae. Kongliga Svenska Vetenskaps akademiens Handlingar 2 (8): 1-158.

Material examined: 3. Station, 1500m, 15.08.2009, 1♀: 1. Station, 1000 m, 15.08.2009, 2♂♂: 2. Station, 1300 m, 15.08.2009, 2♂♂.

General distribution: Western Palearctic, European (YU & HORSTMANN 1997b), Bulgaria (KOLAROV 1987).

Hotspots: Mediterranean Basin.

Result and discussion

The dominance (D) describes the relative abundance of a species within a community. (TISCHER 1977).

It is measured as:

$$D = \frac{b}{a} \times 100$$

b= number of individuals of the species
a= number of individuals of all species

Dominance scale logarithmic after ENGELMANN (MACZEY 2004).

- Eudominant >32.0 ≤ 100 %
- Dominant > 10.0 ≤ 32.0 %
- Subdominant > 3.2 ≤ 10.0 %
- Recedent > 1.0 ≤ 3.2 %
- Subrecedent > 0.32 ≤ 1.0 %
- Sporadic ≤ 0.32 %
- Missing = 0 %

According to ENGELMANN's (1978) dominance scale, *Exetastes gracilicornis* and *Probles versutus* are dominant species. *Oxyrrherix carbonator*, *Cryptus spinosus*, *Dichrogaster longicaudata*, *Diplazon laetatorius*, *Coelichneumon rudis* and *Pristomerus*

vulnerator are subdominant species. *Pimplaspuria*, *Itopectis maculator*, *Zaglyptus multicolor*, *Acrodactyla quadrisculpta*, *Tryphon atriceps*, *Netelia fuscicornis* and *Chorinaeus funebris* are recedent species. There is not any eudominant, subrecedent, sporadic and missing species in the research area (Table 2).

Table 2: Dominance of species and hotspots

Species	1. Station	2. Station	3. Station	D %	Category	Hotspots
<i>Pimplaspuria</i>	+	-	-	2.38	Recedent	Common
<i>Itopectis maculator</i>	-	-	+	2.38	Recedent	Common
<i>Zaglyptus multicolor</i>	+	-	-	2.38	Recedent	Common
<i>Acrodactyla quadrisculpta</i>	-	-	+	2.38	Recedent	Mediterranean Basin and Caucasus
<i>Oxyrrherix carbonator</i>	+	-	+	9.52	Subdominant	Common
<i>Tryphona triceps</i>	+	-	-	2.38	Recedent	Common
<i>Netelia fuscicornis</i>	-	-	+	2.38	Recedent	Common
<i>Cryptus spinosus</i>	+	-	+	7.14	Subdominant	Mediterranean Basin
<i>Dichrogaster longicaudata</i>	+	+	-	7.14	Subdominant	Mediterranean Basin
<i>Diplazon laetatorius</i>	+	-	-	4.76	Subdominant	Mediterranean Basin and Caucasus
<i>Exetastes gracilicornis</i>	+	+	+	16.67	Dominant	Mediterranean Basin and Caucasus
<i>Chorinaeus funebris</i>	+	-	-	2.38	Recedent	Mediterranean Basin
<i>Coelich neumonrudis</i>	+	+	-	9.52	Subdominant	Mediterranean Basin
<i>Probles versutus</i>	+	+	+	11.90	Dominant	Mediterranean Basin
<i>Pristomerus vulnerator</i>	-	-	+	7.14	Subdominant	Common

D= Dominance, (+)= Species are shown in the research area, (-)= Species are not shown in the research area

Within this study identified two species are new records for Turkish fauna. These are *Probles versutus* and *Acrodactyla quadrisculpta*. Turkey has three hotspots. These are Caucasus, Irano-Anatolian and Mediterranean Basin (MYERS et al. 2000). The Türkmen Mountain is not included in these three hotspots, but Türkmen Mountain is near the Irano-Anatolian and Mediterranean Basin hotspots. *Acrodactyla quadrisculpta* found Mediterranean Basin and Caucasus hotspots (Table 2). Probably, *Acrodactyla quadrisculpta* will arrive at Irano-Anatolian hotspot, because Türkmen Mountain is near the Irano-Anatolian hotspot. Also *Probles versutus* was found only in the Mediterranean Basin hotspot. The results showed that Türkmen Mountain is an important transit area between Mediterranean Basin and Irano-Anatolian hotspots.

Acknowledgments

We would like to thank Dr. Janko Kolarov (Plovdiv University, Bulgaria) and Dr. M. Faruk Gürbüz (Süleyman Demirel University, Turkey) for his valuable helps. This is part of the Eroğlu's Msc. Thesis.

Zusammenfassung

Die Ichneumonidenfauna (Hymenoptera, Ichneumonidae) des Berges Türkmen in der türkischen Provinz Eskişehir wurde im Zeitraum Mai 2009 bis November 2009 untersucht. 15 Arten aus 15 Gattungen und 9 Unterfamilien wurden nachgewiesen, darunter 5 Arten Pimplinae, je zwei Arten Tryphoninae und Cryptinae, sowie je eine Art aus den Unterfamilien Tersilochinae, Cremastinae, Ichneumoninae, Diplozantinae, Banchinae und Metopiinae. Zwei Arten (*Acrodactyla quadrisculpta*, *Probes versutus*) stellen Erstnachweise für die Türkei dar. Auf Basis von Literaturangaben wurden Angaben zu Verbreitungen und Synonymien gegeben.

References

- BUNCUKÇU A. (2008): Isparta İli Merkezive Adana-Yumurtalık İlçesi Halep Çamlığı Ichneumonidae Türlerinin Tespiti ve Kültüre Edilebilen Türlerin Biyolojisinin Araştırılması. — Süleyman Demirel Üni. Fen Bilimleri Enst., Yüksek Lisans Tezi, 74 pp, Isparta.
- ÇORUH S. (2005): Erzurum ve Çevre İllerdeki Pimplinae (Hymenoptera: Ichneumonidae) Türleri Üzerinde Faunistik, Sistemantik ve Ekolojik Çalışmalar. — Atatürk Üni. Fen Bilimleri Enst., Doktora Tezi, 211pp, Erzurum.
- ÇORUH S., ÖZBEK H. & J. KOLAROV (2002): New and rare Taxa of Ichneumonidae (Hymenoptera) from Turkey. — J. Ent. Res. Soc. **4** (1): 1-4.
- ÇORUH S., ÖZBEK H. & J. KOLAROV (2005): A Contribution to the Knowledge of Ichneumonidae (Hymenoptera) from Turkey. — Journal of the Entomological Research Society: **7** (3): 53-57.
- ENGELMANN H.D. (1978): Zur Dominanzklassifizierung von Bodenarthropoden. — Pedobiologia **18**: 378-380.
- GÜRBÜZ M.F. (2004): Isparta İli Ichneumonidae (Hymenoptera) Familyası Türleri Üzerine Faunistik ve Sistemantik Araştırmalar. — Süleyman Demirel Üni. Fen Bilimleri Enst., Doktora Tezi, 68pp, Isparta.
- GÜRBÜZ M.F. & J. KOLAROV (2008): A Study of the Turkish Cryptini IV. (Cryptinae: Ichneumonidae: Hymenoptera). — Turk. J. Zool. **32**: 373-377.
- GÜRBÜZ M.F., LJUBOMIROV T., KOLAROV J., YURTCAN M., TABUR M.A., ÇORUH S. & A. BUNCUKÇU (2008): Investigation of the Ichneumonidae, Ampulicidae, Crabronidae and Sphecidae (Hymenoptera, Insect) Fauna in Natural Protection Zones of East Mediterranean Region in Turkey. — TBAGU/ **168** (106T189): 126.
- KIRTAY H. (2008): Isparta Kasnak Meşesi Ormanı Tabiatı Koruma Alanı Ichneumonidae (Hymenoptera) Faunası Üzerine Bir Araştırma. — Süleyman Demirel Üni. Fen Bilimleri Enst., Yüksek Lisans Tezi, 77pp, Isparta.
- KOLAROV J. (1987): Ichneumonidae (Hymenoptera) from Balkan Peninsula and some Adjacent Regions. I. Pimplinae, Tryphoninae, Cryptinae. — Turk. Ent. D. **11** (1): 11-28.
- KOLAROV J. (1989a): Ichneumonidae (Hymenoptera) from Balkan peninsula and some adjacent regions. III. Ophioninae, Anamaloniae, Metopiinae, Mesochorinae, Acaenitinae, Oxytorinae, Orthopelmatinae, Collyriinae, Orthocentrinae, Diplazontinae and Ichneumoninae. — Türkiye Entomoloji Dergisi **13** (3): 131-140.

- KOLAROV J. (1994): Nocturnal Ichneumonidae from Bulgaria and Turkey with description of a new species. — *Entomofauna* **15**: 93-97.
- KOLAROV J. (1995): A catalogue of the Turkish Ichneumonidae (Hymenoptera). — *Entomofauna* **7**: 137-188.
- KOLAROV J. (1995a): A catalogue of the Turkish Ichneumonidae (Hymenoptera). — *Entomofauna* **7**: 137-188.
- KOLAROV J. (1995b): Cremastinae (Hymenoptera: Ichneumonidae) from Italy and some adjacent regions. — *Linzer biol. Beitr.* **27** (2): 1104-1114.
- KOLAROV J. (1997b): A review of the Cremastinae of the Balkan Peninsula, Turkey and Cyprus with zoogeographical notes. — *Beitr. Ent.* **47** (1): 169-199.
- KOLAROV J. & A. BEYARSLAN (1994b): Investigations on the Ichneumonidae (Hymenoptera) fauna of Turkey I. Pimplinae and Tryphoninae. — *Türk. Ent. D.* **18**(3): 133-140.
- KOLAROV J., BEYARSLAN A. & M. YURTCAN (1997): New and Rare Ichneumonidae (Hymenoptera) from Turkey. I. Pimplinae, Tryphoninae, Phyeuontinae, Banchinae and Ctenopelmatinae. — *Acta. Zoo. Bulg.* **3-4**: 10-12.
- KOLAROV J., BEYARSLAN A. & M. YURTCAN (1997b): New and Rare Ichneumonidae (Hymenoptera) from Turkey. I. Pimplinae, Tryphoninae, Phyeuontinae, Banchinae and Ctenopelmatinae. — *Acta Zoologica Bulgarica*: 3-4, 10-12.
- KOLAROV J. & M.F. GÜRBÜZ (2007): A Study of the Turkish Ichneumonidae (Hymenoptera). V. Cryptinae. — *Phygadeuontini*: 987-992.
- KOLAROV J., ÖZBEK H. & E. YILDIRIM (1999): New distributional data of the Turkish Ichneumonidae (Hymenoptera). I. Pimplinae and Tryphoninae. — *J. Ent. Soc. Res.* **1** (2): 9-15.
- KOLAROV J., YURTCAN M. & A. BEYARSLAN (2002): Ichneumonidae species of the Turkish Aegean Region. Parasitic Wasp: evolution, systematics, biodiversity and biological control. — International symposium Kőszeg-Hungary, George Melika and Csaba Thuroczy (editors): 299-305.
- MACZEY N. (2004): The Auchenorrhyncha Communities of Chalk Grassland in southern England. — *Naturwissenschaft Fachbereich 3: Mathematik-Naturwissenschaften Universität Koblenz, Dissertation, Landau* 230pp.
- MYERS N., MITTERMEIER R.A., MITTERMEIER C.G., da FONSECA G.A.B. & J. KENTS (2000): Biodiversity Hotspots for Conservation Priorities. — *Nature* **403**: 853-858.
- ÖNCÜER C. (1991): A Catalogue of the Parasites and Predators of Insect Pests of Turkey. — *Ege Üniv. Ziraat. Fak. Yayınları* **505**: 1-354.
- ÖZDEMİR Y. (2001): İç Anadolu Bölgesinde Saptanan Diplazontinae ve Tryphoninae (Hymenoptera: Ichneumonidae) Türleri. — *Türkiye Ent. Der.* **25** (3): 183-191.
- ÖZDEMİR Y. & N. KILINÇER (1990): The Species of Pimplinae and Ophioninae from Central Anatolia. — *Türk II. Bio. Múc. Kong. Ankara*: 309-318.
- TISCHLER W. (1977): Einführung in die Ökologie. — Fischer, Stuttgart.
- TUATAY N., KALKANDELEN A. & N. ÇAĞATAY (1972): Nebat Koruma Müzesi Böcek Katalogu. — *Zir. Múc. VeZir. Karan. Gn. Md. Yayınları Mesleki Kitaplar Serisi*: 119.
- YU D. & K. HORSTMANN (1997b): Catalogue of World Ichneumonidae (Hymenoptera). — *The Amer. Ent. Inst. Part 1*, **2**: 1-1558.
- YURTCAN M. (2004): Trakya Bölgesi Pimplinae (Hymenoptera: Ichneumonidae) Faunasının Taksonomik ve Faunistik Yönünden Araştırılması. — *Trakya Üni. Fen bilimleri Enst., Doktora Tezi*, 110pp, Edirne.
- YURTCAN M., BEYARSLAN A. & J. KOLAROV (1999): Investigations on the Ichneumonidae (Hymenoptera) Fauna of Turkey. V. Diplazontinae and Ichneumoninae. — *Acta. Ent. Bulg.* **5**(1): 34-36.
- YURTCAN M., BEYARSLAN A. & J. KOLAROV (2002): The Species of Tryphoninae (Hymenoptera: Ichneumonidae) in Turkish Thrace. — *Turk. J. Zool.* **26**: 77-95.

Addresses of the authors: Fatma EROĞLU
Süleyman Demirel University, Faculty of Arts and Sciences
Department of Biology
TR-32260 Isparta, Turkey
E-Mail: fatma.eroglu73@gmail.com

Akın KIRAC
Süleyman Demirel University, Faculty of Arts and Sciences
Department of Biology
TR-32260 Isparta, Turkey
E-mail: akirac20@hotmail.com

Onur BİROL (Corresponding Author)
Süleyman Demirel University, Faculty of Arts and Sciences
Department of Biology
TR-32260 Isparta, Turkey
E-mail: onurbirol@sdu.edu.tr

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Linzer biologische Beiträge](#)

Jahr/Year: 2011

Band/Volume: [0043_2](#)

Autor(en)/Author(s): Eroglu Fatma, Kirac Akin, Birol Onur

Artikel/Article: [A Faunistic study on Ichneumonidae\(Hymenoptera\) in Türkmen Mountain, Turkey 1219-1228](#)