

Linzer biol. Beitr.	45/1	673-679	31.7.2013
---------------------	------	---------	-----------

## Contribution to the Knowledge of the Ladybird beetles (Coleoptera, Coccinellidae), Predators of Mealybugs (Hemiptera, Pseudococcidae) in Hormozgan Province, Southern Iran

M. FALLAHZADEH, R. ABDIMALEKI & N. SAGHAEI

**A b s t r a c t :** The Coccinellid predators (Coleoptera, Coccinellidae) associated with mealybug species (Hemiptera, Pseudococcidae) in Hormozgan province, south of Iran was studied during years 2009-2010. A total of 8 species representing 5 genera belonging to 3 subfamilies were collected and identified. *Nephus (Sidis) hieckei riyadhensis* FÜRSCH 1979 is new record for the fauna of Iran. Locality and date of collection, host(s) and distribution data for each species are provided.

**K e y   w o r d s :** Coleoptera, Coccinellidae, Predators, Pseudococcidae, Hormozgan, Iran.

### Introduction

The Coccinellidae comprise one of the largest and most abundant families of Cucujoidae with over 6.000 nominal species known throughout the world (CANEPARI 2011). Most Coccinellidae are entomophagous, predators of a wide range of hosts from hemipteran pests such as aphids, mealybugs and scale insects, as well as thrips (Thysanoptera) and mites (Acarina) in all parts of the world (RAIMUNDO et al. 2006, 2008). They represent the largest group of predators of mealybugs (Hemiptera, Pseudococcidae) (MAJERUS 1994). Several species of coccinellid beetles have been used or are used today for suppression of various mealybug crop pests (BARTLETT 1978, BEN-DOV 1994, BEN-DOV et al. 2012). Recently, some faunistic studies on Iranian Coccinellids fauna have been generally focused on predators associated with mealybugs (ASADEH & MOSSADEGH 1991, MOSSADEGH & KOCHEILI 1993, NOVIN 2000, LOTFALIZADEH & AHMADI 2000, FALLAHZADEH & HESAMI 2004, FALLAHZADEH et al. 2006, 2010, 2011). Here, we provide some additional information on the distribution of Coccinellid predators associated with mealybug in Hormozgan province southern Iran, add one subspecies to the published records, and provide some new host records.

### Materials and Methods

Surveys for predators of mealybug (Hemiptera, Pseudococcidae) were conducted on cultivated and non-cultivated plants in different area of Hormozgan province, Iran, from

2009 to 2010. During each sample, predators were collected using an aspirator, when observed, or by shaking plant to dislodge and collect predators in a funnel below. In the laboratory, immature stages of the predators were reared, on mealybugs, to the adult stage for better identification (using keys by YAZDANI 1990, RAIMUNDO & VAN HARTEN 2000, RAIMUNDO et al. 2006, 2008). Voucher specimens were deposited in the Entomology Department of the Agriculture Faculty, Islamic Azad University, Jahrom Branch, Iran.

## Results and Discussion

A total of 8 Coccinellid species representing 5 genera belonging to 3 subfamilies were collected. The list of species and their distributional data are given below.

### **Subfamily Scymninae MULSANT 1846**

#### **Tribe Scymniini MULSANT 1846**

##### **Genus *Scymnus* KUGELANN 1794**

###### ***Scymnus (Scymnus) nubilus* MULSANT 1850**

**M a t e r i a l :** 2♀♀, 3♂♂, Rodan, 15.09.2009, ex *Nipaecoccus viridis* (NEWSTEAD) on *Citrus* (Rutaceae); 13♀♀, 1♂, Minab, 18.02.2010, ex *N. viridis* on *Citrus*; 3♀♀, 4♂♂, Minab, 18.02.2010, ex *Maconellicoccus hirsutus* (GREEN) on *Abutilon fruticosum* GUILL & PERR. (Malvaceae); 8♀♀, 4♂♂, Rodan, 9.10.2010, ex *N. viridis* on *Citrus*; 14♀♀, 10♂♂, Bandar-e-Abbas, 14.10.2010, ex *M. hirsutus* on *Ricinus communis* L. (Euphorbiaceae); 6♀♀, 1♂, Bandar-e-Abbas, 14.10.2010, ex *M. hirsutus* on *Helianthus annuus* L. (Asteraceae); 4♂♂, Bandar-e-Abbas, 14.10.2010, ex *M. hirsutus* on *Ficus religiosa* L. (Moraceae).

**R e m a r k s :** It has been recorded as a predator of the mealybug *Maconellicoccus hirsutus* (GREEN) (KUMAR & CHAKRABORTY 1997).

**D i s t r i b u t i o n :** Southern Europe, Atlantic islands eastwards to Japan and Micronesia, Africa (CANEPARI 2001), Iran, Yemen and UAE (YAZDANI 1990, RAIMUNDO & VAN HARTEN 2000, RAIMUNDO et al. 2006, 2008).

###### ***Scymnus (Pullus) syriacus* (MARSEUL 1868)**

**M a t e r i a l :** 7♂♂, Bandar-e-Abbas, 23.04.2009, ex *Planococcus ficus* (NASONOV) on *Ficus* sp. (Moraceae); 13♀♀, 1♂, Bandar-e-Abbas, 30.5.2009, ex *Phenacoccus solenopsis* (TINSLEY) on *Abutilon fruticosum* GUILL & PERR. (Malvaceae), 3♀♀, 2♂♂, 4.12.2009, ex *P. solenopsis* on *Acacia nilotica* (L.) WILLD. ex DELILE (Mimosaceae); 6♀♀, 11♂♂, Minab, 18.02.2010, ex *P. solenopsis* on *Hibiscus rosa-sinensis* L. (Malvaceae); 8♀♀, 5♂♂, Rodkhaneh, 15.06.2010, ex *P. solenopsis* on *Gossypium* L. (Malvaceae).

**R e m a r k s :** It has been recorded as a predator of the mealybug *Nipaecoccus viridis* (NEWSTEAD) (SHARAF & MEYERDIRK 1987), *Ferrisia virgata* (COCKERELL) (MANGOUD 2008).

**D i s t r i b u t i o n :** Eastern Mediterranean region especially Jordan, Lebanon, Turkey, Iran, Libya, Yemen and Egypt (YAZDANI 1990, RAIMUNDO & VAN HARTEN 2000, RAIMUNDO et al. 2006).

## Genus *N e p h u s* MULSANT 1846

### *Nephus (Sidis) hieckei riyadhensis* FÜR SCH 1979

M a t e r i a l : 13♀♀, 6♂♂, Bandar-e-Abbas, 30.11.2009, ex *P. solenopsis* on *Sesamum orientale* L. (Pedaliaceae); 20♀♀, 4♂♂, Bandar-e-Abbas, 14.06.2010, ex *P. solenopsis* on *Lantana camara* L (Verbenaceae); 5♀♀, 2♂♂, Minab, 18.09.2010, ex *P. solenopsis* on *Hibiscus rosa-sinensis* L. (Malvaceae); 7♀♀, Rodan, 19.04.2010, ex *P. solenopsis* on *Solanum nigrum* L. (Solanaceae).

R e m a r k s : The nominal species has been recorded as a predator of the mealybug *Planococcus citri* (RISSE) (FÜR SCH 1979).

D i s t r i b u t i o n : The nominal species has been recorded from Portugal, Spain, Greece, Turkey, Palestine, Lebanon and Iraq. The subspecies was only known from Egypt, Syria, Yemen, Saudi Arabia and UAE (FÜR SCH 1979, RAIMUNDO & VAN HARTEN 2000, RAIMUNDO et al. 2006, 2008)

### *Nephus (Geminosipho) fenestratus* (SAHLBERG 1913)

M a t e r i a l : 6♀♀, 10♂♂, Rodan, 15.09.2009, ex *N. viridis* on *Citrus* (Rutaceae); 9♀♀, 1♂, Minab 18.02.2010, ex *N. viridis* on *Citrus*; 1♀, 3♂♂, Rodan, 3.09.2010, ex *N. viridis* on *Citrus*; 4♀♀, 2♂♂, Minab 12.06.2010, ex *N. viridis* on *Citrus*; 5♀♀, 1♂, Rodan, 16.07.2010, ex *N. viridis* on *Citrus*; 4♀♀, 2♂♂, Rodan, 12.10.2010, ex *N. viridis* on *Citrus*.

R e m a r k s : It has been recorded as a predator of the mealybug *Nipaecoccus viridis* (NEWSTEAD) (ASA DEH & MOSSADEGH 1991, MOSSADEGH & KO CHEILI 1993, NOVIN 2000, HESAMI & FALLAHZADEH 2004), *Maconellicoccus hirsutus* (GREEN) (FALLAHZADEH & HESAMI 2004).

D i s t r i b u t i o n : Egypt, Syria, Iran, Saudi Arabia (FÜR SCH 1979), Yemen and UAE (RAIMUNDO & VAN HARTEN 2000; RAIMUNDO et al. 2006, 2008).

### *Nephus (Bipunctatus) includens* (KIRSCH 1870)

M a t e r i a l : 4♂♂, Rodan, 15.09.2009, ex *N. viridis* on *Citrus* (Rutaceae); 19♀♀, 3♂♂, Minab 18.02.2010, ex *N. viridis* on *Citrus*; 17♀♀, 16♂♂, Rodan, 3.09.2010, ex *N. viridis* on *Citrus*; 12♀♀, 4♂♂, Minab 12.06.2010, ex *N. viridis* on *Citrus*; 1♀, 2♂♂, Rodan, 16.07.2010, ex *N. viridis* on *Citrus*; 54♀♀, 3♂♂, Rodan, 14.10.2010, ex *N. viridis* on *Citrus*; 6♀♀, Rodan, 17.12.2010, ex *N. viridis* on *Citrus*; 1♂, Rodan, 20.12.2010, ex *N. viridis* on *Citrus*.

R e m a r k s : It has been recorded as a predator of the mealybug *Nipaecoccus viridis* (Newstead) (ASA DEH & MOSSADEGH 1991, MOSSADEGH & KO CHEILI 1993, NOVIN 2000), *Planococcus citri* (RISSE) (ARGYRIOU et al. 1976, KONTODIMAS et al. 2005).

D i s t r i b u t i o n : West Africa (Gambia, Guinea-Bissau, Cape Verde, Canary Islands); Mediterranean Europe, Africa and Near East (Portugal, Greece, Sicily, Turkey, Egypt, Israel, Syria, Ethiopia, Saudi Arabia, Iran, Yemen and UAE (FÜR SCH 1979, 1986, RAIMUNDO 1992, 1994, RAIMUNDO & VAN HARTEN 2000, RAIMUNDO et al. 2006, 2008).

## Tribe Hypoperaspidini MULSANT 1846

### Genus *Hypoperaspis* DEJEAN 1833

#### *Hyperaspis polita* WEISE 1885

Materiel: 4♀♀, 1♂, Minab, 30.04.2009, ex *M. hirsutus* on *Abutilon fruticosum* GUILL & PERR. (Malvaceae); 6♀♀, 1♂, Rodan, 15.09.2009, ex *N. viridis* on *Citrus*; 8♀♀, Bandar-e-Abbas, 30.11.2009, ex *P. solenopsis* on *Sesamum orientale* L. (Pedaliaceae); 1♂, Bandar-e-Abbas, 30.05.2009, ex *N. viridis* on *Ziziphus mauritiana* Lam. (Rhamnaceae); 2♀♀, 6♂♂, Rodan, 3.09.2010, ex *N. viridis* on *Citrus*; 3♂♂, Bandar-e-Abbas, 30.11.2009, ex *P. solenopsis* on *Sesamum orientale* L. (Pedaliaceae); 2♂♂, Bandar-e-Abbas, 14.06.2010, ex *P. solenopsis* on *Lantana camara* L (Verbenaceae); 4♀♀, 1♂, Minab, 18.09.2010, ex *P. solenopsis* on *Hibiscus rosa-sinensis* L. (Malvaceae).

Remarks: It has been recorded as a predator of the mealybug *Nipaecoccus viridis* (NEWSTEAD) (ASADEH & MOSSADEGH 1991, MOSSADEGH & KOCHEILI 1993, NOVIN 2000, HESAMI & FALLAHZADEH 2004), *Planococcus vovae* (NASONOV) (LOTFALIZADEH & AHMADI 2000), *Maconellicoccus hirsutus* (GREEN) (FALLAHZADEH & HESAMI 2004), *Planococcus ficus* (SIGNORET) (FALLAHZADEH et al. 2011).

Distribution: Lebanon and the coastal areas of the Eastern Mediterranean region, Turkey, Iran to Pakistan and dry regions in South Western Asia (YAZDANI 1990, RAIMUNDO & VAN HARTEN 2000, RAIMUNDO et al. 2006).

## Subfamily Chilocorinae MULSANT 1846

### Genus *Exochomus* REDTENBACHER 1843

#### *Exochomus melanocephalus* (ZOUBKOFF 1833)

Materiel: 2♀♀, 6♂♂, Bandar-e-Abbas, 30.05.2009, ex *N. viridis* on *Ziziphus mauritiana* LAM. (Rhamnaceae); 20♀♀, 1♂, Bandar-e-Abbas, 21.06.2009, ex *N. viridis* on *Z. mauritiana*; 4♀♀, 2♂♂, Bandar-e-Abbas, 7.08.2010, ex *N. viridis* on *Z. mauritiana*.

Remarks: It has been recorded as a predator of the mealybug *Ferrisia virgata* (COCKERELL) (LE PELLEY 1968).

Distribution: South Russia, Transcaucasia, Iran (MADER 1955, YAGHMAEI & KHARAZI PAKDEL 1995).

## Subfamily Coccinellinae LATREILLE 1807

### Genus *Menochilus* TIMBERLAKE 1943

#### *Menochilus sexmaculatus* (FABRICIUS 1781)

Materiel: 2♀♀, Minab, 30.04.2009, ex *M. hirsutus* on *Abutilon fruticosum* GUILL & PERR. (Malvaceae); 1♀, 7♂♂, Bandar-e-Abbas, 30.11.2009, ex *P. solenopsis* on *Sesamum orientale* L. (Pedaliaceae); 1♀, Bandar-e-Abbas, 14.06.2010, ex *P. solenopsis* on *Lantana camara* L (Verbenaceae); 5♂♂, Minab, 18.09.2010, ex *P. solenopsis* on *Hibiscus rosa-sinensis* L. (Malvaceae); 5♀♀, 9♂♂, Rodan, 19.04.2010, ex *P. solenopsis* on *Solanum nigrum* L. (Solanaceae); 4♂♂, Bandar-e-Abbas, 14.10.2010, ex *M. hirsutus* on *Ricinus communis* L. (Euphorbiaceae).

Remarks: It has been recorded as a predator of the mealybug *Phenacoccus*

*solenopsis* (TINSLEY) (ZHOU et al. 2012).

Distribution: Japan, China, Formosa, India, Philippines, Ceylon, Malaysia, New Guinea, Micronesia, Indonesia, Timor and UAE (RAIMUNDO et al. 2008).

Members of the family Coccinellidae are questionably one of the most useful groups in biological control, whereas several species of them are used in biological control programs of scale insect species (MAJERUS 1994), particularly on the mealybugs (BEN-DOV et al. 2012). In our survey, each species was directly observed feeding on eggs, nymphs, and adults of mentioned mealybug species. The *Nephus (Sidis) hieckeи riyadhensis* FÜRSCH 1979 is new record for Iran. More work is needed to document the effectiveness of the native lady beetles, found throughout the world's regions, as mealybug predators.

### Acknowledgements

The authors are indebted to Dr. H. FÜRSCH (Germany) and Dr. C. CANEPARI (Italy) for invaluable help with the project. The research was supported by Islamic Azad University, Jahrom Branch, Iran.

### Zusammenfassung

Die räuberischen Coccinelliden (Coleoptera, Coccinellidae), assoziiert mit Schmierläusen (Hemiptera, Pseudococcidae), wurden in der südiranischen Provinz Hormozgan im Zeitraum 2009-2010 studiert. 8 Arten aus 5 Gattungen und 3 Unterfamilien konnten nachgewiesen und dokumentiert werden, *Nephus (Sidis) hieckeи riyadhensis* FÜRSCH 1979 stellte sich als Erstnachweis für den Iran heraus.

### References

- ARGYRIOU L.C., STAVRAKI H.G. & P.A. MOURIKIS (1976): A list of recorded entomophagous insects of Greece. — Benaki Phytopathological Institute, Athens. 73 pp.
- ASADEH G.A. & M.S. MOSSADEGH (1993): Important natural enemies of mealybugs (*Pseudococcus* spp.) in the Khuzestan province Iran. — Scientific Journal of Agriculture Shahid Chamran University **16** (1-2): 46-52.
- BARTLETT B.R. (1978): Pseudococcidae. — In: CLAUSEN C.P. (ed.), Introduced Parasites and Predators of Arthropod Pests and Weeds: a World Review. — Agricultural Research Service, United States Department of Agriculture, Washington, D.C.: 137-170.
- BEN-DOV Y. (1994): A Systematic Catalogue of the Mealybugs of the World (Insecta: Homoptera: Coccoidea: Pseudococcidae and Putoidae) with Data on Geographical Distribution, Host Plants, Biology and Economic Importance. — Intercept Ltd., Andover, U.K., 686 pp.
- BEN-DOV Y., MILLER D.R. & G.A.P. GIBSON (2012): ScaleNet. — Available on-line at: <http://www.sel.barc.usda.gov/scalenet/scalenet.htm>. Accessed: 2 February 2012.
- CANEPA RI C. (2011): Contribution to the knowledge of the Coccinellidae of Sardinia (Coleoptera). — Conservazione Habitat Invertebrati **5**: 501-516.
- FALLAHZADEH M. & S. HESAMI (2004): Study of the natural enemies of *Maconellicoccus hirsutus* (Homoptera: Pseudococcidae) in Jahrom region of Fars province. — Proceeding of the 16<sup>th</sup> Plant Protection Congress of Iran, Tabriz 28 August-1 September 2004 Tabriz, University of Tabriz, Iran, pp 47.

- FALLAHZADEH M., KAYDAN M.B. & F. KOZÁR (2010): Description of a new species of *Chorizococcus* (Hemiptera: Coccoidea: Pseudococcidae) infesting *Vitis vinifera* in Iran. — *Türkiye Entomoloji Dergisi* **34** (2): 157-163.
- FALLAHZADEH M., JAPOSHEVILI G., SAGHAEI N. & K.M. DAANE (2011): Natural enemies of *Planococcus ficus* (Hemiptera: Pseudococcidae) in Fars Province vineyards, Iran. — *Biocontrol Science and Technology* **21** (4): 427-433.
- FALLAHZADEH M., SHOJAEI M., OSTOVAN H. & K. KAMALI (2006): The first report of two parasitoid wasps on the larvae of *Nephus bipunctatus* (Col., Coccinellidae) from Iran. — *Journal of Entomological Society of Iran* **26** (1): 95-96.
- FÜRSCH H. (1979): Coleoptera: Fam. Coccinellidae. — *Fauna of Saudi Arabia* **1**: 235-248.
- HESAMI S. & M. FALLAHZADEH (2004): Study of the natural enemies of the citrus mealybug *Nipaecoccus viridis* (Homoptera: Pseudococcidae) in Jahrom region of Fars province. — Proceeding of the 16<sup>th</sup> Plant Protection Congress of Iran, Tabriz 28 August-1 September 2004 Tabriz, University of Tabriz, Iran, pp 50.
- KONTODIMAS D.C., LYKOURRESSIS D., KARANDINOS M.G., KATSOYANNOS P., STATHAS G.J., ELIOPoulos P.A. & L.P. ECONOMOU (2005): The effect of temperature on the development of *Nephus includens* and *Nephus bisignatus*, predators of *Planococcus citri*. — In: ERKİLÇ L. & M.B. KAYDAN (eds), Proceedings of the X International Symposium on Scale Insect Studies, held at Plant Protection Research Institute, Adana♀ Turkey, 19-23 April 2004. Adana Zirai Muscadele Arastirma Enstitusu., Adana, Turkey. 359 pp.
- KUMAR M.V.S. & N. CHAKRABORTY (1997): Biology of *Scymnus nubilus* MULSANT (Coleoptera: Coccinellidae), a native predator of pink mealy bug, *Maconellicoccus hirsutus* (GREEN) (Hemiptera: Pseudococcidae). — *Journal of Entomological Research* **21** (4): 329-334.
- LE PELLEY R.H. (1968): *Pests of coffee*. — Longmans, London. 588 pp.
- LOTFALIZADEH H. & A.A. AHMADI (2000): Natural enemies of cypress tree mealybug, *Planococcus vovae* (NASONOV), and their parasitoids in Shiraz, Iran. — *Iran Agricultural Research* **19** (2): 145-154.
- MADER L. (1955): Evidenz der paläarktischen Coccinelliden und ihrer Aberrationen in Wort und Bild. 2. — *Entomologische Arbeiten aus dem Museum G. FREY Tutzing bei Muenchen* **6**: 765-1035.
- MAJERUS M.E.N. (1994): *Ladybirds*. — London, UK: Hopper Collins. 367pp.
- MANGOUD A.A.H. (2008): Manipulation of the two predators, *Scymnus syriacus* and *Hyperaspis vingigurrae* for augmentative release for controlling the virgatus mealybug, *Ferrisia virgata*. — In: BRANCO M., FRANCO J.C. & C.J. HODGSON (eds), Proceedings of the XI International Symposium on Scale Insect Studies, Oeiras, Portugal, 24-27 September 2007. ISA Press, Lisbon, Portugal. 294 pp.
- MOSSADEGH M.S. & F. KOCHEILI (1993): Some of the ladybirds (Coccinellidae) fauna of Khuzestan, southwest, Iran. — *Scientific Journal of Agriculture, Shahid Chamran University* **16** (1-2): 23-30.
- NOVIN M. (2000): The biology and population fluctuations of *Nipaecoccus viridis* (NEWS.) and its natural enemies in citrus orchards of Dezful. — M. Sc. Thesis, Shahid Chamran University of Ahvaz. 152 pp.
- RAIMUNDO A.A.C. & A. VAN HARTEN (2000): An annotated checklist of Coccinellidae (Insecta: Coleoptera) of Yemen. — *Fauna of Arabia* **18**: 211-243.
- RAIMUNDO A.C., FÜRSCH H. & A. van HARTEN (2006): Additional notes on the ladybird beetles (Coleoptera: Coccinellidae) of Yemen with descriptions of new species. — *Fauna of Arabia* **21**: 217-245.
- RAIMUNDO A.C., FÜRSCH H. & A. van HARTEN (2008): Order Coleoptera, family Coccinellidae. — In: van HARTEN, A., (ed.), *Arthropod Fauna of the UAE*. Dar Al Ummah Printing, Adu Dhabi, UAE **1**: 217-239.

- SHARAF N.W. & D.E. MEYERDIRK (1987): A review on the biology, ecology and control of *Nipaecoccus viridis* (Homoptera: Pseudococcidae). — Miscellaneous Publications of the Entomological Society of America **66**: 1-18.
- YAGHMAEI F. & A. KHARAZI PAKDEL (1995): A faunistic survey of Coccinellids in Mashhad region. — Proceeding of the 12<sup>th</sup> Plant Protection Congress of Iran, Karj, 2-7 September 1995, Junior College of Agriculture, Karaj, Iran, 307 pp.
- YAZDANI A. (1990): The coccinellids (Col.; Coccinellidae) Fauna of Fars province. — M.Sc. Thesis, Shiraz University, 145 pp.
- ZHOU A., LU Y., ZENG L., XU Y. & G. LIANG (2012): Does Mutualism Drive the Invasion of Two Alien Species? The Case of *Solenopsis invicta* and *Phenacoccus solenopsis*. — PLoS One **7** (7): 1-10.

Authors' addresses:

Dr. Majid FALLAHZADEH  
Department of Entomology  
Jahrom Branch, Islamic Azad University  
Jahrom, Iran  
P. O. Box: 74135-355  
E-mail: mfalahm@yahoo.com

Roohafza ABDIMALEKI  
Department of Entomology  
Jahrom Branch, Islamic Azad University  
Jahrom, Iran  
E-mail: roohafzaabdimaleki@yahoo.com

Dr. Nazila SAGHAEI  
Department of Entomology  
Marvdasht Branch, Islamic Azad University  
Marvdasht, Iran  
E-mail: nazila\_saghaei@yahoo.com

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Linzer biologische Beiträge](#)

Jahr/Year: 2013

Band/Volume: [0045\\_1](#)

Autor(en)/Author(s): Fallahzadeh Majid, Abdimaleki Roohafza, Saghaei Nazila

Artikel/Article: [Contribution to the Knowledge of the Ladybird beetles \(Coleoptera, Coccinellidae\), Predators of Mealybugs \(Hemiptera, Pseudococcidae\) in Hormozgan Province, Southern Iran 673-679](#)