

Linzer biol. Beitr.	46/1	683-686	31.7.2014
---------------------	------	---------	-----------

A new species *Afrorabigus scopus* nov.sp. from Tanzania (Coleoptera: Staphylinidae: Philonthina)

L. HROMÁDKA

A b s t r a c t : *Afrorabigus scopus* nov. sp. is described from Tanzania and diagnosed from similar *A. uromastyx* HROMÁDKA 2011. The male genitalia and important external male characters of both species are illustrated.

K e y w o r d s : Coleoptera, Staphylinidae, Philonthina, *Afrorabigus*, Afrotropical region, Tanzania, taxonomy, new species.

Introduction

The endemic Afrotropical philonthine genus *Afrorabigus* was described by Levasseur (1965). At this time, it is known nine species of this genus. The representatives of the genus *Afrorabigus* are similar in habitus to those of *Rabigus* MULSANT & REY 1876, but differ from them by the relative lengths of maxillary palpomeres: palpomere 3 is longer than palpomere 2.

Material, methods and measurements

The material referred in this study is deposited in the following institutions and private collection:

BMNH The Natural History Museum, London, United Kingdom (Maxwell Barclay, Roger Booth)

cHPC author's private collection

NMPC National Museum, Praha, Czech Republic (Jiří Hájek)

A double slash (//) is used to divide separate labels of the type specimen. All measurements were taken from the beetles with extended abdomen. Ratios mentioned in the description can be converted to length using the following formula: 20 units = 1 mm.

Results

Afrorabigus scopus nov.sp. (Figs 1-3)

Type material: Holotype ♂: Tanzania, Tango, viii.1902, Methner //Holotype *Afrorabigus scopus* nov. sp. Hromádka det., 2013 [red oblong printed label] (NMPC). Paratype ♀: same label data as holotype (cHPC).

Description: Body length 4.8 mm, length of fore body (to end of elytra) 2.4 mm.

Colouration: Head black, anterior half of pronotum orange, posterior half blackish, scutellum black, elytra orange, sides and suture narrowly blackish, sometimes shoulders and around scutellum blackish, abdomen visible tergites 1-4 black, posteriot margin of each tergite narrowly and paratergites orange, visible tergites 5-6 in anterior third orange-yellow, posterior two thirds darker. Maxillary and labial palpi, antennomeres 1-2 and base of antennomere 3 dirty yellow, legs yellow.

Head wider than long (ratio 18 : 15), distinctly narrowed posteriad. Posterior angles obtusely rounded, bearing one long black bristle. Four punctures present between eyes, distance between medial punctures four times as large as distance between medial and lateral puncture. Eyes slightly convex, longer than temples (ratio 8 : 5), posterior margin with two coarse punctures, temporal area with several varying large punctures. Surface with traces of very fine microsculpture in places.

Antennae long, slightly widened distally, exceeding posterior margin of pronotum by the lenght of antennomeres 10-11 when reclined. Antennomeres 1-6 and 11 distinctly longer than wide, antennomeres 7-10 slightly longer than wide. Antennomere 1 almost twice longer than antennomere 11, antennomere 2 as long as antennomere 3.

Pronotum slightly longer than wide (ratio 21 : 18), parallel-sided, anterior angles conspicuously deflexed, vaguely obtusely rounded, posterior angles markedly rounded. Each dorsal row with six aproximately equidistant, punctures,each sublateral row with two punctures, puncture two distinctly shifted laterally. Surface with microsculpture similar to that on head.

Scutellum very finely punctured, diameter of punctures approximately as large as eye-facets, separated by distance one puncture diameter or slightly larger. Surface with eye like microsculpture; setation dark.

Elytra as long as wide, slightly widened posteriad. Punctuation very fine and very dense, diameter of punctures slightly larger than eye-facets, separated by distance smaller than one puncture diameter. Surface without microsculpture; setation greyish.

Legs. Metatibia approximately as long as metatarsus, metatarsomere 1 longer than metatarsomere 5, slightly shorter than metatarsomeres 2-4 combined.

Abdomen wide, almost parallel-sided, first three visible tergites with two basal lines, elevated area between lines densely and finely punctate. Punctuation at base of tergites similar to that on elytra, becoming sparser towards posterior margin of each tergite. Surface without microsculpture; setation similar to that on elytra.


Male. Protarsomeres 1-3 dilated and sub-bilobed, each covered with modified pale setae ventrally, protarsomere 4 narrower than preceding ones. Sternite IX (Fig. 3), aedeagus (1-2).

Female. Protarsomeres 1-3 less dilated than in male, each covered with modified pale setae venrally, protarsomere 4 narrower than preceding ones.

C o m p a r a t i v e n o t e s : *Afrorabigus scopus* nov.sp. it differs from all species of this genus by having 6 punctures in dorsal rows of pronotum, (the other species have 4 or 5 punctures in dorsal rows).

E t y m o l o g y : The name of this species, a noun in apposition, is the Latin generic name of African Hamerkop *Scopus umbretta* GMELIN 1789.

D i s t r i b u t i o n : Tanzania.



Figs 1-3: *Afrorabigus scopus* nov.sp. (1) aedeagus, ventral view, (2) aedeagus, lateral view, (3) male sternite IX, ventral view. **Figs 4-6:** *Afrorabigus tropheus* HROMÁDKA 2011. (4) aedeagus, ventral view, (5) aedeagus, lateral view, (6) male sternite IX, ventral view.

Afrorabigus tropheus HROMÁDKA 2011 (Figs 4-6)

M a t e r i a l e x a m i n e d : Holotype ♂: Ethiopia: Bale, S of Omar: 07.07°N, 40.60°E, 1200 m 0704N4036E. CHPC).

Acknowledgements

I wish to express my thanks to Jiří Janák (Rtyň nad Bílinou, Czech Republic) for help in preparing the manuscript.

Zusammenfassung

Afrorabigus scopus wird von Tanzania beschrieben und von der ähnlichen *A. tropheus* HROMÁDKA 2011 unterschieden. Merkmale insbesondere des männlichen Geschlechts beider Arten werden abgebildet.

References

- BERNHAUER M. (1915): Zur Staphyliniden-Fauna des tropischen Afrika. — Annales Musei Nationalis Hungarici **13**: 95-189.
- HERMAN L.H. (2001): Catalog of the Staphylinidae (Insecta: Coleoptera), 1758 to the end of the second Millennium. V. Staphylinine group (Part 2): Diochini, Maorothiini, Othiini, Platyprosopini, Staphylinini. (Amblyopinina, Anisolinina, Hyptiomima, Philonthina) — Bulletin of the American Museum of Natural History **265**: 2674-2694.
- HROMÁDKA L. (2011): Revision of the Afrotropical species of the genus *Afrorabigus* (Coleoptera: Staphylinidae: Philonthina). — Journal of the National Museum (Prague), Natural History Series **180** (6): 63-79.
- LEVASSEUR L. (1965): *Afrorabigus*, gen.nov. descriptions d'espèces nouvelles du Katanga Gabon (Col., Staphylinidae, Philonthini). — Annales da la Société Entomologique de France (n.s.) **1** (1) 57-62.
- LEVASSEUR L. (1967): Nouvelles espèces de Staphylinidae de Côte d'Ivoire, du Sudan et du Katanga. — Bulletin de l'Institut Fondamental d'Afrique Noire (A) **29**: 1624-1644.
- TOTTENHAM C.E. (1962): Mission zoologique de l'I.R.S.A.C. en Afrique orientale (P. Basilewsky et N. Leleup, 1957). LXXXVI. Coleoptera, Staphylinidae, Staphilininae. — Annales du Musée de l'Afrique Centrale, Sciences Zoologique **(8)** 110: 132-258.

Author's address:

Lubomír HROMÁDKA
Anny Letenské 7
CZ-120 00 Praha 2, Czech Republic
E-mail: hromadka@seznam.cz

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Linzer biologische Beiträge](#)

Jahr/Year: 2014

Band/Volume: [0046_1](#)

Autor(en)/Author(s): Hromadka Lubomir

Artikel/Article: [A new species Afrorabigus scopus nov.sp. from Tanzania \(Coleoptera: Staphylinidae: Philonthina 683-686](#)