

Linzer biol. Beitr.	47/1	449-458	31.7.2015
---------------------	------	---------	-----------

A faunistic study on Scelionidae (Hymenoptera: Platygastroidea) from some regions of Iran

Hassan GHAHARI & Peter Neerup BUHL

A b s t r a c t : The Scelionidae is a large family of parasitic Hymenoptera whose members specialize in egg parasitism of insects and arachnids. This paper deals with the faunistic data of scelionid wasps (Hymenoptera: Scelionidae) from some regions of Iran. In total 11 species from 4 genera, *Gryon* HALIDAY, *Scelio* LATREILLE, *Telenomus* HALIDAY, *Trissolcus* ASHMEAD and 2 subfamilies Scelioninae and Telenominae were collected and identified.

K e y w o r d s : Hymenoptera, Scelionidae, fauna, Iran.

Introduction

The Scelionidae are exclusively egg parasitoids, with a wide host array, with host-dependent morphological specialization. They show high specificity in host partitioning at tribal level (GALLOWAY & AUSTIN 1983). There are 3.308 valid species of Scelionidae (JOHNSON 2010). Scelionids have been used quite successfully in classic biological control programs directed principally against agricultural pest. Scelionids are virtually all solitary idiobiont primary parasitoids, and reports that diverge from this biology are rare and often unsubstantiated (EBERHARD 1975; GALLOWAY & AUSTIN 1984; DANGERFIELD et al. 2001). Many of the characteristics considered most desirable in a natural enemy can be found within scelionid wasps. They have high searching abilities and reproductive rates, synchrony with host populations, positive host-density responsiveness, simple adult diets, can be reared easily, and lack of hyperparasitoids (DANGERFIELD et al. 2001; SAMIN et al. 2011b).

Of the three scelionid subfamilies, the Telenominae, the Scelioninae and the Teleasinae, the first two are highly abundant, diverse and quite significant to agroecosystems. While the Lepidoptera and the Heteroptera are principal host groups for telenomines, the Orthoptera, the Heteroptera, the Coleoptera, the Odonata, the Mantodea and the araenomorph spiders form the major host assemblage of the Scelioninae (AUSTIN et al. 2005).

The fauna of Iranian Scelionidae was studied well by different researchers (RADJABI & AMIR NAZARI 1989; RADJABI 1994; NOORBAKHSH & RAZAVI 1995; IRANIPOUR et al. 1998; TAGHADDOSI & RAJABI 1998; NOORI & ASGARI 2000; MANSOUR GHAZI & RADJABI 2000; HASHEMI RAD et al. 2000, 2002; RADJABI 2001; HAGHSHENAS 2004;

HASHEMI RAD 2008; RAKHSHANI et al. 2008; SAMIN 2010; IRANPOUR & JOHNSON 2010; FATHI et al. 2011; GHAHARI et al. 2009; HEJAZI et al. 2011; SHAFAEI et al. 2011; RAHNEMAYE-SHASAVARI et al. 2011; SAMIN et al. 2010a, b, c; 2011 a, b, c; 2012; SAMIN & ASGARI 2012a, b). Additionally some checklists were prepared on Iranian Scelionidae e.g. MODARRES AWAL 1997, SCELIONIDAE; SAMIN et al. 2010d, *Telenomus*; GHAHARI et al. 2011, *Trissolcus*). Recently, GHAHARI et al. (2015) catalogued all the data about Iranian Scelionidae which totally fifty four species have been recorded from Iran so far.

Materials and Methods

The materials of this research were collected from some regions of Iran by rearing of some eggs of Pentatomidae and Scutelleridae (Hemiptera), *Locusta migratoria* (LINNAEUS) (Orthoptera: Acrididae), and *Sesamia cretica* LEDERER (Lepidoptera: Noctuidae) for emergence of parasitoids inside the host. Egg masses were placed in plastic bags and in optimum condition ($26\pm2^{\circ}\text{C}$, $65\pm5\%$ RH, 14: 10 L: D) in incubator. Classification, nomenclature and distributional data of Scelionidae suggested by KOZLOV (1978), KOZLOV & KONOVOVA (1983), KONOVOVA (1992, 1995), JOHNSON (1992, 2010), KONOVOVA & KOZLOV (2008) have been followed.

Results

Totally eleven scelionid species from four genera, *Gryon* HALIDAY, *Scelio* LATREILLE, *Telenomus* HALIDAY, *Trissolcus* ASHMEAD and two subfamilies, Scelioninae and Telenominae are given in this paper. The list of species is given below alphabetically with distribution data.

Family Scelionidae (HALIDAY, 1840)

Subfamily Scelioninae FÖRSTER, 1856

Genus *Gryon* HALIDAY, 1833

***Gryon monspeliense* (PICARD, 1924)**

Hadronotus monspeliensis PICARD, 1924; *Hadronotus afanasievi* MEIER, 1949; *Hadronotus telengai* RJACHOVSKY, 1959; *Gryon afanasievi* KOZLOV, 1963; *Hadronotellus monspeliensis* SZABO, 1966.

M a t e r i a l e x a m i n e d : Lorestan province, Azna, $33^{\circ}28'N$ $49^{\circ}25'E$, 2♀♀, ex *Dolycoris baccarum* LINNAEUS, 14.v.2012.

G e n e r a l d i s t r i b u t i o n : France (PICARD 1924), Turkey (LODOS 1982), Russia, Ukraine, Azerbaijan, Belgium, Bulgaria, Italy, Romania (KONOVOVA & KOZLOV 2008).

Genus *Scelio* LATREILLE, 1805

Scelio flavibarbis (MARSHALL, 1874)

Aleria flavibarbis MARSHAL, 1874.

M a t e r i a l e x a m i n e d : Hamadan province, Malayer, 34°20'N 48°45'E, 3♀ ♀, ex *Locusta migratoria* (LINNAEUS) (Orthoptera: Acrididae), 11.vi.2011.

G e n e r a l d i s t r i b u t i o n : Corsica (MARSHAL 1874), Bulgaria, Kazakhstan, Russia Ukraine, (KONONOVA & KOZLOV 2008).

Subfamily T e l e n o m i n a e THOMSON, 1860

Genus *Telenomus* HALIDAY, 1833

Telenomus busseolae GAHAN, 1922

Platytenomus hylas NIXON, 1935; *Platytenomus busseolae* MASNER, 1965.

M a t e r i a l e x a m i n e d : Golestan province, Gorgan, 36°50'N 54°30'E, 2♀ ♀, 1♂, ex *Sesamia cretica* LEDERER (Lepidoptera: Noctuidae), 21.vii.2013.

G e n e r a l d i s t r i b u t i o n : Israel (KOZLOV & KONONOVA 1983), Bangladesh, Cameroon, Egypt, Ghana, Greece, Iran, Iraq, Kenya, Mauritius, Nigeria, Reunion, Senegal, South Africa, Sudan, Uganda (POLASZEK & KIMANI 1990).

Telenomus chloropus (THOMSON, 1861)

Phanurus chloropus THOMSON, 1861; *Telenomus sokolowi* MAYR, 1897; *Telenomus mayri* SOKOLOV, 1904; *Prophanurus Sokolowi* KIEFFER, 1912; *Telenomus tischleri* NIXON, 1939; *Telenomus sokolovi* MEIER, 1940.

M a t e r i a l e x a m i n e d : Tehran province, Taleqan, 35.48°N 50.58°E, 3♀ ♀, ex *Eurygaster integriceps* PUTON, 9.viii.2012.

G e n e r a l d i s t r i b u t i o n : Ukraine (KIEFFER 1926), Turkey (LODOS 1961), England (JAVAHERY 1968), Armenia, Azerbaijan, Georgia, Kazakhstan, Moldova, Russia (European part, Altai, Far East) (KOZLOV & KONONOVA 1983; KONONOVA 1995), Ireland (O'CONNOR & MINEO 2009), France, Hungary, Spain, Sweden, Japan, USA (Mississippi) (JOHNSON 1984).

Genus *Trissolcus* ASHMEAD, 1893

Trissolcus basalis (WOLLASTON, 1858)

Telenomus basalis WOLLASTON, 1858; *Telenomus maderensis* WOLLASTON, 1858; *Telenomus megacephalus* ASHMEAD, 1895; *Telenomus megalcephalus* SCHULZ, 1906; *Telenomus piceipes* DODD, 1920; *Liophanurus megacephalus* KIEFFER, 1926; *Microphanurus basalis* NIXON, 1935; *Asolcus basalis* DELUCCHI, 1961; *Trissolcus basalis* MASNER, 1965; *Trissolcus maderensis* MASNER, 1965; *Trissolcus piceipes* MASNER, 1965; *Asolcus lodosi* SZABO, 1981.

M a t e r i a l e x a m i n e d : Qazvin province, Abyek, 36°03'N 50°20'E, 3♀ ♀, 1♂, ex *Eurygaster maura* (LINNAEUS), 14.vii.2011. West Azarbaijan province, Salmas, 38°11'N 44°44'E, 2♀ ♀, 2♂ ♂, ex *Eurygaster integriceps* PUTON, 24.ix.2012. Kordestan province, Qorveh, 35°15'N 47°40'E, 4♀ ♀, 3♂ ♂, ex *Aelia acuminata* (LINNAEUS), 27.vi.2013.

General distribution: Afro-tropical, Australasian, Nearctic, Neotropical, Oriental, and Palaearctic Regions (JOHNSON 1992).

***Trissolcus djadetshko* (RJACHOVSKY, 1959)**

Microphanurus djadetshko RJACHOVSKY, 1959: *Asolcus djadetshko* VIKTOROV, 1964.

Material examined: Kordestan province, Bijar, 35°52'N 47°36'E, 1♀, ex *Eurygaster integriceps* PUTON, 29.vi.2013.

General distribution: Armenia, Azerbaijan, Kazakhstan, Moldova, Russia (European part, Altai, Primorskii krai), Ukraine, Uzbekistan, Mongolia (KOZLOV & LE 1988; KONOVOVA 1995), Denmark (BUHL 1999), Turkey (KOÇAK & KILINÇER 2000, 2003), Romania (POPOVICI 2004).

***Trissolcus festivae* (VIKTOROV, 1964)**

Asolcus festivae VIKTOROV, 1964.

Material examined: Zanjan province, Zanjan, 36°35'N 48°15'E, 2♀, ex *Eurygaster integriceps* PUTON, 16.viii.2012. Chaharmahal & Bakhtiari province, Boroojen, 31°54'N 51°12'E, 1♀, 1♂, ex *Eurydema ornatum* LINNAEUS, 7.ix.2013.

General distribution: Armenia, Azerbaijan, Kazakhstan, Moldova, Mongolia, Russia (North Caucasus, European part, Altai), Turkmenistan, Ukraine, Uzbekistan (KOZLOV & LE 1988; KOZLOV & KONOVOVA 1983), Romania (FABRITIUS 1974), Turkey (TARLA 1997).

***Trissolcus grandis* (THOMSON, 1861)**

Telenomus grandis THOMSON, 1861; *Telenomus nigripes* THOMSON, 1861; *Telenomus frontalis* THOMSON, 1861; *Telenomus nigrita* THOMSON, 1861; *Telenomus nigritus* DALLA TORE, 1898; *Aphanurus nigripes* KIEFFER, 1912; *Aphanurus grandis* KIEFFER, 1912; *Aphanurus nigrita* KIEFFER, 1912; *Aphanurus frontalis* KIEFFER, 1912; *Microphanurus nigripes* KIEFFER, 1926; *Microphanurus grandis* KIEFFER, 1926; *Microphanurus nigritus* KIEFFER, 1926; *Microphanurus frontalis* KIEFFER, 1926; *Asolcus grandis* DELUCCHI, 1961; *Asolcus nixomartini* JAVAHERY, 1968; *Asolcus silwoodensis* JAVAHERY, 1968; *Trissolcus nigripes* FERGUSSON, 1978; *Trissolcus nixomartini* FERGUSSON, 1978; *Trissolcus silwoodensis* FERGUSSON, 1978; *Telenomus nigripes* FERGUSSON, 1984; *Telenomus nixomartini* FERGUSSON, 1984; *Telenomus silwoodensis* FERGUSSON, 1984.

Material examined: Yazd province, Yazd, 31°53'N 54°27'E, 3♀, 1♂, ex *Eurygaster maura* (LINNAEUS), 20.iv.2012. Chaharmahal & Bakhtiari province, Koohrang, 32°25'N 50°00'E, 2♀, 3♂, ex *Dolycoris baccarum* (LINNAEUS), 6.ix.2013. Markazi province, Khomeyn, 33°40'N 50°00'E, 2♀, 4♂, ex *Graphosoma lineatum* (LINNAEUS), 15.v.2014.

General distribution: Denmark (THOMSON 1861), Armenia, Kazakhstan, Middle Asia, Moldova, Russia (South of Far East), Ukraine (KOZLOV & KONOVOVA 1983; KONOVOVA 1995), Belgium (DEBAUCHE 1947), Syria (REMAUDIÈRE & SKAF 1963), Morocco (VOEGELÉ 1964), England (JAVAHERY 1968), Romania (FABRITIUS 1974), Italy (VIGGIANI & MINEO 1974), Turkey (KOÇAK 2007), Iraq (ALI 2011).

***Trissolcus rufiventris* (MAYR, 1908)**

Telenomus rufiventris MAYR, 1907; *Prophanurus rufiventris* KIEFFER, 1912; *Dissolcus rufiventris* KIEFFER, 1926; *Microphanurus anitus* NIXON, 1939; *Asolcus rufiventris* MASNER, 1959.

M a t e r i a l e x a m i n e d : South Khorasan province, Sarbisheh, 32°24'N 60°05'E, 2♀♀, 1♂, ex *Aelia furcula* FIEBER, 19.iv.2013. Kermanshah province, Sonqor, 34°50'N 47°30'E, 2♀♀, ex *Eurygaster integriceps* PUTON, 11.viii.2014.

G e n e r a l d i s t r i b u t i o n : Morocco (VOEGELÉ 1964), Romania (FABRITIUS 1974), Middle Asia, Moldova, Mongolia, West Kazakhstan, West Europe, Russia, Ukraine (North Caucasus, European part, Altai), Africa (KOZLOV & KONOVOVA 1983; KONOVOVA 1995), Turkey (TUATAY et al. 1972; KOÇAK 2007), Iraq (ALI 2011).

***Trissolcus semistriatus* (NEES, 1834)**

Teleas semistriatus NEES, 1834; *Telenomus ovulorum* THOMSON, 1861; *Telenomus semistriatus* MAYR, 1879; *Aphanurus semistriatus* KIEFFER, 1912; *Microphanurus semistriatus* KIEFFER, 1926; *Microphanurus alexeevi* MEIER, 1949; *Microphanurus schtepelnikovae* MEIER, 1949; *Asolcus semistriatus* MASNER, 1959.

M a t e r i a l e x a m i n e d : Isfahan province, Kashan, 34°00'N 51°20'E, 2♀♀, 3♂♂, ex *Carpocoris fuscipinus* (BOHEMAN), 5.vi.2011. Semnan province, Garmsar, 35°00'N 52°20'E, 4♀♀, 2♂♂, ex *Dolycoris baccarum* LINNAEUS, 25.iv.2012. Golestan province, Azadshahr, 37°07'30"N 55°06'00"E, 2♀♀, 2♂♂, ex *Eurygaster maura* (LINNAEUS), 21.vii.2013.

G e n e r a l d i s t r i b u t i o n : Armenia, Austria, Azerbaijan, Denmark, France, Germany, Moldova, Russia (European part, Altai, Primorskii krai), Ukraine (KOZLOV & KONOVOVA 1983; KONOVOVA 1995), England (JAVAHERY 1968), Portugal (GRAHAM 1984), Romania (FABRITIUS 1974), Turkey (LODOS 1961), Morocco (VOEGELÉ 1964), Iraq (ALI 2011).

***Trissolcus vassilievi* (MAYR, 1903)**

Telenomus Vassiliewi MAYR, 1903; *Microphanurus vassiliewi* KIEFFER, 1926; *Microphanurus vassilievi* MEIER, 1940; *Trissolcus (Microphanurus) vassilievi* RJACHOVSKY, 1959; *Asolcus vassilievi* DELUCCHI, 1961.

M a t e r i a l e x a m i n e d : East Azarbaijan province, Shabestar, 38°12'N 45°44'E, 4♀♀, 1♂, ex *Carpocoris mediterraneus* (TAMANINI), 17.vi.2013. North Khorasan province, Shirvan, 37°32'N 57°54'E, 3♀♀, ex *Graphosoma lineatum* (LINNAEUS), 28.viii.2014.

G e n e r a l d i s t r i b u t i o n : Turkey (LODOS 1961; KOZLOV & KONOVOVA 1983; KONOVOVA 1995), Syria (REMAUDIÈRE & SKAF 1963), Morocco (VOEGELÉ 1964), Armenia, Moldova, Russia (European part, North Caucasus, European part), Central Asia, Mongolia, Ukraine (KOZLOV & KONOVOVA 1983; KONOVOVA 1995), Iraq (ALI 2011).

Discussion

The several conducted researches on Iranian Scelionidae which were catalogued by GHAHARI et al. (2015) indicate that the fauna of these beneficial insects is very diverse in Iran. Scelionid wasps can have efficient role in biological control programs if the conservation methods are done effectively in agroecosystems. The main conservation of these efficient egg parasitoids can be done with management in pesticides application. Although fewer than 30 species have been used in classical biological control attempts, several of these have produced excellent results. In despite of rich faunistic works on Iranian Scelionidae but since Iran contains various geographical regions and especially diverse fauna of Pentatomidea (GHAHARI et al. 2009, 2014) as the main hosts of

scelionid wasps, we expect that there are some other species which carrying on the faunistic syrveys will result to finding new data.

Acknowledgments

The authors are grateful to E. Kocak (Turkey) for identification of materials, L. Masner (Canada) and N. Samin (Iran) for providing the necessary articles, F. Gusenleitner (Austria) for editing and providing the manuscript. The research was supported by Shahre Rey Islamic Azad University.

Zusammenfassung

Vorliegende Arbeit behandelt die Faunistik der Scelionidae (Hymenoptera) aus verschiedenen Regionen Irans. Insgesamt gelang der Nachweise von 11 Arten aus den 4 Gattungen *Gryon HALIDAY*, *Scelio LATREILLE*, *Telenomus HALIDAY* und *Trissolcus ASHMEAD*, welche den Unterfamilien Scelioninae and Telenominae angehören.

References

- ALI W.Kh. (2011): The level of sunn pest oophagous parasitoids (Hymenoptera: Scelionidae) in infested wheat field of northern Governorate in Iraq with an identification key of *Trissolcus* species. — Bulletin of the Iraq Natural History Museum **11** (4): 7-15.
- AUSTIN A.D., JOHNSON N.F. & M. DOWNTON (2005): Systematics, evolution and biology of Scelionid and Platygastrid wasps. — Annual Review of Entomology **50**: 553-582.
- BUHL P.N. (1999): Tillaeg til fortægnelsen over Danmarks proctotruper (Hymenoptera: Proctotropoidea s.l.). 2. del: Scelionidae, Platygastridae, Megaspilidae og Ceraphronidae. — Flora og Fauna **105** (1): 9-13.
- DANGERFIELD P., AUSTIN A. & G. BAKER (2001): Biology, ecology and systematics of Australian *Scelio* (Hymenoptera: Scelionidae). — Melbourne: CSIRO, 254 pp.
- DEBAUCHE H.R. (1947): Scelionidae de la faune belge (Hymenoptera Parasitica). — Bulletin et Annales de la Societe Entomologique de Belgique **83**: 255-285.
- EBERHARD W.G. (1975): The ecology and behavior of a subsocial pentatomid bug and two scelionid wasps: strategy and counter strategy in a host and its parasites. — Smithsonian Contribution Zoology **205**: 1-39.
- FABRITIUS K. (1974): Die Telenominen (Hymenoptera: Scelionidae) Rumäniens, eine faunistische Studie in unmittelbarer Verbindung mit der biologischen Schädlingsbekämpfung. — Folia Entomologica Hungarica **27** (Supplement): 339-344.
- FATHI A.A., NOURI GHANBALANI Gh. & P. HONARMAND (2011): Parasitoids of sunn pest (*Eurygaster integriceps* PUTON) and percent parasitism of egg in Quraim, Ardabil province. — The Scientific Journal of Agriculture (Plant Protection) **34** (1): 47-57.
- GALLOWAY I.D. & A.D. AUSTIN (1983): Revision of the Scelioninae (Hymenoptera: Scelionidae) in Australia. — Australian Journal of Zoology Supplementary Series **99**: 1-138.
- GHAHARI H., HAVASKARY M., TABARI M., OSTOVAN H., SAKENIN H. & A. SATAR (2009): An annotated catalogue of Orthoptera (Insecta) and their natural enemies from Iranian rice fields and surrounding grasslands. — Linzer biologische Beiträge **41** (1): 639-672.
- GHAHARI H., CHEROT F., LINNAVUORI R.E. & H. OSTOVAN (2009): Annotated catalogue of Iranian burrower bugs (Heteroptera, Pentatomidae, Cydnidae). — ZooKeys **26**: 1-31.

- GHAHARI H., BUHL P.N. & E. KOCAK (2011): Checklist of Iranian *Trissolcus* ASHMEAD (Hymenoptera: Platygastroidea: Scelionidae: Telenominae). — International Journal of Environmental Studies, October 2011: 1-9.
- GHAHARI H., MOULET P. & D.A. RIDER (2014): An annotated catalog of the Iranian Pentatomoidae (Hemiptera: Pentatomomorpha). — Zootaxa **3837** (1): 1-95.
- GHAHARI H., BUHL P.N., KOCAK E. & Sh. IRANPOUR (2015): An annotated catalogue of the Iranian Scelionidae (Hymenoptera: Platygastroidea). — Entomofauna **36**: 349-376.
- GRAHAM M.W.R. DE V. (1984): Madeira insects, mainly Hymenoptera Proctotrupoidea, Ceraphronoidea, and Bethyloidea. — Boletim do Museu Municipal do Funchal **36**: 83-110.
- HAGHSHENAS A. (2004): Protection and conservation of sunn pest tachinid ad scelionid parasitoids in Chahar Mahal va Bakhtiari province. — Proceedings of 16th Iranian Plant Protection Congress, University of Tabriz, p. 5.
- HASHEMI RAD H. (2008): Study on egg parasitoids *Trissolcus agriope* and *Psix* sp. using egg of five bug species in the laboratory condition. — Proceedings of 18th Iranian Plant Protection Congress, University of Bu-Ali Sina Hamedan, p. 47.
- HASHEMI RAD H., SOLEYMAN NEJADIAN E., RADJABI Gh. & Sh. IRANPOUR (2000): Introduction of egg-parasitoids of pentatomid pistachio bugs in Kerman province. — Proceedings of 14th Iranian Plant Protection Congress, Isfahan University of Technology, p. 269.
- HASHEMI RAD H., SOLEYMAN NEJADIAN E. & Gh. RADJABI (2002): Population fluctuation of the pistachio green stink bugs (*Brachynema* spp. & *Acrosternum* spp.) and their egg parasitoids in Rafsanjan. — Proceedings of 14th Iranian Plant Protection Congress, Isfahan University of Technology, p. 107.
- HEJAZI M.J., KARIMZADEH R. & S. IRANPOUR (2011): *Eurygaster integriceps* egg parasitoids in Gharamalek. — Proceedings of Global conference on Entomology, 5-9 March 2011, Chiang Mai, Thailand, p. 45.
- IRANPOUR SH., KHARRAZI PAKDEL A., ESMAILI M. & Gh. RAJABI (1998): Introduction of two species of egg parasitoids of pentatomid bugs from genus *Trissolcus* (Hym.: Scelionidae). — Proceedings of 13th Iranian Plant Protection Congress, Campus of Agriculture and Natural Resources, University of Tehran, Karaj, p. 4.
- IRANPOUR SH. & N.F. JOHNSON (2010): *Trissolcus radjabii* n.sp. (Hymenoptera: Platygastridae), an egg parasitoid of the shield bug, *Apodiphus amygdali* (Heteroptera: Pentatomidae) and the sunn pest, *Eurygaster integriceps* (Heteroptera: Scutelleridae). — Zootaxa **2515**: 65-68.
- JAVAHERY M. (1968): The egg parasite complex of British Pentatomoida (Hemiptera): taxonomy of Telenominae (Hymenoptera: Scelionidae). — Transactions of the Royal Entomological Society of London **120**: 417-436.
- JOHNSON N.F. (1984): Systematics of Nearctic *Telenomus*: classification and revisions of the podisi and phymatae species groups (Hymenoptera: Scelionidae). — Bulletin of the Ohio Biological Survey **6** (3): 1-113.
- JOHNSON N.F. (1992): Catalog of world Proctotruipoidea excluding Platygastridae. Memoirs of the American Entomological Institute **51**: 1-825.
- JOHNSON N.F. (2010): Fauna Europaea: Scelionidae. — In: NOYES J. (ed.), Fauna Europaea: Hymenoptera. Fauna Europaea version 2.2, <http://www.faunaeur.org> (accessed 12 October 2010).
- KIEFFER J.J. (1926): Scelionidae. — Das Tierreich. Vol. **48**. Walter de Gruyter & Co., Berlin, 885 pp.

- KOÇAK E. (2007): Egg parasitoids of sunn pest in Turkey: A review, pp. 225-235. — In: PARKER B.L., SKINNER M., BOUHSSINI M.E. & KUMARI S.G. (eds), Proceedings of Second International Conference on Sunn Pest (19-22 July 2004, Aleppo, Syria). Sunn pest management, a decade of progress, 1994-2004. The Arab Society for Plant Protection, Beirut, Lebanon, 432 pp.
- KOÇAK E. & N. KILINÇER (2000): *Trissolcus* species (Scelionidae, Hymenoptera) new records for the beneficial fauna of Turkey. — Plant Protection Bulletin **40** (3-4): 169-177.
- KOÇAK E. & N. KILINÇER (2003): Taxonomic Studies on *Trissolcus* sp. (Hymenoptera: Scelionidae), Egg Parasitoids of Sunn Pest (Hemiptera: Scutelleridae: *Eurygaster* sp.), in Turkey. — Turkish Journal of Zoology **27** (4): 301-317.
- KONONOVA S.V. (1992): Fauna of Ukraine. Volume 11. Parasitic Hymenoptera. Part 10. Proctotrupoid scelionid parasitoids of the subfamilies Scelioninae and Telenominae. — Naukova Dumka, Kiev. 255 pp.
- KONONOVA S.V. (1995): Family Scelionidae, pp. 57-121. — In: LEHR P.A. (ed.), Key to insects of Russian Far East in six volume. vol. 4. Neuropteroidea, Mecoptera, Hymenoptera. Part 2. Hymenoptera. — Dal'nauka, Vladivostok, 600 pp.
- KONONOVA S.V. & M.A. KOZLOV (2008): [Scelionids of the Palearctic (Hymenoptera, Scelionidae). Subfamily Scelioninae]. — Tovarishchestvo Nauchnykh Izdanii KMK, Saint Petersburg, 489 pp.
- KOZLOV M.A. (1978): [Superfamily Proctotropoidea]. Pages 538-664. — In: MEDVEDEV G.S., [Determination of insects of the European portion of the USSR.] Vol. 3, part 2. — Nauka, Leningrad, 758 pp.
- KOZLOV M.A. & S.V. KONONOVA (1983): Telenominae of the fauna of the USSR (Hymenoptera, Scelionidae, Telenominae). — Leningrad Nauka Publisher **136**, 336 pp.
- KOZLOV M.A. & X.H. LEE (1988): *Trissolcus*, pp. 1110-1179. — In: MEDVEDEV G.S. (ed.), Keys to the Insects of the European Part of the USSR. III. Hymenoptera. Part II. Leningrad: Nauka Publisher, 336 pp.
- LODOS N. (1961): Investigations on sunn pest (*Eurygaster integriceps* PUT.) in Turkey, Iran and Syria. (Distributions, damages, biology, parasites and control). — Aegean University Agricultural Faculty Publications **51**, 115 pp. (in Turkish).
- LODOS N. (1982): Turkish Entomology Volume II (Genel, Uygulamalı ve Faunistik). — Aegean University Agricultural Faculty Publications **429**, Izmir, 580 pp. (in Turkish).
- MANSOUR GHAZI M. & G.R. RADJABI (2000): Sunn pest tachinid and scelionid parasitoids in Kurdistan. — Proceedings of 14th Iranian Plant Protection Congress, Isfahan University of Technology, p. 219.
- MARSHALL T.A. (1874): Descriptions of a new genus and two new species of European *Oxyura*. — Entomologists Monthly Magazine **10**: 207-209.
- MODARRES AWAL M. (1997): Family Scelionidae (Hymenoptera), pp. 279-280. — In: MODARRES AWAL M. (ed.), List of agricultural pests and their natural enemies in Iran. Ferdowsi University Press, 429 pp.
- NOORBakhsh S.H. & S. RAZAVI (1995): Distribution of sunn pest (*Eurygaster integriceps* PUT.) and its natural enemies in Chaharmahal Bakhtiari. — Proceedings of 12th Iranian Plant Protection Congress, University of Tehran, p. 15.
- NOORI H. & Sh. ASGARI (2000): Study and identification of sunn pest egg parasitoid in Qazvin Province. — Proceedings of 14th Iranian Plant Protection Congress, Isfahan University of Technology, p. 218.
- O'CONNOR J. P. & G. MINEO (2009): *Telenomus chloropus* (THOMSON) new to Ireland and second Irish records of *T. nitidulus* (THOMSON) and *Trissolcus flavipes* (THOMSON) (Hym., Scelionidae). — Entomologists Monthly Magazine **145**: 106.
- PICARD F. (1924): Description d'un nouveau Proctotrypide du genre *Hadronotus* [Hym.]. — Bulletin de la Societe Entomologique de France **1924**: 107-109.

- POLASZEK A. & S.W. KIMANI (1990): *Telenomus* species (Hymenoptera: Scelionidae) attacking eggs of pyralid pests (Lepidoptera) in Africa: a review and guide to identification. — Bulletin of Entomological Research **80**: 57-71.
- POPOVICI O. (2004): New Scelionidae species (Hymenoptera: Platygastroidea: Scelionidae) for Romania's fauna. — Analele Științifice ale Universității "Al. I. Cuza" Iași, s. Biologie animală, Tom L: 103-106.
- RADJABI Gh. & M. AMIR NAZARI (1989): Egg parasites of sunn pest in the central part of Iranian plateau. — Entomologie et Phytopathologie Appliquées **56** (1-2): 1-12.
- RADJABI Gh. (1994): First report of the existence of sunn pest egg parasitoid, *Trissolcus festivae* VIKTOROV (Hym.: Scelionidae) in Iran and some preliminary studies on its biology. — Journal of Entomological Society of Iran **14**: 1-7.
- RADJABI Gh. (2001): *Trissolcus esmailii* sp.n. (Hym.: Scelionidae), an egg parasitoid of *Eurygaster integriceps* PUT. and *Dolycoris baccarum* L. in Iran. — Journal of Agricultural Science **7** (2) 106-112 (in Persian, English summary).
- RAHNEMAYE-SHASAVARI M., LOTFALIZADEH H., IRANIPOUR S. & S. POURMOHAMMADI (2011): First report of the subfamily Teleasinae (Hym.: Scelionidae) from Iran. — Journal of Field crop Entomology **1** (1): 77-79.
- RAKHSHANI H., EBADI R. & E. RAKHSHANI (2008): Report of *Telenomus chrysopae* (Hym.: Scelionidae) from Iran. — Journal of Entomological Society of Iran **27** (2): Supplement, pp. 21-22.
- REAUDIÈRE G. & R. SKAF (1963): Analyse du complexe des Hyménoptères parasites oophages d'*Eurygaster integriceps* PUT. [Hes. Pentatomidae] en Syrie. — Revue de Pathologie végétale et Dé Entomologie Agricole de France **XLII** (1): 15-25.
- SHAFAEI F., IRANIPOUR S., KAZEMI M.H. & E. ALIZADEH (2011): Diversity and seasonal fluctuations of sunn pest's egg parasitoids (Hymenoptera: Scelionidae) in central regions of west-Azarbaijan province, Iran. — Journal of Field crop Entomology **1** (1): 39-54.
- SAMIN N. (2010): Scelionid wasps (Hymenoptera: Scelionidae) and dominant species in wheat fields of Varamin and vicinity. — M.Sc thesis of Entomology, Islamic Azad University, Tehran Science and Research Branch, 161 pp. (in Persian, English summary).
- SAMIN N., SHOJAI M., KOÇAK E. & M. HAVASKARY (2010a): An annotated list of the Platygastroidea (Hymenoptera) from the Arasbaran biosphere and vicinity, northwestern Iran. — Far Eastern Entomologist **210**: 1-8.
- SAMIN N., SHOJAI M., GHAHARI H. & E. KOÇAK (2010b): A contribution to the Scelionid wasps (Hymenoptera: Platygastroidea: Scelionidae), egg parasitoids of Pentatomidae (Heteroptera) in Tehran province, Iran. — Efflatounia **10**: 7-14.
- SAMIN N., SHOJAI M., ASGARI SH., GHAHARI H. & E. KOÇAK (2010c): Sunn pest (*Eurygaster integriceps* PUTON, Hemiptera: Scutelleridae) and its scelionid (Hymenoptera: Scelionidae) and tachinid (Diptera: Tachinidae) parasitoids in Iran. — Linzer biologische Beiträge **42** (2): 1421-1435.
- SAMIN N., KOÇAK E., GHAHARI H. & M. SHOJAI (2010d): A checklist of Iranian *Telenomus* Haliday (Hymenoptera: Platygastroidae: Scelionidae: Telenominae). — Linzer biologische Beiträge **42** (2): 1437-1444.
- SAMIN N., SHOJAI M., KOÇAK E. & H. GHAHARI (2011a): Distribution of scelionid wasps (Hymenoptera: Platygastroidea: Scelionidae) in Western Iran. — Klapalekiana **47**: 75-82.
- SAMIN N., GHAHARI H., KOÇAK E. & Gh. RADJABI (2011b): A contribution to the scelionid wasps (Hymenoptera: Scelionidae) from some regions of Eastern Iran. — Zoosystematica Rossica **20** (2): 299-304.
- SAMIN N., SHOJAI M., ASGARI Sh., GHAHARI H. & H. KHODDAM (2011c): Population fluctuations of the sunn pest, *Eurygaster integriceps* (Hemiptera: Scutelleridae) in the wheat and barley fields, and introducing of its important parasitoids in aestivation and hibernation shelters in Varamin and Shahre Rey, Iran. — Journal of Agronomy Sciences **5** (6): 78-91 [in Persian, English summary].

- SAMIN N., RADJABI GH. & S. ASGARI (2012): Contribution to the knowledge of Scelionidae (Hymenoptera: Platygastroidea) from Khuzestan province, Southwestern Iran. — *Entomofauna* **33** (3): 17-24.
- SAMIN N. & S. ASGARI (2012a): A study on the Scelionidae (Hymenoptera: Platygastroidea) of Golestan province, Iran. — *Calodema* **200**: 1-5.
- SAMIN N. & S. ASGARI (2012b): A study on the fauna of scelionid wasps (Hymenoptera: Platygastroidea: Scelionidae) in the Isfahan province, Iran. — *Archives of Biological Science Belgrade* **64** (3): 1073-1077.
- TAGHADDOSI M.V. & Gh. RAJABI (1998): Sunn pest egg parasitoids in Zandjan province. — Proceedings of 13th Iranian Plant Protection Congress, Campus of Agriculture and Natural Resources, University of Tehran, Karaj, p. 10.
- TARLA Ş. (1997): Determination of sunn pest, *Eurygaster integriceps* PUT.'s egg parasitoid in Antakya region and investigation on mass culture property of them. — Mustafa Kemal University, Institute of Natural and Applied Sciences, M. Sc. Thesis, Antakya, 57 pp.
- THOMSON C.G. (1861): Sveriges Proctotruper. Tribus IX. Telenomini. Tribus X. Dryinini. — *Öfversigt af Kongliga Vetenskaps-Akademiens Förfhandlinger* **17**: 169-181.
- TUATAY N., KALKANDELEN A. & N. AYSEV (1972): Insect catalogue of plant protection museum (1961-1971). — Plant protection and Quarantine General Directory Publications, 119 pp. (in Turkish).
- VIGGIANI G. & G. MINEO (1974): Identificazione dei parassitoidi del *Gonocerus acuteangulatus* (Goeze). — *Bollettino dell'Istituto di Entomologia Agraria e dell'Osservatorio di Fitopatologia di Palermo* **8**: 143-163.
- VOEGELE J. (1964): *Asolcus bennisi* n. sp. (Hymenoptera, Proctotrupoidea) parasite oophage de *Graphosoma lineata* L. (Het., Pentatomidae). — *Entomophaga* **9**: 119-122.

Authors' addresses:

Hassan GHAHARI
Department of Plant Protection
Yadegar – e- Imam Khomeini (RAH) Branch
Islamic Azad University
Tehran, Iran
E-mail: hghahari@yahoo.com

Peter Neerup BUHL
Plantvägen 1, SE-260 51 Ekeby
Sweden
E-mail: pnbuhl@hotmail.com

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Linzer biologische Beiträge](#)

Jahr/Year: 2015

Band/Volume: [0047_1](#)

Autor(en)/Author(s): Ghahari Hassan, Buhl Peter Neerup

Artikel/Article: [A faunistic study on Scelionidae \(Hymenoptera: Platygastroidea\) from some regions of Iran 449-458](#)