

Linzer biol. Beitr.	47/1	687-690	31.7.2015
---------------------	------	---------	-----------

A newly recorded species of the genus *Dendrothrips* (Thysanoptera: Thripidae) from Iran

Majid MIRAB-BALOU & Azra DOSTY

Abstract: Two genera and five species of Dendrothripinae (Thysanoptera: Thripidae) are recorded in Iran: *Dendrothrips* with 4 species, and *Pseudodendrothrips* with one species. In this paper, *Dendrothrips jasminum* (RAMAKRISHNA & MARGABANDHU, 1939) collected on dill from Qazvin Province is recorded in Iran for the first time. A key is provided to distinguish Dendrothripine thrips from Iran. Diagnostic morphological characters and geographical distribution of the newly recorded species are briefly discussed.

Key words: *Dendrothrips*, new record, identification, Iran.

1. Introduction

The Dendrothripinae (Thysanoptera) is one of the four subfamily recognized in the family Thripidae. The members in this subfamily are leaf-feeding, and can be distinguished by having an enlarged metathoracic furca, which associated with leg muscles; it makes this species with strong jumping abilities (MASUMOTO & OKAJIMA 2003). This subfamily comprises a fauna of 16 genera and 100 species (THRIPSWIKI 2014), of which five species in two genera have been recorded from Iran (Table 1) (MIRAB-BALOU 2013). The members of this subfamily usually feed on dictyledonous tree leaves (MOUND 1997) and also some of them are associated with grasses (such as *Edissa* and *Petrothrips*) (MASUMOTO & OKAJIMA 2003).

The recorded species of this group are not usually considered pests in Iran, although the mulberry thrips, *Pseudodendrothrips mori* is sometimes a minor problem on trees grown for silkworms, and the privet thrips, *Dendrothrips ornatus*, can cause some distortion to the leaves of its host plant (MOUND 1997).

Here, a key is constructed for all Dendrothripine thrips from Iran; with *D. jasminum* is newly recorded for fauna of this country.

2. Materials and methods

The specimens of *D. jasminum* were collected from Qazvin Province, Iran, and prepared and mounted on slide following MIRAB-BALOU & CHEN (2010). The specimens are

deposited in the collection of Department of Plant Protection, College of Agriculture, Ilam University, Iran (ILAMU).

3. Results

Key to genera of the Iranian Dendrothripinae

- 1 Head usually with 2 pairs of ocellar setae. Fore wing with anterior margin curved at apex to join posterior margin, all cilia at apex small, fringe cilia on anterior margin arise ventrally well behind the margin *Dendrothrips* UZEL
- Head usually with 3 pairs of ocellar setae. Fore wing not curved at apex, with long fringe cilia and stout, long apical setae, fringe cilia on anterior margin arise close to the margin *Pseudodendrothrips* SCHMUTZ

Table 1: Dendrothripine thrips recorded from Iran.

Taxa (* new record for Iran)	Distribution in Iran by provinces
<i>Dendrothrips degeeri</i> UZEL	Khorasan-e-Shomali
<i>Dendrothrips jasminum</i> (RAMAKRISHNA & MARGABANDHU) *	Qazvin
<i>Dendrothrips karnyi</i> PRIESNER	Khuzestan
<i>Dendrothrips phyllireae</i> (BAGNALL)	Golestan
<i>Dendrothrips saltator</i> UZEL	Kerman, Hamedan, Khorasan-e-Shomali, Khuzestan
<i>Pseudodendrothrips mori</i> (NIWA)	Guilan, Hamedan, Alborz, Kordestan, Azarbaijan-e-Gharbi

Dendrothrips UZEL

Dendrothrips UZEL, 1895: 159.

This genus currently includes 52 described species in the world (THRIPSWIKI 2014). They are usually found on plants with small hard leaves, such as Oleaceae shrubs and trees, rather than on herbaceous plants (MOUND & KIBBY 1998).

Generic diagnosis. Head wider than long, with anterior margin inset between eyes; eyes large; ocelli widely spaced; ocellar setae pair I absent; postocular setae absent. Antennae 7- to 9-segmented; segments III and IV with simple or forked sense cones. Maxillary palps 2-segmented. Mouth cone short to long. Pronotum wider than long, with a pair of distinct posteroangular setae. Mesospinasternum fused to metasternum. Tarsi 1-segmented, hind tarsi not exceptionally elongated. Fore wings with posterior margin straight, costal setae minute; apex of wing without long seta; anterior fringe cilia inserted on ventral side, far from costal margin; posterior fringe cilia straight. Metathoracic furca lyre-shaped. Meso- and metasternal spinulae present. Abdominal tergites granulated with hexagonal reticulations on the sides, tergite VIII with a posterior comb of setae; tergites I-VIII each with setae S1 elongate and closely spaced. Males without sternal pore plates.

Key to Iranian species of *Dendrothrips* UZEL

- 1 Pronotum in the posterior angles without long setae, or setae that as small as discal setae. Antennae 7, 8- or 9-segmented.....2
- Pronotum in the posterior angles with at least single setae, that clearly longer than discal setae. Antennae 8-segmented.....3
- 2 Antennae 8- or 9-segmented*D. degeeri*
- Antennae 7-segmented*D. phyllireae*
- 3 Antennal segment II pale brown to dark brown, always clearly darker than I4
- Antennal segments I–V uniformly white*D. karnyi*
- 4 Antennal segments III and IV with forked sense cones..... *D. jasminum*
- Antennal segments III and IV with simple sense cones *D. saltator*

Dendrothrips jasminum (RAMAKRISHNA & MARGABANDHU, 1939), new record

Dendrothripiella jasminum RAMAKRISHNA & MARGABANDHU, 1939: 36.

D i a g n o s i s . Body color grayish yellow, all legs yellowish; antennal segments I, III & IV yellowish, segments II, IV–VIII brown to dark brown; wings transparent and white.

F e m a l e m a c r o p t e r a . Head wider than long, with two pairs of ocellar setae; ocelli present; eyes large; mouth cone reaching to prosternum. Antennae 8-segmented; segment II globular and stouter than segment I; segments III and IV with forked sense cones; segments III & IV in equal length. Maxillary palps 2-segmented. Pronotum wider than long, with a pair of distinct posteroangular setae that longer than pronotal discal setae. Mesonotum longitudinally striate; metascutum with small sculpture, median setae situated behind anterior margin. Tarsi 1-segmented. Fore wings first vein with the basal four small setae placed regularly up to the middle of the wing, second vein with 2 small setae at the middle far apart each other; posterior fringe cilia straight. Metathoracic furca lyre-shaped. Meso- and meta-sternal spinulae present. Abdomen elongate and oval-shaped, narrowing towards last segments, with ovipositor well-developed. Abdominal tergites granulated with hexagonal reticulations on the sides, tergite VIII with a comb at posterior margin; tergite I to VIII each with setae S1 elongate and closely spaced; tergite IX with two pairs of setae S1 & S2 long and equal in length; tergite IX with anterior half smooth, posterior half with microtrichia. Abdominal sternites without discal setae, with three pairs of posteromarginal setae; sternite VII with posteromarginal setae situated on posterior margin.

Male. Unknown.

M a t e r i a l e x a m i n e d : IRAN: Qazvin Province: Razjerd, 2 ♀♀ Qazvin, from dill, *Anethum graveolens* (Apiaceae), 29.viii.2013, coll. A. Dosty, (in ILAMU).

D i s t r i b u t i o n . Iran (Qazvin province); India (RAMAKRISHNA & MARGABANDHU 1939).

Acknowledgements

To Islamic Azad University, Takestan Branch, for the financial support given to the second author during her M.Sc. course. We are also grateful to Dr. Kaomud Tyagi, Zoological Survey of India for supplying some references.

References

- MASUMOTO M. & S. OKAJIMA (2003): Two new Dendrothripinae (Insecta: Thysanoptera: Thripidae) from Japan. — *Species Diversity* **8**: 35-46.
- MIRAB-BALOU M. & X.X. CHEN (2010): A new method for preparing and mounting thrips for microscopic examination. — *Journal of Environmental Entomology* **32** (1): 115-121.
- MIRAB-BALOU M. (2013): A checklist of Iranian thrips (Insecta: Thysanoptera). — *Far Eastern Entomologist* **267**: 1-27.
- MOUND L.A. & G. KIBBY (1998): *Thysanoptera: An Identification Guide*. — 2nd Ed. CABI, Oxford and New York, 70 pages.
- MOUND L.A. (1997): Biological diversity. — In: LEWIS T. (ed.), *Thrips as crop pests*. — Wallingford (UK): CAB International. pp. 197-215.
- RAMAKRISHNA T.V. & V. MARGABANDHU (1939): Notes on Indian Thysanoptera with descriptions of new species. — *Indian Journal of Entomology* **1**: 35-48.
- THRIPSWIKI (2014): ThripsWiki – providing information on the World's thrips. — Retrieved June 20, 2014. from <http://thrips.info/wiki/>.
- UZEL H. (1895): *Monographie der Ordnung Thysanoptera*. — Königgrätz, Bohemia pp. 1-472.

Autors' addresses:

Majid MIRAB-BALOU
Department of Plant Protection, College of Agriculture,
Ilam University, Ilam, Iran
Corresponding author: E-mail: majid.mirab@gmail.com

Azra DOSTY
Department of Plant protection, College of Agriculture,
Takestan Branch, Islamic Azad University, Takestan, Iran

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Linzer biologische Beiträge](#)

Jahr/Year: 2015

Band/Volume: [0047_1](#)

Autor(en)/Author(s): Mirab-Balou Majid, Dosty Azra

Artikel/Article: [A newly recorded species of the genus Dendrothrips \(Thysanoptera: Thripidae\) from Iran 687-690](#)