

Linzer biol. Beitr.	49/1	677-686	28.7.2017
---------------------	------	---------	-----------

New records of leafrollers reared in Azerbaijan (Lepidoptera: Tortricidae)

Sheyda MAHARRAMOVA & Hamit AYBERK

A b s t r a c t : 17 species of tortricid moths had been defined for the eastern parts of Azerbaijan during the years of 1994-2015. Five of these, *Ptycholoma lecheana* (LINNAEUS, 1758), *Cacoecimorpha pronubana* HÜBNER, 1799, *Eudemis profundana* ([DENIS & SCHIFFERMÜLLER], 1775), *Hedya salicella* (LINNAEUS, 1758), and *Epinotia demarniana* (FISCHER VON RÖSLERSTAMM, 1840), are new to the Azerbaijan fauna; two of them (*Cacoecimorpha pronubana* and *Epinotia demarniana*) are new to the Caucasian fauna, as well. Leafrollers were collected in the larval and pupal stage from March to September on their food plants according to the sampling methods. The early stages were kept in the laboratory until adult emergence, after which they were killed and pinned. Data on newly recorded leafrollers are given in the text including species name, collection area, coordinates of area, data of collection, food plants on which they were recorded, and sex of specimen(s).

K e y w o r d s : Tortricids, damage, host plants, Azerbaijan, fauna.

Introduction

Tortricidae, commonly known as leafrollers, are one of the most diverse families in the Microlepidoptera. The number of leafroller species in countries bordering Azerbaijan differs among the adjacent regions: 469 species are recorded from Turkey (KOÇAK & KEMAL 2012), 145 species from Iran (KOÇAK & KEMAL 2012), and 139 species from Georgia (ESARTIA 1988). A total of 128 species of leafrollers has been reported for Azerbaijan (MAHARRAMOVA 2016).

Field studies have been conducted between 1994 and 2015 focused on the tortricid fauna of eastern Azerbaijan: rearing 17 species of tortricid moths. Five of these, *Ptycholoma lecheana* (LINNAEUS, 1758), *Cacoecimorpha pronubana* HÜBNER, 1799, *Eudemis profundana* ([DENIS AND SCHIFFERMÜLLER], 1775), *Hedya salicella* (LINNAEUS, 1758), and *Epinotia demarniana* (FISCHER VON RÖSLERSTAMM, 1840), are new to the Azerbaijan fauna; two of them (*Cacoecimorpha pronubana* and *Epinotia demarniana*) are new to the Caucasian fauna, as well.

Materials and Methods

R e s e a r c h A r e a : Gonagkend in the north-east; Absheron-Gobustan in the east and Lankaran region in the south east have been selected as the study areas belonging to

the eastern parts of Azerbaijan (Fig. 1). In the southeast and northeast areas, the plant communities consist of native vegetation such as *Quercus castaneifolia*, *Q.longires*, *Crataegus caucasica*, *Parrotia persica*, *Ulmus carpinifolia*, *Salix kuznetsovii*, *Populus hyrcana*, *Fraxinus excelsior*, *Tilia caucasica* and etc.; in the eastern areas dominated ornamental plants such as *Maclura romifera*, *Amorpha fruticosa*, *Ligustrum lucidum*, *Laurus nobilis*, *Colutea arborescens* and etc. introduced from around the world.

M e t h o d s : Leafrollers were collected in the larval and pupal stage from March to September on their food plants according to the sampling methods of FASULATI (1971). All the specimens were kept in the reserves of Azerbaijan National Academy of Sciences, Institute of Zoology. The early stages were kept in the laboratory until adult emergence, after which they were killed and pinned. Nomenclature follows AARVIK et al. (2013) and ZICHA (2014).

A b b r e v i a t i o n s :

penins.....	peninsula
settl.....	settlement
v	village
s.l.....	sea level
a.s.l.....	above sea level
b.s.l.....	below sea level




Fig. 1: Distribution of leafrollers in Eastern Azerbaijan:

- (1) *Ptycholoma lechaena* (LINNAEUS, 1758)
- (2) *Cacoecimorpha pronubana* (HÜBNER, 1800)
- (3) *Eudemis profundana* (DENIS & SCHIFFERMÜLLER, 1775)

- (4) *Hedya salicella* (LINNAEUS, 1758)
- (5) *Epinotia demarniana* (FISCHER von RÖSLERSTAMM, 1840)

Results and Discussion

The family Tortricidae includes three subfamilies - Tortricinae, Olethreutinae and Chlidanotinae. 17 species belonging to 14 genera and two subfamilies: Tortricinae (9

species) and Olethreutinae (8 species) have been reared in laboratory conditions. Of the 17 species, five (*Ptycholoma lecheana* (L.), *Cacoecimorpha pronubana* HUB., *Eudemis profundana* ([DEN.AND SCHIFF.], *Epinotia demarniana* (FISCH.), *Hedya salicella* (L.) are new to the Azerbaijan fauna, two of which (*Cacoecimorpha pronubana*, *Epinotia demarniana*) are new to the Caucasian fauna (Table 1).

Data on newly recorded leafrollers are given below including species name, collection area, coordinates of area, data of collection, food plants on which they were recorded, and sex of specimen(s). Information about distribution of leafrollers was taken from AARVIK et al. (2013).

T o r t r i c i n a e

Archipini PIERCE & METCALFE 1922

Ptycholoma lecheana (LINNAEUS, 1758)

Ptycholoma lecheanum (LINNAEUS, 1758)

Phalaena (Tortrix) lecheana LINNAEUS, 1758

Tortrix circumclusana CHRISTOPH, 1881

Ptycholoma circumclusana CHOI et al., 2004

Cacoecia magnifica HERRICH-SCHIFFER, 1861

Cacoecia lecheana nipponica OKA, 1925

Orthotaenia obsoletana OKA, 1925

Ptycholoma obsoletana STEPHENS, 1834

S p e c i m e n s e x a m i n e d . Absheron penins., Mardakan settl. (40.490051° 50.160848°, 6 m a.s.l.), 22.05.2001, *Quercus castaneifolia*, 2♀♀ Absheron penins., Mardakan settl. (40.487881° 50.159309°, 4 m a.s.l.), 23.05.2001, *Malus sylvestris*, 3♀♀ Absheron penins., Mardakan settl. (40.491343° 50.165834°, 1 m b.s.l.), 27.05.2013, *Quercus castaneifolia*, 3♀♀

H o s t p l a n t s . *Acer negundo* (Aceraceae) (YASUDA 1975), *Quercus robur* L. (Fagaceae) (DISQUE 1908), *Fraxinus* sp. (Oleaceae) (HANNEMANN 1961), *Abies* sp., *Larix decidua* (Pinaceae) (BRADLEY et al. 1973), *Crataegus* sp. (HANNEMANN 1961), *Malus domestica* (SYLVÉN 1958), *Malus sylvestris* (BRADLEY et al. 1973), *Prunus divaricata*, *P.spinosa* (BUDASHKIN 2009), *Prunus* sp. (BRADLEY et al. 1973), *Sorbus* sp. (Rosaceae) (HANNEMANN 1961), *Populus* sp. (BRADLEY et al. 1973), *Salix* sp. (Salicaceae) (BRADLEY et al. 1973), *Tilia* sp. (Tiliaceae), *Ulmus* sp. (Ulmaceae) (HANNEMANN 1961).

D i s t r i b u t i o n . Europe (Albania, Austria, Belgium, Bosnia & Herzegovina, Britain, Bulgaria, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Ireland, Italy, Latvia, Lithuania, Luxembourg, Macedonia, Norway, Poland, Portugal, Romania, Russia (except Northern parts of Russia), Asia (Asian parts of Turkey, Caucasian countries, Lebanon, Georgia, Arabian peninsula, Armenia, Syria, Jordan, Israel, Iran, Iraq, Korea, China, Japan), Africa (Sinai Peninsula-Egypt).

R e m a r k s . This species was collected in larval stage. Larvae are greenish. Early instars skeletonize leaves; later instars damage buds, flowers, and newly emerged leaves. Larvae attach young leaves by silk and feed inside. In early spring larvae damage buds by eating them from inside. Then they join together several buds, flowers, and/or young leaves with silk. Pupae are black or dark brown. The pupa develops within 9-12 days. Adults emerge in late May or early June. Eggs are yellowish-green. Egg clusters contain 15 to 18 eggs. New to Azerbaijan fauna.

This species has one generation throughout its range. Second and third instars overwinter in fairly dense white cocoons in cracks of the bark of trunks and branches, under leaf litter, fixed to the cortex, as well as under dry bud scales on the branches. Overwintered larvae hatch from their winter cocoons in April and May. In May and early June they gnaw blossoming buds and twisted on treetops leaves and flowers. Older instar larvae can gnaw unripe apples and other fruits outside. Pupate in feeding places or in the crevices of trunks. Adults hatch from the end of May to July. Eggs are laid on the leaves, sometimes along the veins (KUZNETSOV 1994).

***Cacoecimorpha pronubana* HÜBNER, 1799**

Tortrix pronubana HÜBNER, [1796-1799]

Tortrix ambustana FRÖLICH, in GEYER & HÜBNER, 1830

Tortrix hermineana DUPONCHEL, in GODART, 1834

Tortrix herminiana DUPONCHEL, in GODART, 1834

Tortrix isolatana LUCAS, 1848

Tortrix musculana obsoletana STRAND, 1901 (form)

Tortrix perochreana HERRICH-SCHÄFFER, 1856

S p e c i m e n s e x a m i n e d . Absheron penin., Mardakan settl. (40.488998° 50.159894°, 6 m a.s.l.), 06.06.1997, *Laurus nobilis*, 1♀; Absheron penin., Mardakan settl. (40.490087° 50.163831°, 8 m a.s.l.), 12.06.2000, *Laurus nobilis*, 1♀; Lankaran district, Hirkan National Park (38.607268° 48.783872°, 97 m a.s.l.), 27.06.2003, *Ligustrum lucidum*, 2♂♂; Lankaran district, Hirkan National Park (38.608021° 48.803839°, 103 m a.s.l.), 18.06.2008, *Ligustrum lucidum*, 1♂.

H o s t p l a n t s . Larvae are polyphagous, feeding on more than 160 species of plants in 42 families (GILLIGAN & EPSTEIN 2014): *Aegopodium podagraria* (Apiaceae) (ALLEN 1992), *Vinca* sp. (Apocynaceae) (DUGDALE et al. 2005), *Hieracium* sp. (Asteraceae) (BRADLEY et al. 1973), *Brassica* sp. (Brassicaceae) (GARDINER 1982), *Dianthus* sp. (Caryophyllaceae) (BRADLEY et al. 1973), *Euonymus japonicus* (BRADLEY et al. 1973; EMMET 1992; NIELSEN 1993), *Cupressocyparis leylandii* (Cupressaceae) (SOKOLOFF 1983), *Hippophae rhamnoides* (Elaeagnaceae) (BRADLEY et al. 1973), *Vaccinium corymbosum* (Ericaceae) (EASTERBROOKE 1986), *Vaccinium* sp. (Ericaceae) (CALVO AND MOLINA 2003), *Euphorbia amygdaloides* (Euphorbiaceae), *Robinia pseudoacacia* (Fabaceae), *Laurus nobilis* (Lauraceae) (BRADLEY et al. 1973), *Narcissus* sp. (Liliaceae) (ALLEN 1987), *Ligustrum* sp. (Oleaceae), *Fuchsia* sp. (Onagraceae) (BRADLEY et al. 1973), *Pinus halepensis* (Pinaceae) (CASTRESANA et al. 1996), *Punica granatum* (Punicaceae) (VASILYeva 1982), *Fragaria* sp. (Rosaceae) (BRADLEY et al. 1973), *Citrus reticulata* (Rutaceae) (BRADLEY 1987), *Lycopersicon esculentum* (Solanaceae) (BRADLEY et al. 1973), *Vitis* sp. (Vitaceae) (JACOBS 1978).

D i s t r i b u t i o n . North America (Oregon state of U.S.A.), Europe (Albania, Austria, Belgium, Bosnia and Herzegovina, Britain, Cyprus, France (Corsica island), Greece, Hungary, Ireland, Balearic Is., Italy, Malta, Portugal, Romania, Sicily, Slovenia, Spain, Switzerland, The Netherlands (Madeira island), Africa (Sinai Peninsula (Egypt)), Asia (Asian Turkey, Caucasian countries, Lebanon, Syria, Jordan, Israel, Arabian peninsula, Iran, Iraq).

R e m a r k s . This species was collected as larvae in the Eastern and South-Eastern territories of Azerbaijan. The yellow or brownish-green larvae attach 2 or 3 leaves together with silk and eat the parenchyma tissue on the inside of the leaves. At the end of 3rd instar, the larvae cover the leaves with a silken web.

Larvae of all instars can overwinter in fruit on the tree, in fallen fruit, or in fallen leaves. They continue to feed in the winter when the weather is favorable, especially on evergreens. Pupation occurs in a silken cocoon on the food plant. Adults fly from late June until July. Eggs are light green, pupae are blackish brown. The number of generations varies from 2 to 6 depending upon latitude (KUZNETSOV 1994). Newly reported for the Caucasian fauna.

Olethreutinae

Olethreutini WALSINGHAM 1895

Eudemis profundana ([DENIS & SCHIFFERMÜLLER], 1775)

Tortrix profundana [DENIS & SCHIFFERMÜLLER], 1775

Tortrix aethiopiana HAWORTH, [1811]

Tortrix alphonsiana DUPONCHEL, in GODART, 1834

Paedisca nebulana DONOVAN, [1806]

Eudemis profundana f. obscurata GIBEAUX & LUQUET, 1998

Eudemis profundana f. satans GIBEAUX & LUQUET, 1998

Phalaena (Tinea) triangulella GOEZE, 1783

Tortrix wellensiana HÜBNER, [1811-1813]

S p e c i m e n s e x a m i n e d : Azerbaijan: Lankaran district, v.Alekseyevka (38.675763° 48.803327°, at s.l.), 19.05.1995, *Quercus castaneifolia*, 2♂ 4♀ ♀; Lankaran district (38.678006° 48.781246°, 66 m a.s.l.), 10.06.1999, *Quercus castaneifolia*, 12♂ 2♀ ♀; Siyazan district, v.Galaalty (41.083321° 49.022541°, 384 m a.s.l.), 11.05.2001, *Quercus iberica*, 2♂ ♂; Lankaran district, Hirkhan National Park (38.676011° 48.779312°, 104 m a.s.l.), 25.05.2002, *Quercus castaneifolia*, 3♀ ♀; Lankaran district (38.678197° 48.7778540°, 116 m a.s.l.), 09.06.2003, *Quercus castaneifolia*, 2♂ ♂; Siyazan district, v.Galaalty (41.084228° 49.022972°, 389 m a.s.l.), 12.05.2005, *Quercus iberica*, 1♂ 6♀ ♀; Lankaran district, v.Avrora (38.661756° 48.802450°, 2 m a.s.l.), 12.05.2006, *Quercus castaneifolia*, 5♂ 5♀ ♀; Siyazan district, v.Galaalty (41.084182° 49.021693°, 345 m a.s.l.), 02.06.2008, *Quercus iberica*, 11♂ 7♀ ♀; Lenkeran district, v.Alekseyevka (38.675763° 48.803327°, at s.l.), 28.05.2009, *Quercus castaneifolia*, 9♀ ♀; Siyazan district, v.Galaalty (41.084243° 49.021411°, 333 m a.s.l.), 22.05.2010, *Quercus iberica*, 6♀ ♀; Siyazan district, v.Galaalty (41.083653° 49.021133°, 330 m a.s.l.), 10.05.2011, *Quercus iberica*, 11♂ 12♀ ♀; Masally district, v.Geyechol (39.005293° 48.715136°, 10 m b.s.l.) 19.05.2011, *Quercus castaneifolia*, 12♂ 7♀ ♀; Lankaran district, Hirkhan National Park (38.653407° 48.758074°, 104 m a.s.l.), 25.05.2011, *Quercus castaneifolia*, 5♂ 14♀ ♀; Siyazan district, v.Galaalty (41.083647° 49.022791°, 387 m a.s.l.), 14.05.2012, *Quercus iberica*, 32♂ 24♀ ♀; Masally district (38.977001° 48.602869°, 98 m a.s.l.), 22.05.2012, *Quercus castaneifolia*, 18♂ 12♀ ♀.

H o s t p l a n t s : Main food plant is oak. However, the species is found on different forest trees and bushes: *Quercus aliena* (KIM & SOHN 1999), *Quercus pubescens* (BUDASHKIN 2009), *Quercus robur* (DISQUE 1908), *Quercus serrata*, *Malus toringo*, *Prunus padus* (KAWABE 1982), *Prunus ssiori* (Rosaceae) (BAE & SAKAMAKI 1995).

D i s t r i b u t i o n : Europe (Albania, Austria, Belgium, Britain, Bulgaria, Czech Republic, Denmark, Estonia, Finland, France, Germany, Hungary, Ireland, Italy, Latvia, Lithuania, Luxembourg, Macedonia, Norwich, Poland, Portugal, Romania, Russia (except Russia North and Russia Northwest), Sardinia, Sicily, Slovakia, Slovenia, Spain, Sweden, Switzerland, The Netherlands, Ukraine, former Yugoslavia), Africa (Sinai Peninsula (Egypt), Asia (Asian Turkey, Caucasian Russian republics, Arabian peninsula, Lebanon, Syria, Israel, Jordan, Iran, Iraq).

R e m a r k s : Larvae and pupae were collected in North-Eastern and South-Eastern Azerbaijan on different forest plants. Last instar larvae are dark green, very active, and 2.0-2.3 cm in length. Caterpillars damage *Quercus castaneifolia* and *Quercus iberica*.

Caterpillars are found in the web of leaves or young shoots twisted together. Young caterpillars feed on buds, then they move on to the leaves (BRADLEY et al. 1979). They pupate in a leaf roll along the main vein of the leaf (KUZNETSOV 1994) or sometimes on the ground amongst leaf litter in June (BRADLEY et al. 1979).

Newly reported for Azerbaijan fauna.

Olethreutinae: Olethreutini WALSINGHAM 1895

Hedya salicella (LINNAEUS 1758)

S p e c i m e n s e x a m i n e d . Siyazan district, v.Galaalty (41.084632° 49.022847°, 383 m a.s.l.), 14.05.2001, *Salix kuznetsovii*, 1♂; Siyazan district, v.Galaalty (41.083933° 49.022913°, 388 m a.s.l.), 18.05.2005, *Salix kuznetsovii*, 2♂♂ 3♀♀; Siyazan district, v.Galaalty (41.084730° 49.021817°, 345 m a.s.l.), 09.05.2007, *Populus hibrido*, 2♀♀; Siyazan district, v.Galaalty (41.084811° 49.021521°, 332 m a.s.l.), 23.05.2010, *Salix kuznetsovii*, 6♂♂ 4♀♀, *Populus hibrido*, 3♂♂ 2♀♀; Siyazan district, v.Galaalty (41.083030° 49.022369°, 385 m a.s.l.), 03.05.2012, *Salix kuznetsovii*, 4♀♀.

H o s t p l a n t s . *Populus nigra*, *P. tremula* (BRADLEY et al. 1987; JAROS and SPITZER 2002), *Populus* (BUDASHKIN 2009), *Salix alba*, *S. atrocinerea*, *S. caprea* (BRADLEY et al. 1987), *S. cinerea* (Salicaceae) (JAROS & SPITZER 2002).

D i s t r i b u t i o n . Europe (Albania, Austria, Belgium, Bosnia and Herzegovina, Britain, Bulgaria, Czech Republic, Danish mainland, Estonia, Finland, France, Germany, Greece, Hungary, Italy (Sardinia island), Latvia, Lithuania, Luxembourg, Norwich mainland, Poland, Romania, Russia (except northern parts), Slovakia, Slovenia, Spain, Sweden, Switzerland, The Netherlands), Asia (Asian Turkey, Caucasian countries, Georgia, Armenia, Lebanon, Syria, Israel, Jordan, Arabian peninsula, Iran, Iraq) Africa (Sinai Peninsula (Egypt)).

R e m a r k s . Larvae were collected in the North-Eastern territories of Azerbaijan, on the leaves of *Salix kuznetsovii* and *Populus hibrido* (Salicaceae). Adults emerged in late May and early June. Newly recorded for the Azerbaijan fauna.

O le th r e u t i n a e : Eucosmini MEYRICK, 1909

Epinotia demarniana (FISCHER VON RÖSLERSTAMM, 1840)

Paedisca demarniana FISCHER VON RÖSLERSTAMM, 1840

S p e c i m e n s e x a m i n e d . Lankaran district, Hirkan National Park (38.646542° 48.814932°, 14 m b.s.l.), 15.05.1995, *Alnus barbata*, 3♂♂; Lankaran district, Hirkan National Park (38.647164° 48.819530°, 19 m b.s.l.), 27.05.2001, *Alnus barbata*, 3♂♂ 1♀; Lankaran district, Hirkan National Park (38.646154° 48.819281°, 18 m b.s.l.), 01.06.2015, *Alnus barbata*, 2♀♀.

H o s t p l a n t s . *Alnus glutinosa* (BRADLEY 1987), *Betula pubescens* (Betulaceae) (DISQUE 1908), *Salix caprea* (Salicaceae) (BRADLEY 1987).

D i s t r i b u t i o n . Europe (Austria, Belgium, Britain, Czech Republic, Denmark, Estonia, Finland, France (Corsica), Germany, Hungary, Italy (Sardinia, Sicily), Latvia, Lithuania, Luxembourg, Norway, Poland, Romania, Russia, Slovakia, Slovenia, Spain, Sweden, Switzerland, Netherlands).

Remarks. Larvae were collected on *Alnus barbata* in the Hirkan National Park in Lankaran. Adults are on the wing from June to July. The larvae feed within the catkins of *Betula*, *Alnus* and *Salix caprea*. Pupation takes place on the ground in a cocoon in the leaf litter. Newly recorded for the Caucasian fauna.

Acknowledgments

The authors express their deep gratitude to Dr. Mustafa OZDEMIR, Plant Protection and Taxonomy Museum, Ankara, Turkey for determining the tortricids and to Dr. John W. BROWN, Systematic Entomology Laboratory, PSI, Agricultural Research Service, U.S. Department of Agriculture, c/o National Museum of Natural History, Washington, D.C., USA for manuscript editing.

Zusammenfassung

17 Wickler (Tortricidae) konnten im Zeitraum 1994 bis 2015 für den östlichen Teil Aserbaidschans nachgewiesen werden. *Ptycholoma lecheana* (LINNAEUS, 1758), *Cacoecimorpha pronubana* HÜBNER, 1799, *Eudemis profundana* ([DENIS & SCHIFFERMÜLLER], 1775), *Heda salicella* (LINNAEUS, 1758) und *Epinotia demarniana* (FISCHER VON RÖSLERSTAMM, 1840) sind Erstnachweise für Aserbaidschan. Die einzelnen Wickler wurden im Raupen- und Puppenstadium an ihren Futterpflanzen gesammelt, bis zum Adultstadium gezüchtet und anschließend genadelt. Die Neunachweise wurden mit den vorhandenen Informationen in der vorliegenden Arbeit angeführt.

References

- AARVIK L.E. (2013): Fauna Europaea: Tortricidae. — In: KARSHOLT O. & E.J. NIEUKERKEN van, Fauna Europaea: Lepidoptera, Moths. Fauna Europaea version 2.6.2. <http://www.faunaeuropae.org>.
- ABDULLAYEVA Sh.Y. (1988): On the fauna of the leaf-rollers (Lepidoptera, Tortricidae) of the east regions of Azerbaijan (I information). — News of the Academy of Sciences of Azerbaijan, series of biological sciences 4: 69-74 (in Azerbaijani).
- ABDULLAYEVA Sh.Y. (1990): On the fauna of the leaf-rollers (Lepidoptera, Tortricidae) of the east regions of Azerbaijan (II information). — News of the Academy of Sciences of Azerbaijan, series of biological sciences 3: 70-73 (in Azerbaijani).
- AKHUNDOVA-TUAEVA L.M. (1960): To studies on lepidopteron damaging the trees and shrubs in Lankaran zone. Scientific notes of the Azerbaijan State University named after S.M.KIROV, N1 (in Russian).
- ALLEN A.A. (1987): Food plants of *Cacoecimorpha pronubana* HÜBN. — Entomologist's Record and Journal of Variation 99: 152.
- ALLEN A.A. (1992): *Cacoecimorpha pronubana* (HÜBNER) (Lep., Tortricidae) bred from flowers and seed-head of *Aegopodium*. — Entomologist's Record and Journal of Variation 104: 288.
- BAE Y.S. & Y. SAKAMAKI (1995): New larval food plant records of Tortricidae and Carposinidae (Lepidoptera) from Japan. — Tyo to Ga 45: 263-268.
- BOGACHEV A.B. (1951): Leaf-rollers – Tortricidae. — In: Animal world of Azerbaijan, Baku, Publ. AzSSR, 382-383 (in Russian).
- BRADLEY J.D. (1987): Card catalogue of identified reared material received by Bradley for identification from about 1955 to 1987. — original in BMNH Microlepidoptera Section library.

- BRADLEY J.D., TREMEWAN W.G. & A. SMITH (1973): British tortricoid moths. *Cochylidae and Tortricidae: Tortricinae*. — The Ray Society, London, 251 pp.
- BRADLEY J.D., TREMEWAN W.G. & A. SMITH (1979): British tortricoid moths. *Tortricidae: Olethreutinae*. — The Ray Society, London, 336 pp.
- BROWN J.W. (2005): World catalogue of insects. V. 5. *Tortricidae (Lepidoptera)*. — Apollo books, 741 pp.
- BUDASHKIN Yu.I. (1987): Revision of leaf-roller fauna (Lepidoptera, Tortricidae) of Crimea — 95 years Karadag scientific station. 30 years Karadag Nature National Academy of Sciences of Ukraine, 158-207 (in Russian).
- CALVO D. & J.M. MOLINA (2003): Incidencia de *Cacoecimorpha pronubana* (HÜBNER, [1799]) (Lep., Tortricidae) sobre variedades de arándano americano (*Vaccinium* spp., Ericaceae) con bajos requerimientos de horas frío en Andalucía Occidental. — Boletín de Sanidad Vegetal Plagas **29**: 553-561.
- CASTRESANA L., A.NOTARIO, J.M. IGLESIAS (1996): [Note on a tortricid *Cacoecimorpha pronubana* (HÜBNER) which attacks pine trees.] — Boletín de Sanidad Vegetal Plagas **22**: 469-473 (in Spanish).
- Castresana, L., A. Notario, and C. Iglesias. 1996.
- DANILEVSKII A.S. & V.I. KUZNETSOV (1968): Tortricidae, tribe Laspeyresiini. — In: BYKHOVSKII B.E. (ed.), Fauna of the USSR, new ser. N **98**. Lepidoptera. V. 5. Moscow & Leningrad: AS USSR, 636 pp. (in Russian).
- DISQUE H. (1908): Versuch einer microlepidopterologischen Botanik. — Deutsche Entomologische Zeitschrift Iris **21**: 34-147.
- DUGDALE J.S., GLEESON D., CLUNIE L.H. & P.W. HOLDER (2005): A diagnostic guide to Tortricidae encountered in field surveys and quarantine inspections in New Zealand: morphological and molecular characters. — MAF Biosecurity Authority, Wellington, New Zealand, 163 pp.
- EASTERBROOK M.A. (1986): Damage to blueberry (*Vaccinium corymbosum*) by *Cacoecimorpha pronubana* (HÜBNER). — Entomologist's Record and Journal of Variation **98**: 218.
- EMMET A.M. (1992): Life history and habits of the British Lepidoptera. pp. 61-300. — In: EMMET A.M. & J. HEATH (eds), The Moths and Butterflies of Great Britain and Ireland 7. 400 pp., Harley Books, Colchester.
- ESARTIYA G.K. (1988): Leaf-rollers (Lepidoptera; Tortricidae) of the Eastern Georgia: Abstract of dissertation for the degree of candidate of biological sciences. — Leningrad, 24. (In Russian).
- FASULATI K.K. (1971): Field study of terrestrial invertebrates. Moscow, p. 424 (in Russian)
- GARDINER B.O.C. (1982): *Cacoecimorpha pronubana* HNB. (Lep.: Tortricidae) successfully reared on artificial diet, with a note on its diapauses requirements. — Entomologist's Record and Journal of Variation **94**: 122-123.
- GILLIGAN T.M., BAIXERAS J., BROWN J.W. & K.R. TUCK (2012): T@RTS: Online World Catalogue of the Tortricidae (Ver. 2.0).
- GILLIGAN T.M. & M.E. EPSTEIN (2014): Tortricids of Agricultural Importance. Interactive Keys developed in Lucid 3.5. Last updated August 2014.
- HANNEMANN H.J. (1961): Kleinschmetterlinge oder Microlepidoptera. I. Die Wickler (s. str.) (Tortricidae). — In: DAHL F. (ed.), Die Tierwelt Deutschlands **48**: 1-233.
- JACOBS S.N.A. (1978): *Cacoecimorpha pronubana* HÜBNER (Lep., Tortricidae): a greenhouse pest. — Entomologist's Record and Journal of Variation **90**: 266.
- JAROS J. & K. SPITZER (2002): Food plants of Lepidoptera associated with an alder care forest in South Bohemia (Central Europe). — Zoological Studies in the Vrbenske Rybniky Reserve (South Bohemia, CZ). Ročník **42**, Supplement, 5-60.

- KAWABE A. (1982): Tortricidae and Cochylidae, pp. 62-151. — In: INOUE H., SUGI S., KUROKO H., MORIUTI S. & A. KAWABE (eds), The Moths of Japan, Part 1: 62-258, Part 2: 158-183.
- KIM S.S. & J.C. SOHN (1999): Records of host plants of Korean moth species. — Journal Lepidoptera Society Korea **11**: 45-51.
- KOÇAK A.O. & M.K. KEMAL (2012): List of the hitherto recorded pterygot taxa of Turkey (Insecta) (Temporary report of the Entomofauna Project of Turkey-10). — Centre for Entomological Studies Ankara, Memoirs **6**: 1-1649, Van, 1 fig.
- KUZNETSOV V.I. (1978): Tortricidae. — Keys to the European Part of the USSR. V. **4**, Part 1. Lepidoptera, Leningrad: Nauka, pp. 193-680. (in Russian).
- KOSTYUK Yu.A. (1980): Tortricidae. Tortricinae. — In: DOLIN V.G., ed. Fauna of Ukraine, V. **15**. Kiev: Naukova dumka. 424 pp. (in Ukrainian).
- KUZNETSOV V.I. (1994): Family Tortricidae. — In: KUZNETSOV V.I. (ed.), Insects and mites - pests of agricultural plants. V. 3. Lepidoptera. St.Petersburg: Nauka, pp. 51-234 (in Russian).
- KUZNETSOV V.I. (2005): Family Tortricidae (Olethreutidae, Grapholitini). — In: LER P.A. (ed.), Keys to the insects if the Russian Far East. V. 5. Trichoptera and Lepidoptera. Vladivostok: Dalnauka, p. 575 (in Russian).
- MAHARRAMOVA S.M. (2016): Species composition and parasitoid complex of leaf-rollers (Lepidoptera: Tortricidae), damaging to wood-fruit cultures in the Eastern part of Azerbaijan. — Baku, PhD thesis, 27 pp.
- RAZOWSKI J. (2001): Tortricinae and Chlidanotinae. Tortricidae of Europe. — Bratislava: Slamka **1**: 247 pp.
- RAZOWSKI J. (2003): Olethreutinae. Tortricidae of Europe. Bratislava. — Slamka **2**. 301 pp.
- REKACH V.N. & T.A. DOBRETSOVA (1935): Fam. Tortricidae – leaf-rollers. — In: Overview of the pests of technical and forage crops in the cotton areas of the Caucasus. Tiflis pp. 122-123 (in Russian).
- SINEV S.YU. (2008): Tortricidae / SINEV S.Yu., NEDOSHIVINA S.V. — Catalogue of Lepidoptera (Lepidoptera) of Russia. Moscow, pp. 114-148 (in Russian).
- SOKOLOFF P. (1983): *Cacoecimorpha pronubana* (HÜBNER) (Lepidoptera: Tortricidae) on conifers. — Entomologist's Gazette **34**: 124.
- SYLVÉN E. (1958): Studies on fruit leaf tortricids (Lepidoptera), with special reference to the periodicity of the adult moths. — Statens Vaxtskyddsanstalt, Meddelanden [Swedish State Plant Protection Institute Contributions] 11, 74. Stockholm, pp. 135-296.
- VASILYEVA E.A. (1982): Carnation leafroller – a dangerous pest of pomegranate and ornamental plants. — Proceedings of Governmental Nikitin Botanical Garden **87**: 54-62 (In Russian).
- YASUDA T. (1975): The Tortricinae and Sparganothinae of Japan (Lepidoptera: Tortricidae) (Part II). — Bulletin of the University of Osaka Prefecture (series B) **27**: 9-251.
- ZICHA O. (ed.) 1999–2014: Biolib. Dostupné z www:
<http://www.biolib.cz/cz/taxonmap/id361/> [cit. 20.VIII.2014]

Authors' addresses:

Dr. Sheyda MAHARRAMOVA
 Azerbaijan National Academy of Sciences
 Institute of Zoology, Baku, Azerbaijan
 E-mail: mm_sheyda@hotmail.com

Dr. Hamit AYBERK
 Istanbul University, Faculty of Forestry
 Department of Forest Entomology and Protection
 TR-34473 Istanbul, Turkey
 E-mail: hayberk@istanbul.edu.tr

Tab. 1. Trophic relations of leafrollers in eastern Azerbaijan.

Leafrollers	New		Host plants (by studying areas)	Physical- geographica l regions
		Collecting years		
		Mortality (%)	Imago hatching (%)	
	Total			
<i>Ptycholoma lecheana</i> (L.)	Azerbaijan	8	37.5	62.5
<i>Cacoecimorpha pronubana</i> HUB.	Caucasus	5	40	60
<i>Eudemis profundana</i> (DEN. et SCHIFF.)	Azerbaijan	224	62.5	37.5
<i>Hedya salicella</i> L.	Azerbaijan	27	55.5	44.5
<i>Epinotia demarniana</i> (FISCH.)	Caucasus	18	16.6	83.4
				1995, 2001, 2015
				<i>Quercus castaneifolia</i> <i>Malus sylvestris</i>
				<i>Laurus nobilis</i> <i>Ligustrum lucidum</i>
				<i>Quercus castaneifolia</i> <i>Quercus iberica</i>
				<i>Quercus iberica</i> <i>Salix kuznetsovii</i> <i>Populus hibrido</i>
				<i>Alnus barbata</i>
				East
				East south-east
				south-east north-east
				north-east
				south-east

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Linzer biologische Beiträge](#)

Jahr/Year: 2017

Band/Volume: [0049_1](#)

Autor(en)/Author(s): Maharramova Sheyda Mamed, Ayberk Hamit

Artikel/Article: [New records of leafrollers reared in Azerbaijan \(Lepidoptera: Tortricidae\) 677-686](#)