

Linzer biol. Beitr.	49/1	691-696	28.7.2017
---------------------	------	---------	-----------

Notes on Cantharidae (Coleoptera) species occurring in sweet cherry orchards in Kemalpaşa (Izmir) province of western Turkey

Alpay ONARAL & Serdar TEZCAN

A b s t r a c t : In this paper, information on four species, namely *Cantharis delagrangei* DELKESKAMP, 1939, *C. livida* LINNAEUS, 1758, *C. prusiensis* MARSEUL, 1864 and *Rhagonycha fulvaliena* ŠVIHLA, 1995 have been given from sweet cherry orchards in Kemalpaşa (Izmir) province of western Turkey.

K e y w o r d s : Cantharidae, Fauna, Sweet cherry, Turkey.

Introduction

Sweet cherry is a fruit which has significant role in Turkish economy. There are 20 615 760 sweet cherry trees in Turkey and annual production is 535 600 tons (ANONYMOUS 2015a). Cherries are extensively grown in the Aegean, Marmara and Mediterranean regions of Turkey. In the Aegean region, Kemalpaşa (Izmir) is the area where they are commonly widespread (ANONYMOUS 2015b) and 8,8% of Turkish sweet cherry production is realized in this province (ANONYMOUS 2015a). According to 2015 data, there are 2112800 trees bearing fruits in 8681,1 hectares of land in Kemalpaşa province (ANONYMOUS 2015a).

There have been many studies on the pests and the beneficial species found in sweet cherry orchards in Turkey. However, apart from ERTOP & ÖZPINAR (2011) no other study was found which went as far down as the species belonging to Cantharidae (Coleoptera). In this study *Cantharis decipiens* BAUDI, 1871 was reported from sweet cherry orchards in Çanakkale province of Turkey without giving an information on feeding effects and damage (ERTOP & ÖZPINAR 2011).

SAHLBERG (1912-1913), WITTMER (1967-1968, 1969, 1971, 1972, 1975), GÜL-ZÜMREOĞLU (1972), TUATAY et al. (1972), ŠVIHLA (1993, 1994, 1995, 1998, 1999, 2002, 2004, 2009), LODOS (1998), SAYAN (2010), YILDIRIM et al. (2011), SILKIN (2012) and DEMİRÖZER & KARACA (2014) were the other studies conducted on Cantharidae fauna of Turkey.

In some other studies, species belonging to Cantharidae have been evaluated by means of their feeding habits as predators (LANGENSTÜCK & HEIMBACH 1999; RAMSDALE 2002; ALFORD 2003; PRODANOVIĆ et al. 2010). But there is not detailed information on feeding

of these species in sweet cherry orchards. For the purpose of overcoming this absence, conducting a study in the sweet cherry orchards in Kemalpaşa was deemed necessary.

Material and Methods

The material of this study consisted of species within the Cantharidae collected from sweet cherry orchards from the villages of Kemalpaşa province in the months between April and June of 2015.

Studies were conducted in 142 sweet cherry orchards in 24 localities with sizes ranging between 1 to 5 decares. These localities and the number of orchards which were sampled were as follows: Akalan (3), Armuthu (14), Aşağıkızılca (3), Bağyurdu (14), Bayramlı (3), Beşpinar (3), Central province (13), Çambel (3), Damlacık (3), Kamberler (3), Kızılızüm (3), Kurudere (3), Ovacık (3), Ören (20), Örnekköy (8), Sarilar (3), Sinancılar (3), Sütçüler (5), Ulucak (6), Vişneli (5), Yenikurudere (3), Yenmiş (3), Yiğitler (10), Yukarıkızılca (5). - 142 Orchards in total.

In these orchards by walking along the transects of the orchard with the beating tray, falling samples by 25 impact on randomly selected trees in every orchard was collected with the aspirator; sampling was conducted using 25 swings of sweep net for collection of the individuals likely to be found on lower vegetation. Furthermore, in each orchard for 10 minutes the insects on trees were observed and insect samples were collected with aspirator. This was repeated three times for the early, mid-season and late maturing varieties of cherries. The confirmation and identification of the samples were done by Dr. Andreas Kopetz (Naturkundemuseum, Erfurt, Germany).

Results

As a result of this study, a total of four species, namely *Cantharis delagrangei* DELKESKAMP, 1939, *C. livida* LINNAEUS, 1758, *C. prusiensis* MARSEUL, 1864 and *Rhagonycha fulvaliena* ŠVIHLA, 1995 were determined and they were indicated in Table 1.

Those species were reported for the first time from sweet cherry orchards in Turkey. The most common species in the province is *C. livida* sampled 118 orchards (83,09%) from total of 142. This species was followed by *R. fulvaliena* (112 orchards, 78,87%), *C. prusiensis* (20 orchards, 14,08%) and *C. delagrangei* (1 orchard, 0,70%), respectively.

When the numerical and percentage status of the collected samples within the total sample was observed *C. livida* was the most prominent one with 846 samples (59,12%). It was followed by *R. fulvaliena* with 487 samples (34,03%), *C. prusiensis* with 97 samples (6,78%) and *C. delagrangei* with 1 sample (0,07%).

A total of 1431 samples which belong to four species collected. Among those, 664 samples (46,40%) were collected with beating tray, 577 (40,32%) were with aspirator and 190 (13,28%) were with sweep net.

During this study, feeding of *C. prusiensis* and *C. delagrangei* on sweet cherry fruits have not been observed both in nature and in the laboratory. *C. livida* has been observed

as feeding on early and mid-season sweet cherry varieties while *R. fulvaliena* on late season varieties especially on the fruits of high economic variety of 0900 Ziraat both in nature and in laboratory (Fig. 1).



Fig 1: *Rhagonycha fulvaliena* adults feeding on sweet cherry fruit.

Similar studies in other sweet cherry production areas need to consider in Turkey. In this way, it may contribute to the increase of knowledge on this group of insects.

Acknowledgements

Authors would like to express their sincere thanks to Dr. Andreas Kopetz (Naturkundemuseum, Erfurt, Germany) and Prof. Dr. Erol Yıldırım (Atatürk University, Erzurum, Turkey) for confirmations, identifications and literature supply. Support for this study was provided by the Rectorate of Ege University Research Fund grant 15-ZRF-006.

Zusammenfassung

Vorliegende Arbeit gibt Informationen zu den vier Cantharidae-Arten (Coleoptera) *Cantharis delagrangei* DELKESKAMP, 1939, *C. livida* LINNAEUS, 1758, *C. prusiensis* MARSEUL, 1864 und *Rhagonycha fulvaliena* ŠVIHLA, 1995 aus Untersuchungen in einem Süßkirschen-Obstgarten in Kemalpaşa in der türkischen Provinz Izmir im Westen der Türkei.

References

- ALFORD D.V. (2003): Biocontrol of Oilseed Rape Pests. — Blackwell Science Ltd, USA, 1-368.
- ANONYMOUS (2015a): <https://biruni.tuik.gov.tr/bitkiselapp/bitkisel.zul> — (Access date: 20th March 2016).
- ANONYMOUS (2015b): http://www.kemalpasa.gov.tr/default_B0.aspx?content=1002. — (Access date: 20th March 2016).
- DEMİRÖZER O. & I. KARACA (2014): Isparta İli yağ gülü (*Rosa damascena* MILLER) üretim alanlarında bulunan avcı ve parazitoit türler ve yayılışları. — Türkiye Entomoloji Bülteni 4 (3): 171-184.
- ERTOP E. & A. ÖZPINAR (2011): Çanakkale İli kiraz ağaçlarındaki fitofag ve yararlı türler ile bazı önemli zararlıların popülasyon değişimi. — Türkiye Entomoloji Bülteni 1 (2): 109-118.
- GÜL-ZÜMREOĞLU S. (1972): İzmir Bölge Zirai Mücadele Araştırma Enstitüsü Böcek ve Genel Zararlılar Kataloğu, 1928-1969 (I. Kısım). — İstiklal Matbaası, İzmir: 1-119.
- LANGENSTÜCK V.C. & U. HEIMBACH (1999): Bedeutung von Weichkäfer-Larven (Coleoptera: Cantharidae) als Blattlaus-Prädatoren = Importance of larvae of soldier beetles (Coleoptera: Cantharidae) as aphidophagous predators. — Gesunde Pflanzen 51 (3): 86-89.
- LODOS N. (1998): Türkiye Entomolojisi VI (Genel, Uygulamalı ve Faunistik) (I. Basım). — Ege Üniversitesi Ziraat Fakültesi Yayınları No: 529, Ofset Basımevi, Bornova, İzmir, 1-300.
- PRODANOVIĆ D.J., PROTIĆ L. & L. MIHAJLOVIĆ (2010): Predatori i parazitoidi *Cacopsylla pyri* (L.) (Hemiptera: Psyllidae) u Srbiji. — Pesticidi i Fitomedicina (Beograd) 25 (1): 29-42.
- RAMSDALE A.S. (2002): 64. Cantharidae Imhoff 1856, American Beetles (Polyphaga: Scarabaeoidea through Curculionoidea). — CRC Press, Newyork 2: 202-219.
- SAHLBERG J. (1912-1913): Coleoptera Mediterranean Orientalia, quae in Aegyptia, Palaestina, Syria, Caramania atque in Anatolia occidentali anno 1904. — Ofversigt af Finska Vetenskaps-Societetens Förhandlingar 60 (13): 127-132.
- SAYAN M. (2010): Adana'da Buğday Agro-Ekosistemindeki Böcek Türlerinin Belirlenmesi. — Yüksek Lisans Tezi, Çukurova Üniversitesi Ziraat Fakültesi, 1-80 (unpublished).
- SILKIN N. (2012): Akdeniz Bölgesi *Cantharis*, *Boveycantharis* ve *Malthinus* (Cantharidae) cinsleri faunası ve sistematiği üzerine araştırma. — 21. Ulusal Biyoloji Kongresi, 03-07 Eylül 2012, Ege Üniversitesi, İzmir, Türkiye.
- ŠVIHLA V. (1993): Contribution to the knowledge of the genus *Rhagonycha* ESCHSCH. (Coleoptera, Cantharidae) from Eastern Mediterranea. — Entomologica Basiliensis 16: 255-277.
- ŠVIHLA V. (1994): New taxa of Palaearctic Cantharidae (Coleoptera). — Folia Heyrovskiana 2 (9-10): 116-123.
- ŠVIHLA V. (1995): Contribution to the knowledge of the genus *Rhagonycha* ESCHSCHOLTZ (Coleoptera, Cantharidae) II. — Entomologica Basiliensis 18: 71-90.
- ŠVIHLA V. (1998): Contribution to the knowledge of the subfamily Malthininae (Coleoptera: Cantharidae) of Turkey. — Klapalekiana 34: 227-232.
- ŠVIHLA V. (1999): Contribution to the knowledge of the genus *Cantharis* L. and related genera from Turkey and adjacent regions (Coleoptera, Cantharidae). — Entomologica Basiliensis 21: 135-170.
- ŠVIHLA V. (2002): A contribution to knowledge of the subfamily Malthininae (Coleoptera: Cantharidae) from the Western Palaearctic. — Folia Heyrovskiana 10 (2-3): 119-154.

- ŠVIHLA V. (2004): Contribution to the knowledge of the family Cantharidae (Coleoptera) from the Western Palaearctic. — Časopis Národního muzea, Řada přírodovědná (J. Nat. Mus., Nat. Hist. Ser.) **173** (1-4): 77-88.
- ŠVIHLA V. (2009): New species of the subfamily Malthininae (Coleoptera: Cantharidae) from the Western Palaearctic Region. — Acta Entomologica Musei Nationalis Pragae **49** (1): 191-216.
- TUATAY N., KALKANDELEN A. & N. AYSEV (1972): Nebat Koruma Müzesi Böcek Kataloğu (1961-1971). — T.C. Tar. Bak. Zir. Müc. ve Zir. Kar. Gn. Müd. Yayımları, Ankara, 1-119.
- WITTMER W. (1967-1968): 37. Beitrag zur Kenntnis der palaearktischen Cantharidae und Malachiidae (Col.). — Mitteilungen der Schweizerischen Entomologischen Gesellschaft **40** (3-4): 227-280.
- WITTMER W. (1969): Zur Kenntnis der Gattung *Metacantharis* BOURG. (43. Beitrag zur Kenntnis der palaearktischen Cantharidae, Coleoptera). — Verhandlungen der Naturforschenden Gesellschaft in Basel **80**: 70-93.
- WITTMER W. (1971): 51. Beitrag zur Kenntnis der palaearktischen Cantharidae. — Entomologische Arbeiten aus dem Museum G. Frey Tutzing bei München, 226-239.
- WITTMER W. (1972): 56. Beitrag zur Kenntnis der palaearktischen Cantharidae (Col.). — Mitteilungen der Schweizerischen Entomologischen Gesellschaft **45** (1-3): 61-77.
- WITTMER W. (1975): 61. Beitrag zur Kenntnis der palaearktischen Cantharidae und Malachiidae. — Mitteilungen der Schweizerischen Entomologischen Gesellschaft **48** (3-4): 259-267.
- YILDIRIM E., KOPETZ A. & Y. BULAK (2011): Contribution to the knowledge of the Cantharidae (Coleoptera) fauna of Turkey. — Linzer biologische Beiträge **43** (2): 1641-1646.

Authors' addresses:

Alpay ONARAL
 Ege University, Faculty of Agriculture,
 Department of Plant Protection
 TR-Bornova-Izmir, Turkey
 E-mail: alpayonaral@gmail.com

Prof. Dr. Serdar TEZCAN
 Ege University, Faculty of Agriculture,
 Department of Plant Protection
 TR-Bornova-Izmir, Turkey
 E-mail: serdar.tezcan@gmail.com

Table 1: List of Cantharidae species according to the genera and collection methods in Kemerpaşa (İzmir) province of western Turkey.

Species	Number of surveyed orchards	Number of orchards occurring specimens	Rate of orchards occurring specimens (%)	Collection methods, varieties and number of specimens								Total number and rate of collected specimens (%)	
				Insect net				Beating tray					
				E*	M*	L*	E*	M*	L*	E*	M*	L*	
<i>Cantharis delagrangei</i>	142	1	0,70	1	0	0	0	0	0	0	1	0	0,07
<i>Cantharis livida</i>	118	83,09	105	14	0	283	79	3	310	48	4	698	141
<i>Cantharis pruriens</i>	142	20	14,08	29	0	0	29	7	0	27	5	0	72
<i>Rhagonycha fulvaliena</i>	142	112	78,87	1	3	37	8	42	213	6	22	155	15
				136	17	37	320	128	216	343	75	159	786
Total				190			664			577		1431	100,00

* E: early season varieties, M: mid-season varieties, L: late season varieties

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Linzer biologische Beiträge](#)

Jahr/Year: 2017

Band/Volume: [0049_1](#)

Autor(en)/Author(s): Onaral Alpay, Tezcan Serdar

Artikel/Article: [Notes on Cantharidae \(Coleoptera\) species occurring in sweetcherry orchards in Kemalpasa \(Izmir\) province of western Turkey 691-696](#)