

Linzer biol. Beitr.	52/1	337-363	31.7.2020
---------------------	------	---------	-----------

A catalogue of the Apidae (Hymenoptera, Apoidea) of Eritrea

Michael MADL

Abstract: Currently 72 species of the family Apidae are known from Eritrea including the subfamilies Apinae (48 species), Nomadinae (four species) and Xylocopinae (22 species).

Key words: Apidae, Apinae, Nomadinae, Xylocopinae, catalogue, Eritrea

Introduction

Species of the family Apidae belong to the long-tongued bees as the family Megachilidae. They have diverse biologies ranging from solitary to eusocial behaviour including parasitism.

The apid fauna of Eritrea is poorly known and the records are mainly based on material collected more than 100 years ago. Hitherto 74 species have been recorded including the honey bee *Apis mellifera* LINNAEUS, 1758, which is used in apiculture. The subfamily Apinae is represented by 48 species, the subfamily Nomadinae by four species and the subfamily Xylocopinae by 22 species. Nothing is known about the biology of the Eritrean bees.

Currently there are no records of the dwarf honey bee, *Apis florea* FABRICIUS, 1787, which has been recorded from the neighbour countries Sudan, Ethiopia and Djibouti (MADL 2018).

Abbreviations

Afr. Reg..... Afrotropical Region
biogeogr. biogeography
biol. biology
cat..... catalogue
descr..... description
design. designation
fig. (figs) figure(s)
orig. original
pl. (pls) plate(s)
typ. gen..... typus generis
typ. subgen. typus subgeneris
syn..... synonym
tax. taxonomy

Annotated catalogue

Subfamily A p i n a e

***Amegilla (Megamegilla) acraensis* (FABRICIUS, 1793)**

Podalirius acraensis FABRICIUS, 1793: MAGRETTI 1906: 10 (Sabarguma), 15 (Keren).

Anthophora acraensis (!) (FABRICIUS, 1793): FRIESE 1909b: 261 (cat. Africa), 263 (key ♂), 265 (key ♀), 269 (tax., Keren), 466 (cat. Africa).

Anthophora acraensis (!) (FABRICIUS, 1793): FRIESE 1915: 272 (Keren).

Amegilla acraensis (FABRICIUS, 1793): EARDLEY & URBAN 2010: 433 (cat. Afr. Reg. (Eritrea, not Ethiopia)).

**Podalirius bipartitus* SMITH, 1854: MAGRETTI 1906: 10 (Sabarguma).

**Anthophora bipartita comperei* n.ssp.: COCKERELL 1935: 1 (descr. ♂, Nefasit).

Anthophora bipartita ssp. *comperei* COCKERELL, 1935: BROOKS 1988: 559 (syn.).

D i s t r i b u t i o n : Keren, Nefasit, Sabarguma.

Amegilla acraensis is widely distributed in the Afrotropical Region, but is unknown from the neighbour countries.

***Amegilla antimena* (DE SAUSSURE, 1890)**

Podalirius antimerus (!) DE SAUSSURE, 1890: MAGRETTI 1906: 11 (Ghinda).

D i s t r i b u t i o n : Ghinda.

The record from Eritrea, which is not mentioned in FRIESE (1909b, 1915) and EARDLEY & URBAN (2010: 436), should be checked, because *Amegilla antimena* is endemic to the Malagasy Subregion (Seychelles (Aldabra), Madagascar).

***Amegilla atrocincta* (LEPELETIER DE SAINT-FARGEAU, 1841)**

Podalirius plumipes var. *atrocinctus* (LEPELETIER DE SAINT-FARGEAU, 1841): MAGRETTI 1906: 12 (Mai-Hadegà).

Amegilla atrocincta (LEPELETIER DE SAINT-FARGEAU, 1841): EARDLEY & URBAN 2010: 436 (cat. Afr. Reg. (Eritrea under Ethiopia)).

**Podalirius plumipes* FABRICIUS, 1781: MAGRETTI 1906: 15 (Keren).

Anthophora plumipes FABRICIUS, 1781: FRIESE 1915: 272 (Mai Hadega).

D i s t r i b u t i o n : Keren, Mai Hadega.

Amegilla atrocincta is widely distributed in the Afrotropical Region and is also recorded from Ethiopia.

***Amegilla byssina* (KLUG, 1845)**

Anthophora byssina (KLUG, 1845): FRIESE 1915: 273 (Eritrea without locality).

Amegilla byssina (KLUG, 1845): EARDLEY & URBAN 2010: 437 (cat. Afr. Reg. (Eritrea under Ethiopia)).

D i s t r i b u t i o n : no exact locality.

Amegilla byssina is known from the Afrotropical (Eritrea, Ethiopia) and Palaearctic regions.

***Amegilla caerulea* (FRIESE, 1905)**

Anthophora caerulea FRIESE, 1905: FRIESE 1909b: 262 (cat. Africa), 264 (key ♂), 280 (tax., descr. ♂, biol., Gheleb), 466 (cat. Africa).

Anthophora caerulea FRIESE, 1905: FRIESE 1915: 273 (Gheleb), 290 (tax., descr. ♀, Gheleb).

Amegilla caerulea (FRIESE, 1905): EARDLEY & URBAN 2010: 438 (cat. Afr. Reg. (Eritrea under Ethiopia)).

D i s t r i b u t i o n : Gheleb.

Amegilla caerulea is widely distributed in the Afrotropical Region and is also known from Ethiopia.

***Amegilla calens* (LEPELETIER DE SAINT-FARGEAU, 1841)**

Anthophora calens LEPELETIER DE SAINT-FARGEAU, 1841: WALKER 1871: 55 (tax., Massawa).

D i s t r i b u t i o n : Massawa.

Amegilla calens is widely distributed in the Afrotropical Region and is also known from Sudan. The record from Eritrea is not mentioned in INNES BEY (1911: 107), FRIESE (1909b: 275, 1915) and EARDLEY & URBAN (2010: 438).

***Amegilla cana* (WALKER, 1871)**

Anthophora cana n.sp: WALKER 1871: 56 (descr. ♀ ♂, Harkeko, Massawa).

Podalirius canus (WALKER, 1871): DE DALLA TORRE 1896: 262 (tax., world cat. (Eritrea, not Egypt)).

Anthophora cana WALKER, 1871: INNES BEY 1911: 107 (Harkeko, cat. Lord Collection).

Anthophora cana WALKER, 1871: ALFKEN 1926: 101 (syn. of *Anthophora byssina* KLUG).

Amegilla cana (WALKER, 1871): BROOKS 1988: 515 (tax.), 561 (app. 3: world cat.).

D i s t r i b u t i o n : Harkeko, Massawa.

Amegilla cana, which is not mentioned in FRIESE (1909b, 1915) and EARDLEY & URBAN (2010), is only known from Eritrea.

***Amegilla capensis* (FRIESE, 1905)**

Anthophora capensis FRIESE, 1905: FRIESE 1909b: 262 (cat. Africa), 263 (key ♂), 277 (tax., descr. ♂, Asmara), 466 (cat. Africa).

Anthophora capensis FRIESE, 1905: FRIESE 1915: 273 (Asmara).

Amegilla capensis (FRIESE, 1905): EARDLEY & URBAN 2010: 438 (cat. Afr. Reg. (Eritrea, not Ethiopia)).

D i s t r i b u t i o n : Asmara.

Amegilla capensis is widely distributed in the Afrotropical Region, but is not recorded from West Africa. It is also unknown from the neighbour countries.

***Amegilla circulata* (FABRICIUS, 1781)**

Anthophora circulata (FABRICIUS, 1781): FRIESE 1915: 273 (Eritrea without locality).

Amegilla circulata (FABRICIUS, 1781): EARDLEY & URBAN 2010: 439 (cat. Afr. Reg. (Eritrea, not Ethiopia)).

D i s t r i b u t i o n : no exact locality.

Amegilla capensis is widely distributed in the Afrotropical Region, but is not recorded from the neighbour countries.

***Amegilla eritrina* (FRIESE, 1915)**

Anthophora eritrina n.sp.: FRIESE 1915: 273 (Asmara), 289 (descr. ♀ ♂, Asmara).

Amegilla (Zebramegilla) eritrina (FRIESE, 1915): BROOKS 1988: 504 (tax.), 564 (app. 3: world cat.).

Anthophora eritrina FRIESE, 1915: RASMUSSEN & ASCHER 2008: 46 (cat. taxa described by Friese).

Amegilla eritrina (FRIESE, 1915): EARDLEY & URBAN 2010: 440 (cat. Afr. Reg. (Eritrea, not Ethiopia)).

D i s t r i b u t i o n : Asmara.

Amegilla eritrina is also known from Ghana, Kenya and Nigeria.

***Amegilla niveocincta* SMITH, 1854**

Amegilla (Zonamegilla) niveocincta SMITH, 1854: BROOKS 1988: 511 (tax.), 569 (app. 3: tax., world cat.).

**Anthophora bimaculifera* n.sp.: WALKER 1871: 57 (descr. ♂, Harkeko).

Anthophora bimaculifera WALKER, 1871: BROOKS 1988: 569 (syn.).

**Podalirius bimaculosus* n.nom.: DE DALLA TORRE 1896: 260 (replacement name, world cat. (Eritrea, not Egypt)).

Podalirius bimaculosus DE DALLA TORRE, 1896: FRIESE 1897: 289 (tax., orig. descr.).

Podalirius bimaculosus DE DALLA TORRE, 1896: BROOKS 1988: 569 (syn.).

D i s t r i b u t i o n : Harkeko.

Amegilla niveocincta is known from the Oriental Region: The record from Eritrea is not mentioned in FRIESE (1909b) and EARDLEY & URBAN (2010).

***Amegilla nubica* (LEPELETIER DE SAINT-FARGEAU, 1841)**

Anthophora nubica LEPELETIER DE SAINT-FARGEAU, 1841: MAGRETTI 1884b: 628 (tax., biol., Keren, Khor Lebka).

Anthophora nubica LEPELETIER DE SAINT-FARGEAU, 1841: MAGRETTI 1885: 330 (Keren), 333 (Khor Lebka), 352 (cat. expedition).

Podalirius nubicus (LEPELETIER DE SAINT-FARGEAU, 1841): MAGRETTI 1906: 10 (biol., Massawa), 12 (Mai Hadegà), 15 (Keren), 19 (Ailet).

Anthophora nubica LEPELETIER DE SAINT-FARGEAU, 1841: GUIGLIA 1939a: 202 (tax., Keren, Massawa, Sabarguma).

D i s t r i b u t i o n : Ailet, Keren, Khor Lebka, Mai Hadega, Massawa.

Amegilla nubica is known from the Afrotropical (e.g. Sudan) and Palaearctic regions. The record from Eritrea is not mentioned in FRIESE (1909b: 271, 1915) and EARDLEY & URBAN (2010: 444).

***Amegilla pulverea* (WALKER, 1871)**

Anthophora pulverea n.sp.: WALKER 1871: 57 (descr. ♂, Harkeko).

Podalirius pulvereus (WALKER, 1871): DE DALLA TORRE 1896: 283 (tax., world cat. (Eritrea under Egypt)).

Anthophora pulverea WALKER, 1871: INNES BEY 1911: 107 (Harkeko, cat. Lord Collection).

Amegilla pulverea (WALKER, 1871): BROOKS 1988: 515 (tax.), 571 (app. 3: tax., world cat.).

D i s t r i b u t i o n : Harkeko.

Amegilla pulverea, which is not mentioned in FRIESE (1909b, 1915) and EARDLEY & URBAN (2010), is only known from Eritrea.

***Amegilla punctifrons* (WALKER, 1871)**

- Anthophora punctifrons* n.sp.: WALKER 1871: 57 (descr. ♂, Massawa).
Podalirius punctifrons (WALKER, 1871): DE DALLA TORRE 1896: 283 (tax., world cat. (Eritrea, not Egypt), footnote: tax.).
Anthophora punctifrons WALKER, 1871: INNES BEY 1911: 107 (without locality, cat. Lord Collection).
Amegilla (Zebramegilla) punctifrons (WALKER, 1871): BROOKS 1988: 504 (tax.), 571 (app. 3: tax., world cat.).
Amegilla (Zebramegilla) punctifrons (WALKER, 1871): EARDLEY 1994:38 (key ♂, 40 (fig. 105: map), 43 (tax. descr. ♀ ♂, Eritrea not seen), 45 (figs 106-110), 64 (figs 159, 160).
Amegilla punctifrons (WALKER, 1871): EARDLEY & URBAN 2010: 455 (cat. Afr. Reg. (Eritrea, not Egypt)).

D i s t r i b u t i o n : Massawa.

Amegilla punctifrons is widely distributed in the Afrotropical Region and is also recorded from Sudan. The record from Eritrea is not mentioned in FRIESE (1909b, 1915).

***Amegilla quadrifasciata* (DE VILLERS, 1789)**

- Anthophora quadrifasciata* (DE VILLERS, 1789): MAGRETTI 1884b: 627 (tax., Keren).
Anthophora quadrifasciata (DE VILLERS, 1789): DE DALLA TORRE 1896: 286 (world cat.).
Anthophora quadrifasciata (DE VILLERS, 1789): GUIGLIA 1932: 131 (tax., Ghinda).
Amegilla quadrifasciata (DE VILLERS, 1789): EARDLEY & URBAN 446 (tax., cat. Afr. Reg.).

D i s t r i b u t i o n : Ghinda, Keren.

Anthophora quadrifasciata, which is not mentioned in FRIESE (1909b, 1915), is widely distributed in the Palaearctic Region. In the Afrotropical Region it is recorded from Eritrea and Socotra.

***Anthophora alternans* (KLUG, 1845)**

- Podalirius quadrifasciatus* var. *alternans* (KLUG, 1845): MAGRETTI 1906: 10 (biol., Massawa).

D i s t r i b u t i o n : Massawa.

Anthophora alternans, which is not mentioned in FRIESE (1909b, 1915) and EARDLEY & URBAN (2010), is known from the Palaearctic Region.

***Anthophora armata* FRIESE, 1905**

- Anthophora armata* n.sp.: FRIESE 1905: 234 (descr. ♂, Eritrea without locality).
Anthophora armata FRIESE, 1905: FRIESE 1909a: 139 (tax., cat. Afr. Reg.).
Anthophora armata FRIESE, 1905: FRIESE 1909b: 89 (map 1), 261 (tax., cat. Africa), 263 (key ♂), 264 (key ♀), 269 (tax., descr. ♀ ♂, Eritrea without locality), 467 (cat. Africa).
Anthophora armata FRIESE, 1905: STRAND 1912b: 136 (tax.).
Anthophora armata FRIESE, 1905: FRIESE 1915: 272 (tax., Eritrea without locality).
Anthophora (Paramegilla) armata FRIESE, 1905: BROOKS 1988: 476 (tax.), 560 (app. 3: tax., world cat.).
Anthophora (Paramegilla) armata FRIESE, 1905: EARDLEY & BROOKS 1989: 40 (key), 44 (tax., type design. by Brooks, descr. ♀ ♂, Eritrea without locality), 45 (fig. 175 (map), 46 (figs 176, 177, 182, 183), 55 (figs 206, 207).
Anthophora (Paramegilla) armata FRIESE, 1905: RASMUSSEN & ASCHER 2008: 25 (cat. taxa described by Friese).
Anthophora armata FRIESE, 1905: EARDLEY & URBAN 2010: 449 (cat. Afr. Reg. (Eritrea, not Ethiopia)).

**Anthophora armata tetra* n.var.: FRIESE 1923: 10 (descr. ♂, Eritrea without locality).

Anthophora armata tetra FRIESE, 1922: BROOKS 1988: 560 (syn.).

Anthophora armata tetra FRIESE, 1922: RASMUSSEN & ASCHER 2008: 103 (cat. taxa described by Friese).

D i s t r i b u t i o n : no exact locality.

Anthophora armata is widely distributed in the Afrotropical (missing in West Africa) and Palaearctic (Egypt) regions. It is unknown from the neighbour countries.

***Anthophora basalis* SMITH, 1854**

Podalirius basalis (SMITH, 1854): MAGRETTI 1906: 10 (Sabarguma), 19 (Ailet).

D i s t r i b u t i o n : Ailet, Sabarguma.

Anthophora basalis is only known from southern Africa (Botswana, Namibia, South Africa) and the record from Eritrea is not mentioned in FRIESE (1909b, 1915) and EARDLEY & URBAN (2010: 450). The identification should be checked.

***Anthophora bimaculata* (PANZER, 1798)**

Anthophora bimaculata (PANZER, 1798): FRIESE 1915: 273 (Eritrea without locality).

D i s t r i b u t i o n : no exact locality.

Anthophora bimaculata, which is not mentioned in EARDLEY & URBAN (2010), is widespread in the Palaearctic Region. The identification should be checked.

***Anthophora bimaculifera* WALKER, 1871**

Anthophora bimaculifera n.sp.: WALKER 1871: 56 (descr. ♂, Harkeko).

Podalirius bimaculifer (!) (WALKER, 1871): DE DALLA TORRE 1896: 260 (world cat. (Eritrea, not Egypt).

Podalirius bimaculifer (!) (WALKER, 1871): FRIESE 1897: 23 (cat. Pal. Reg.), 289 (tax., orig. descr.).

Anthophora bimaculifer (!) WALKER, 1871: INNES BEY 1911: 107 (Harkeko, cat. Lord Collection).

D i s t r i b u t i o n : Harkeko.

Anthophora bimaculifera, which is not mentioned in FRIESE (1909b, 1915), BROOKS (1988) and EARDLEY & URBAN (2010), is recorded from the Afrotropical (Eritrea, Sudan) and Palaearctic (Yemen) region. The status is uncertain.

***Anthophora eritrea* BROOKS, 1988**

Anthophora (Heliophila) eritrea n.nom.: BROOKS 1988: 492 (tax.), 564 (replacement name for *A. flaviventris* FRIESE, 1915, world cat.).

Anthophora eritrea BROOKS, 1988: EARDLEY & URBAN 2010: 451 (cat. Afr. Reg. (Eritrea, not Ethiopia)).

**Anthophora flaviventris* n.sp.: FRIESE 1915: 273 (Asmara), 289 (as *A. flaviventris*, descr. ♀, Asmara).

Anthophora flaviventris FRIESE, 1915: HEDICKE 1929: 71 (syn. of *A. kristenseni* FRIESE, 1915, Asmara).

Anthophora flaviventris FRIESE, 1915: RASMUSSEN & ASCHER 2008: 50 (cat. taxa described by Friese).

D i s t r i b u t i o n : Asmara.

Anthophora eritrea is only known from Eritrea. *Anthophora flaviventris* FRIESE, 1915 is a

junior homonym of *Podalirius orientalis flaviventris* FRIESE, 1896 in the genus *Anthophora* LATREILLE, 1803.

***Anthophora hastula* VACHAL, 1910**

Anthophora (!) *hastula* n.sp.: VACHAL 1910: 533 (descr. ♀ ♂, Asmara).

Anthophora hastula VACHAL, 1910: BROOKS 1988: 450 (types not seen), 565 (app. 3: world cat.).

Anthophora hastula VACHAL, 1910: EARDLEY & URBAN 2010: 452 (cat. Afr. Reg. (Eritrea under Ethiopia)).

D i s t r i b u t i o n : Asmara.

Anthophora hastula, which is not mentioned in FRIESE (1915), is known from Eritrea and Ethiopia.

***Anthophora inclyta* WALKER, 1871**

Anthophora inclyta n.sp.: WALKER 1871: 58 (descr. ♂, Rafla).

Podalirius inclytus (WALKER, 1871): DE DALLA TORRE 1896: 273 (tax., world cat. (Eritrea under Egypt)).

Podalirius inclytus (WALKER, 1871): FRIESE 1897: 298 (tax., orig. descr.).

Anthophora (*Paramegilla*) *inclyta* WALKER, 1871: BROOKS 1988: 476 (tax.), 566 (app. 3: tax., world cat.).

D i s t r i b u t i o n : Rafla.

Anthophora inclyta is known from the Afrotropical (Eritrea, Socotra) and Palearctic regions. The record from Eritrea is not mentioned in FRIESE (1909b, 1915), INNES BEY (1911: 107) and EARDLEY & URBAN (2010: 452).

***Anthophora leucomelaena* DE DALLA TORRE, 1896**

**Anthophora melaleuca* WALKER, 1871: INNES BEY 1911: 107 (Massawa, cat. Lord Collection).

D i s t r i b u t i o n : Massawa.

Anthophora leucomelaena, which is not mentioned in FRIESE (1915) and EARDLEY & URBAN (2010), is known from the Palearctic Region (e.g. Yemen). The record from Eritrea by INNES BEY (1911) is doubtful, because WALKER (1871) described the species from Egypt (Cairo).

***Anthophora nigrifacies* FRIESE, 1905**

Anthophora nigrifacies n.sp.: FRIESE 1905: 240 (descr. ♀, without locality).

Anthophora nigrifacies FRIESE, 1905: FRIESE 1909a: 139 (tax., cat. Afr. Reg.), 143 (key ♂).

Anthophora nigrifacies FRIESE, 1905: FRIESE 1909b: 262 (cat. Africa), 265 (key ♀), 279 (tax., descr. ♀, Eritrea without locality), 466 (cat. Africa).

Anthophora nigrifacies (!) FRIESE, 1905: FRIESE 1915: 273 (Eritrea without locality).

Anthophora (*Pyganthophora*) *nigrifacies* FRIESE, 1905: BROOKS 1988: 460 (tax.), 462 (tax.), 464 (tax.), 568 (app. 3: world cat.).

Anthophora nigrifacies FRIESE, 1905: EARDLEY & URBAN 2010: 454 (cat. Afr. Reg. (Eritrea, not Ethiopia)).

D i s t r i b u t i o n : no exact locality.

Anthophora nigrifacies is only known from Eritrea. BROOKS (1988: 462, 464) placed this species in two different species groups.

***Anthophora senescens* LEPELETIER DE SAINT-FARGEAU, 1841**

Anthophora senescens LEPELETIER DE SAINT-FARGEAU, 1841: FRIESE 1915: 273 (Asmara).

D i s t r i b u t i o n : Asmara.

Anthophora senescens, which is known from the southern Palaearctic Region, is not mentioned in EARDLEY & URBAN (2010).

***Apis mellifera* LINNAEUS, 1758**

The honey bee is divided in many races or subspecies, of which two or three are used in Eritrean apiculture. As the taxonomy is not fixed, I did not search for literature.

***Hypotrigona gribodoi* (MAGRETTI, 1884)**

Trigona Gribodoi n.sp.: MAGRETTI 1884b: 630 (descr. ♂, without locality).

Trigona Gribodoi n.sp.: MAGRETTI 1885: 330 (Keren), 353 (cat. expedition).

Trigona gribodoi MAGRETTI, 1884: KOHL 1894: 281 (tax., descr. ♀, Africa without locality).

Trigona gribodoi MAGRETTI, 1884: MAGRETTI 1895: 153 (footnote 1: cat. Afr. Reg.)

Melipona gribodoi (MAGRETTI, 1884): DE DALLA TORRE 1896: 579 (tax., world cat. (Eritrea, not East Africa)).

Trigona gribodoi MAGRETTI, 1884: FRIESE 1909b: 444 (cat. Africa, key ♂), 455 (key ♀), 456 (tax., descr. ♂, Eritrea without locality (MAGRETTI 1884b cited, but Eritrea not mentioned under distribution), 468 (cat. Africa).

Trigona (Hypotrigona) gribodoi (MAGRETTI, 1884): COCKERELL 1934: 47 (tax., typ. subgen.), 54 (tax.), 62 (fig. 2).

Trigona (Hypotrigona) gribodoi (MAGRETTI, 1884): SANDHOUSE 1943: 560 (type cat. genera and subgenera of bees).

Trigona (Hypotrigona) gribodoi (MAGRETTI, 1884): GUIGLIA 1956: 309 (tax., Keren, footnotes 1: (orig. descr.), 2 (distr.), 3 (tax.)).

Hypotrigona gribodoi (MAGRETTI, 1884): MOURE 1961: 220 (typ. gen.), 221 (tab. 11: tax., Eritrea without locality (type examined)), 222 (tax., cat. Afr. Reg.).

Hypotrigona gribodoi (MAGRETTI, 1884): MICHENER 1997: 30 (type cat. genus-group names).

Hypotrigona gribodoi (MAGRETTI, 1884): EARDLEY 2004: 83 (tax., descr. ♀ ♂, biol., Keren), 84 (figs 33-36), 85 (fig. 41: map (missing Keren)), 87 (tab. 3: tax.).

Hypotrigona gribodoi (MAGRETTI, 1884): EARDLEY & URBAN 2010: 479 (cat. Afr. Reg.).

Hypotrigona gribodoi (MAGRETTI, 1884): PAULY & HORA 2013: 12 (key, type design., figs 7a, b, biol., Keren), 13 (figs 8a-d), 17 (fig.13: map Ethiopia).

**Trigona* n.sp.?: MAGRETTI 1884a: 252 (cat. expedition).

D i s t r i b u t i o n : Keren.

Hypotrigona gribodoi, which is not mentioned in FRIESE (1915), is widely distributed in the Afrotropical Region and is also recorded from Ethiopia and Sudan.

***Liotrigona bottegoi* (MAGRETTI, 1895)**

Trigona bottegoi MAGRETTI, 1895: COCKERELL 1935: 1 (biol., Nefasit (Martini's Concession)).

Liotrigona bottegoi (MAGRETTI, 1895): EARDLEY 2004: 84 (figs 38-40), 88 (fig. 43: map (missing)), 89 (tax., descr. ♀ ♂, tab. 4 (tax.), Eritrea without locality).

Liotrigona bottegoi (MAGRETTI, 1895): EARDLEY & URBAN 2010: 481 (cat. Afr. Reg. (Eritrea under Ethiopia)).

D i s t r i b u t i o n : Nefasit (Martini's Concession)).

Trigona bottegoi is widely distributed in the Afrotropical Region and is also recorded from Ethiopia. The record from Eritrea is missing in PAULY & HORA (2013: 18 (under general distribution), 20 (fig. 19a: map)).

Meliponula beccarii (GRIBODO, 1879)

- Trigona Beccarii* n.sp.: GRIBODO 1879: 340 (descr. ♂, Keren).
Trigona beccarii GRIBODO, 1879: MAGRETTI 1884b: 630 (tax., biol., Sogodas).
Trigona beccarii GRIBODO, 1879: MAGRETTI 1885: 299 (Sogodas), 353 (cat. expedition).
Trigona beccarii GRIBODO, 1879: MAGRETTI 1895: 153 (footnote 1: cat. Afr. Reg.).
Melipona beccarii (GRIBODO, 1879): DE DALLA TORRE 1896: 579 (tax., world cat. (Eritrea, not Ethiopia)).
Trigona beccarii GRIBODO, 1879: FRIESE 1909b: 444 (tax., cat. Africa), 445 (key ♂, tax., descr. ♂, Keren), 468 (cat. Africa).
Trigona beccarii GRIBODO, 1879: FRIESE 1915: 267 (biogeogr.), 276 (Eritrea without locality, Keren).
Trigona beccarii GRIBODO, 1879: COCKERELL 1933b: 462-3 (tax., Keren).
Trigona (Meliponula) beccarii GRIBODO, 1879: COCKERELL 1934: 58 (tax., descr. ♂, Keren).
Trigona beccarii GRIBODO, 1879: COCKERELL 1935: 1 (biol., Nefasit (Martini's Concession)).
Meliplebeia beccarii (GRIBODO, 1879): MOURE 1961: 229 (typ. gen.), 231 (tax., descr. ♂, Keren).
Meliponula (Meliplebeia) beccarii (GRIBODO, 1879): EARDLEY 2004: 70 (key), 75 (tax., descr. ♀ ♂, Keren, without locality), 76 (figs 20-23), 77 (fig. 25: map).
Meliponula beccarii (GRIBODO, 1879): EARDLEY & URBAN 2010: 473 (cat. Afr. Reg.).
Meliponula beccarii (GRIBODO, 1879): PAULY & HORA 2013: 6 (tax., lectotype design., biol., Keren), 7 (figs 1a-f), 8 (figs 2a-f), 9 (figs 3a: map Africa, 3b: map Ethiopia), 10 (figs 4a-c: biol.), 11 (figs 5, 6: both biol.).
Meliponula (Meliplebeia) beccarii (GRIBODO, 1879): PENATI & MARIOTTI 2015: 26 (tax., Keren, cat. Hymenoptera described by Gribodo).
Meliplebeia beccarii (GRIBODO, 1879): MADL 2018: 534 (tax., paralectotype design., Bogos (= Keren)), 535 (figs 2, 3).

D i s t r i b u t i o n : Keren, Nefasit (Martini's Concession), Sogodas.

Meliponula beccarii is widely distributed in the Afrotropical Region and is also recorded from Ethiopia and Sudan.

Tetralonia amoena (WALKER, 1871)

- Tetralonia amoena* n.sp: WALKER 1871: 54 (descr. ♂, Rafla).
Eucera amoena (WALKER, 1871): DE DALLA TORRE 1896: 224 (tax., world cat. (Eritrea, not Egypt)).
Eucera (Macrocera) amoena WALKER, 1871: FRIESE 1896: 22 (cat. Pal. Reg.), 177 (tax., orig. descr.).
Tetralonia amoena WALKER, 1871: INNES BEY 1911: 106 (Rafla, cat. Lord Collection).

D i s t r i b u t i o n : Rafla.

Tetralonia amoena, which is only known from Eritrea, is not mentioned in FRIESE (1909b, 1915), EARDLEY (1989) and EARDLEY & URBAN (2010).

Tetralonia blanda (WALKER, 1871)

- Tetralonia blanda* n.sp: WALKER 1871: 53 (descr. ♂, Harkeko).
Eucera blanda (WALKER, 1871): DE DALLA TORRE 1896: 227 (tax., world cat. (Eritrea, not Egypt)).
Eucera (Macrocera) blanda WALKER, 1871: FRIESE 1896: 22 (cat. Pal. Reg.), 178 (tax., orig. descr.).
Tetralonia blanda WALKER, 1871: INNES BEY 1911: 106 (Harkeko, cat. Lord Collection).
Tetralonia blanda WALKER, 1871: COCKERELL 1933a: 372 (tax.).

D i s t r i b u t i o n : Harkeko.

Tetralonia blanda, which is also recorded from Sudan (COCKERELL 1931: 403 as *Tetralonia gossypii* n.sp.), is not mentioned in FRIESE (1909b, 1915), EARDLEY (1989) and EARDLEY & URBAN (2010).

***Tetralonia vetusta* WALKER, 1871**

Tetralonia vetusta n.sp.: WALKER 1871: 53 (desc. ♀, Harkeko).

Eucera vetusta (WALKER, 1871): DE DALLA TORRE 1896: 250 (tax., world cat. (Eritrea, not Egypt)).

Eucera (Megacera) vetusta WALKER, 1871: FRIESE 1896: 22 (cat. Pal. Reg.), 184 (tax., orig. descr. Walker).

Tetralonia vetusta WALKER, 1871: INNES BEY 1911: 106 (Harkeko, cat. Lord Collection).

D i s t r i b u t i o n : Harkeko

Tetralonia vetusta, which is not mentioned in EARDLEY (1989) and EARDLEY & URBAN (2010), is only known from Eritrea.

***Tetraloniella friesei* (MEADE-WALDO, 1914)**

Tetralonia friesei n.nom.: MEADE-WALDO 1914b: 400 not seen

Tetraloniella friesei (MEADE-WALDO, 1914): EARDLEY 1989: 21 (tax.), 23 (key), 53 (fig. 97), 54 (fig. 108: map), 56 (tax., Gheleb).

Tetraloniella friesei (MEADE-WALDO, 1914): EARDLEY & URBAN 2010: 429 (cat. Afr. Reg. (Eritrea, not Ethiopia)).

**Tetralonia fulvicornis* n.sp.: FRIESE 1909b: 256 (cat. Africa), 259 (descr. ♀, Gheleb), 466 (cat. Africa).

Tetralonia fulvicornis FRIESE, 1909: FRIESE 1915: 272 (Gheleb).

Tetralonia fulvicornis FRIESE, 1909: RASMUSSEN & ASCHER 2008: 52 (cat. taxa described by Friese).

D i s t r i b u t i o n : Gheleb.

Tetraloniella friesei is only known from Eritrea. The records from Democratic Republic of Congo are probably misidentifications (EARDLEY 1989: 56). *Tetralonia fulvicornis* FRIESE, 1909 is a junior homonym of *Tetralonia fulvicornis* MORAWITZ, 1895.

***Tetraloniella graja* (EVERSMANN, 1852)**

Tetralonia graja (EVERSMANN, 1852): MAGRETTI 1906: 19 (Ailet).

D i s t r i b u t i o n : Ailet.

Tetralonia graja, which is known from the Palearctic Region, is not mentioned in FRIESE (1909b, 1915), EARDLEY (1989) and EARDLEY & URBAN (2010). The identification should be checked.

***Thyreomelecta bidentata* (KIRBY, 1899)**

**Crocisa valida* MORAWITZ, 1894: MAGRETTI 1906: 15 (Keren).

D i s t r i b u t i o n : Keren.

The record from Eritrea is not mentioned in FRIESE (1909b, 1915), MEYER (1921: 79), LIEFTINCK (1968: 42), RIGHTMYER & ENGEL (2003: 6) and EARDLEY & URBAN (2010). The identification should be checked.

***Thyreus abyssinicus* (RADOSKOWSKI, 1876)**

Crocisa abyssinica RADOSKOWSKI, 1876: MAGRETTI 1906: 10 (Sabarguma), 15 (Keren).

Crocisa abyssinica RADOSKOWSKI, 1876: MEYER 1921: 72 (tax., world cat.), 102 (tax., cat. Afr. Reg.), 104 (key ♂), 106 (key ♀), 107 (cat. Afr. Reg.), 114 (tax., desc. ♀ ♂, Eritrea without locality).

Thyreus abyssinicus (RADOSKOWSKI, 1873 (!)): EARDLEY 1991a: 4 (key), 31 (tax., descr. ♀ ♂, Eritrea (MEYER 1921)), 32 (figs 92-97), 33 (fig. 102: map, Eritrea missing).

Thyreus abyssinicus (RADOSKOWSKI, 1873 (!)): EARDLEY & URBAN 2010: 462 (cat. Afr. Reg. (Eritrea under Ethiopia)).

D i s t r i b u t i o n : Keren, Sabarguma.

Thyreus abyssinicus is widespread in the Afrotropical Region and is also recorded from Ethiopia and Sudan. The record from Eritrea is not mentioned in FRIESE (1909b: 310; 1915: 274).

***Thyreus alfkeni* (BRAUNS, 1909)**

Crocisa alfkeni BRAUNS, 1909: MEYER 1922: 184 (Eritrea without locality).

D i s t r i b u t i o n : no exact locality.

As *Thyreus alfkeni* is only known from South Africa (EARDLEY 1991: 23; EARDLEY & URBAN 2010: 463), the identification should be checked.

***Thyreus calceatus* (VACHAL, 1904)**

Crocisa calceata VACHAL, 1897: MEYER 1921: 72 (tax., world cat.), 103 (tax., cat. Afr. Reg.), 105 (key ♂), 106 (key ♀), 121 (cat. Afr. Reg.), 126 (tax., descr. ♀ ♂, Asmara).

Crocisa calceata VACHAL, 1897: MEYER 1922: 184 (tax., Eritrea without locality).

Thyreus calceatus (VACHAL, 1903 (!)): EARDLEY 1991: 5 (key as *T. calceata*), 41 (figs 131-136), 42 (tax., descr. ♀ ♂, Eritrea only in map), 44 (fig. 137: map).

Thyreus calceatus (VACHAL, 1903 (!)): EARDLEY & URBAN 2010: 464 (cat. Afr. Reg. (Eritrea under Ethiopia)).

D i s t r i b u t i o n : Asmara.

Thyreus calceatus, which is widely distributed in the Afrotropical Region, is also recorded from Ethiopia.

***Thyreus elegans* (MORAWITZ, 1877)**

Crocisa elegans MORAWITZ, 1877: MEYER 1922: 182 (tax., Assab).

Thyreus elegans (MORAWITZ, 1877): LIEFTINCK 1968: 6 (tab. 1: distr.), 12 (tax.), 26-7 (key), 58 (tax., Eritrea (MEYER 1922)), 60 (fig. 10), pl. 2 (figs 9-11).

D i s t r i b u t i o n : Assab.

Thyreus elegans, which is known from the Afrotropical (Eritrea, Sudan) and Palaearctic (e.g. Saudi Arabia) regions, is not mentioned in EARDLEY (1991) and EARDLEY & URBAN (2010).

***Thyreus erythraeensis* (MEYER, 1921)**

Crocisa erythraeensis n.sp.: MEYER 1921: 72 (tax., world cat.), 102 (tax., cat. Afr. Reg.), 105 (key ♂), 107 (key ♀), 121 (cat. Afr. Reg.), 122 (descr. ♀ ♂, Asmara).

Crocisa erythraeensis MEYER, 1921: ALFKEN 1934: 175 (tax., Asmara).

Thyreus erythraeensis (MEYER, 1921): LIEFTINCK 1968: 127 (tax., Asmara).

Thyreus erythraeensis (MEYER, 1921): EARDLEY 1991a: 46 (tax., not seen).

Thyreus erythraeensis (MEYER, 1921): EARDLEY & URBAN 2010: 465 (cat. Afr. Reg. (Eritrea, not Ethiopia)).

D i s t r i b u t i o n : Asmara.

Thyreus erythraeensis is also known from the Democratic Republic of Congo and Sudan.

? *Thyreus histrio* (FABRICIUS, 1775)

Crocisa histrio (FABRICIUS, 1775): CANNAVIELLO 1900: 297 (tax., Adi Caié, Adi Ugri).

Crocisa histrio (FABRICIUS, 1775): FRIESE 1909b: 301 (tax., cat. Africa), 302 (key ♂ as *C. nubica*), 305 (tax., descr. ♀ ♂, Eritrea without locality), 466 (cat. Africa).

Crocisa histrio (FABRICIUS, 1775): FRIESE 1915: 273 (Ghinda, Keren).

Thyreus histrio (FABRICIUS, 1775): EARDLEY & URBAN 2010: 465 (tax., cat. Afr. Reg. partim).

D i s t r i b u t i o n : Adi Caié, Ghinda, Keren, Adi Ugri.

Thyreus histrio is known from the Oriental and Palaearctic regions. The specimens mentioned in FRIESE (1909b, 1915) probably belong to *T. nubicus* (LEPELETIER DE SAINT-FARGEAU, 1841), because FRIESE (1909a) synonymized this species with *T. histrio*, but used the name *nubica* in the key.

***Thyreus hyalinatus* (VACHAL, 1904)**

Crocisa hyalinata VACHAL, 1903 (!): FRIESE 1909b: 301 (tax., cat. Africa), 303 (key ♀), 309 (tax., descr. ♀, Eritrea without locality), 466 (cat. Africa).

Crocisa hyalinata VACHAL, 1903 (!): FRIESE 1915: 274 (Asmara).

Crocisa hyalinata VACHAL, 1903 (!): MEYER 1921: 72 (tax., world cat.), 76 (tax., cat. Pal. Reg.), 78 (key ♀), 101 (cat. Pal. Reg., tax., descr. ♀ ♂, Eritrea without locality).

Crocisa hyalinata VACHAL, 1903 (!): MEYER 1922: 183 (tax., Assab).

Thyreus hyalinatus (VACHAL, 1903 (!)): LIEFTINCK 1968: 6 (tab. 1: distr.), 29-30 (key), 122 (tax., Assab), 123 (fig. 35), 138 (cat. Pal. Reg.).

Thyreus hyalinatus (VACHAL, 1903 (!)): EARDLEY 1991: 5 (key), 15 (figs 47-51), 19 (tax., descr. ♂, included only material examined by himself – Eritrea not mentioned), 22 (fig. 65: map).

Thyreus hyalinatus (VACHAL, 1903 (!)): EARDLEY & URBAN 2010: 465 (cat. Afr. Reg.).

D i s t r i b u t i o n : Asmara, Assab.

Thyreus hyalinatus is known from the Afrotropical (Djibouti, Eritrea, Sudan) and Palaearctic (e.g. Saudi Arabia, United Arab Emirates) regions.

***Thyreus meripes* (VACHAL, 1904)**

Crocisa meripes VACHAL, 1903 (!): FRIESE 1909b: 301 (tax., cat. Africa), 302 (key ♂), 303 (key ♀), 311 (tax., descr. ♀ ♂, Eritrea without locality), 466 (cat. Africa).

Crocisa meripes VACHAL, 1903 (!): FRIESE 1915: 273 (tax., Gheleb).

Crocisa meripes VACHAL, 1903 (!): MEYER 1921: 72 (tax., world cat.), 103 (tax., cat. Afr. Reg.), 105 (key ♀), 106 (key ♂), 121 (cat. Afr. Reg.), 127 (tax., Eritrea without locality).

Crocisa meripes VACHAL, 1903 (!): EARDLEY 1991: 4 (key), 5 (key), 44 (fig. 137 (map)).

Thyreus meripes (VACHAL, 1903): EARDLEY & URBAN 2010: 466 (cat. Afr. Reg. under Ethiopia).

D i s t r i b u t i o n : Gheleb.

Thyreus meripes is widespread in the Afrotropical Region, but is not recorded from the neighbour countries.

***Thyreus nigroventralis* (MEYER, 1921)**

Crocisa nigroventralis n.sp.: MEYER 1921: 72 (tax., world cat.), 103 (tax., cat. Afr. Reg.), 104 (key ♂), 130 (tax.), 131 (descr. ♂, Keren).

Thyreus nigroventris (!) (MEYER, 1921): EARDLEY 1991a: 46 (tax., not seen).

Thyreus nigroventralis (MEYER, 1921): EARDLEY & URBAN 2010: 466 (world cat. (Eritrea, not Ethiopia)).

D i s t r i b u t i o n : Keren.

Thyreus nigroventralis is only known from Eritrea.

***Thyreus nubicus* (LEPELETIER DE SAINT-FARGEAU, 1841)**

Crocisa nubica LEPELETIER DE SAINT-FARGEAU, 1841: MEYER 1922: 183 (tax., Asmara).

D i s t r i b u t i o n : Asmara.

Thyreus nubicus is also known from Cameroon, Nigeria, Sudan and Uganda. The record from Eritrea is not mentioned in EARDLEY & URBAN (2010: 465). See also *T. histrio* (FABRICIUS, 1775).

***Thyreus orbatus* (LEPELETIER DE SAINT-FARGEAU, 1841)**

Crocisa orbata LEPELETIER DE SAINT-FARGEAU, 1841: CANNAVIELLO 1900: 297 (tax., Adi Caié, Godofelassi).

D i s t r i b u t i o n : Adi Caié, Godofelassi.

Thyreus orbatus, which is not mentioned in FRIESE (1909b, 1915), EARDLEY (1991) and EARDLEY & URBAN (2010), is widespread in the Palaearctic Region. The identification should be checked.

***Thyreus scutellaris* (FABRICIUS, 1781)**

Crocisa scutellaris FABRICIUS, 1781: WALKER 1871: 52 (tax., Harkeko).

D i s t r i b u t i o n : Harkeko.

Thyreus scutellaris is known from the Palaearctic Region. The record from Eritrea is not mentioned in FRIESE (1909b, 1915), INNES BEY (1911: 106), MEYER (1921: 91), EARDLEY (1991: 35) and EARDLEY & URBAN (2010). The identification should be checked.

***Thyreus splendidulus* (LEPELETIER DE SAINT-FARGEAU, 1841)**

Crocisa splendidula LEPELETIER DE SAINT-FARGEAU, 1841: MAGRETTI 1906: 11 (Ghinda), 12 (Saganeiti).

D i s t r i b u t i o n : Ghinda, Saganeiti.

Thyreus splendidulus is also Cameroon, Democratic Republic of Congo, Equatorial Guinea, Ghana, Mali, Nigeria and Uganda. The record from Eritrea is not mentioned in FRIESE (1909b: 307; 1915), EARDLEY (1991: 35) and EARDLEY & URBAN (2010: 469).

***Thyreus* sp.**

Crocisa sejuncta DE SAUSSURE, 1891 (var.): MAGRETTI 1906: 15 (Keren).

D i s t r i b u t i o n : Keren.

The identification should be checked.

Subfamily N o m a d i n a e

Ammobates obscuratus MORAWITZ, 1894

Ammobates obscuratus MORAWITZ, 1894: MAGRETTI 1906: 10 (Sabarguma).

D i s t r i b u t i o n : Sabarguma.

Ammobates obscuratus, which is known from the Palaearctic Region, is not mentioned in FRIESE (1909b, 1915), EARDLEY & BROTHERS (1997) and EARDLEY & URBAN (2010). The identification should be checked.

Epeolus kristenseni FRIESE, 1915

Epeolus kristenseni FRIESE, 1915: BISCHOFF 1923: 600 (key, tax., Eritrea without locality).

D i s t r i b u t i o n : no exact locality.

Epeolus kristenseni is also known from Ethiopia. The record from Eritrea is not mentioned in EARDLEY (1991b: 717) and EARDLEY & URBAN (2010: 414).

Pasites appletoni (COCKERELL, 1910)

Pasites appletoni (COCKERELL, 1910): EARDLEY & BROTHERS 1997: 358 (tab. 1: tax.), 363 (tax., tree), 372 (key), 392 (tax., descr. ♀ ♂, Eritrea without locality), 393 (fig. 52), 394 (figs 53-56).

Pasites appletoni (COCKERELL, 1910): EARDLEY & URBAN 2010: 416 (cat. Afr. Reg. (Eritrea, not Ethiopia).

**Omachthes* (!) *abessinicus* n.sp.: FRIESE 1915: 276 (Eritrea without locality), 297 (descr. ♂, Eritrea without locality).

Morgania abessinica (FRIESE, 1915): Cockerell 1919: 190 (tax., cat. Africa).

Morgania (*Omachthes*) *abessinicus* (!) (FRIESE, 1915): BISCHOFF 1923: 589 (key, tax., Eritrea without locality).

Omachthes abessinicus FRIESE, 1915: EARDLEY & BROTHERS 1997: 297 (syn.).

Omachthes abessinicus FRIESE, 1915: RASMUSSEN & ASCHER 2008: 19 (cat. taxa described by Friese).

D i s t r i b u t i o n : no exact locality.

Pasites carnifex is widely distributed in the Afrotropical Region, but is not recorded from the neighbour countries.

Pasites carnifex (GERSTAECKER, 1869)

Omachthes carnifex GERSTAECKER, 1869: FRIESE 1909b: 437 (cat. Africa, tax., descr. ♂, Eritrea without locality), 468 (cat. Africa).

Pasites carnifex (GERSTAECKER, 1869): EARDLEY & BROTHERS 1997: 358 (tab. 1: tax.), 363 (tax., tree), 371 (key), 383 (tax., descr. ♀ ♂, fig. 33, Eritrea (FRIESE 1909)), 384 (figs 34-37).

Pasites carnifex (GERSTAECKER, 1869): EARDLEY & URBAN 2010: 418 (cat. Afr. Reg. (Eritrea, not Ethiopia).

**Omachthes* (!) *nigripes* n.sp.: FRIESE 1915: 276 (Eritrea without locality), 296 (descr. ♀ ♂, Eritrea without locality).

Morgania nigripes (FRIESE, 1915): COCKERELL 1919: 190 (tax., cat. Africa).

Morgania (*Omachthes*) *nigripes* (FRIESE, 1915): BISCHOFF 1923: 591 (key, tax., Eritrea without locality).

Omachthes nigripes FRIESE, 1915: EARDLEY & BROTHERS 1997: 383 (syn.).

Omachthes nigripes FRIESE, 1915: RASMUSSEN & ASCHER 2008: 77 (cat. taxa described by Friese).

D i s t r i b u t i o n : no exact locality.

Pasites carnifex is widely distributed in the Afrotropical Region, but is not recorded from the neighbour countries.

Subfamily Xylcopinae

***Braunsapis albitarsis* (FRIESE, 1924)**

Allodape albitarsis n.sp.: FRIESE 1924: 70 (cat. Africa), 72 (key), 73 (key), 81 (descr. ♀, Eritrea without locality).

Braunsapis albitarsis (FRIESE, 1924): MICHENER 1975: 86 (key ♂), 87 (key ♀), 96 (figs 96-98), 109 (tax., descr. ♀ ♂, Eritrea without locality), 111 (figs 109, 110).

Braunsapis albitarsis (FRIESE, 1924): MCGINLEY 1989: 15 (cat. immature Apoidea).

Allodape albitarsis FRIESE, 1924: RASMUSSEN & ASCHER 2008: 22 (cat. taxa described by Friese).

Braunsapis albitarsis (FRIESE, 1924): EARDLEY & URBAN 2010: 396 (cat. Afr. Reg. (Eritrea, not Ethiopia)).

D i s t r i b u t i o n : no exact locality.

Braunsapis albitarsis is also known from Kenya.

***Braunsapis bouyssoui* (VACHAL, 1904)**

Braunsapis bouyssoui (VACHAL, 1903 (!)): MICHENER 1975: 75 (biogeogr.), 85 (key ♂), 87 (key ♀), 96 (figs 56-61), 99 (tax., descr. ♀ ♂, Nefasit (Martini's Concession)), 100 (figs 67-75).

Braunsapis bouyssoui (VACHAL, 1903 (!)): EARDLEY & URBAN 2010: 397 (cat. Afr. Reg. (Eritrea, not Ethiopia)).

**Allodape citrifloris* n.sp.: COCKERELL 1935: 3 (descr. ♂, biol., Nefasit (Martini's Concession)).

Allodapula citrifloris (COCKERELL, 1935): COCKERELL 1936: 11 (tax.).

Allodape citrifloris COCKERELL, 1935: MICHENER 1975: 101 (syn.).

**Allodape comperei* n.sp.: COCKERELL 1935: 3 (descr. ♂, biol., Nefasit (Martini's Concession)).

Allodape comperei COCKERELL, 1935: MICHENER 1975: 101 (syn.).

D i s t r i b u t i o n : Nefasit (Martini's Concession).

Braunsapis bouyssoui, which is widespread in the Afrotropical Region, is unknown from the neighbour countries.

***Braunsapis facialis* (GERSTAECKER, 1858)**

Braunsapis facialis (GERSTAECKER, 1858): MICHENER 1975: 72 (fig. 1), 75 (biogeogr.), 83 (fig. 3), 85 (key ♂), 86 (keys ♀ ♂), 87 (tax., descr. ♀ ♂, Eritrea – not Ethiopia), 89 (figs 8-21), 91 (figs 22-37), 93 (tab. 3: tax.), 122 (fig. 8).

Braunsapis facialis (GERSTAECKER, 1858): EARDLEY & URBAN 2010: 398 (cat. Afr. Reg. (Eritrea under Ethiopia)).

**Allodape taita* STRAND, 1914: COCKERELL 1935: 2 (tax., descr. ♂, biol., Nefasit (Martini's Concession)).

D i s t r i b u t i o n : Nefasit (Martini's Concession).

Braunsapis facialis, which is widely distributed in the Afrotropical Region, is unknown from the neighbour countries.

***Braunsapis foveata* (SMITH, 1854)**

Braunsapis foveata (SMITH, 1854): MICHENER 1975: 75 (biogeogr.), 86 (key ♂), 87 (key ♀), 122 (figs 181-187), 128 (tax., descr. ♀ ♂, Eritrea without locality), 131 (figs 225 – 234).

Braunsapis foveata (SMITH, 1854): EARDLEY & URBAN 2010: 399 (cat. Afr. Reg. under Ethiopia).

**Allodape albicincta* n.sp.: FRIESE 1924: 70 (cat. Africa), 74 (key), 76 (descr. ♂, Eritrea without locality).

Allodape albicincta FRIESE, 1924: MICHENER 1975: 128 (syn. (type seen)).

Allodape albicincta FRIESE, 1924: RASMUSSEN & ASCHER 2008: 21 (cat. taxa described by Friese).

D i s t r i b u t i o n : no exact locality.

Braunsapis foveata, which is widespread in the Afrotropical Region, is unknown from the neighbour countries.

***Ceratina citriphila* COCKERELL, 1935**

Ceratina citriphila n.sp.: COCKERELL 1935: 1 (descr. ♂, biol., Nefasit (Martini's Concession)).

Pithitis citriphila (COCKERELL, 1935): DALY 1983a: 36 (key), 45 (fig. 34), 50 (fig. 50), 59 (tax., descr. ♀ ♂, Nefasit (Martini's Concession)).

Pithitis citriphila (COCKERELL, 1935): DALY 1983b: 500 (key), 501 (tax., Nefasit (Martini's Concession)).

Ceratina citriphila COCKERELL, 1935: DALY & ELSE 1988: 89 (tax., lectotype design., Nefasit (Martini's Concession)).

Ceratina citriphila COCKERELL, 1935: EARDLEY & URBAN 2010: 372 (cat. Afr. Reg. under Ethiopia).

**Ceratina tarsata* MORAWITZ, 1872: HIRASHIMA 1969: 655 (fig. 7).

D i s t r i b u t i o n : Nefasit (Martini's Concession).

Ceratina citriphila, which is known from the Afrotropical (Eritrea, Ethiopia, Somalia, Sudan) and Palaearctic (e.g. Yemen) regions, is not mentioned in COCKERELL (1937) and HIRASHIMA (1969).

***Ceratina tarsata* MORAWITZ, 1872**

Pithitis tarsata (MORAWITZ, 1872): HIRASHIMA 1969: 654 (tax.), 656 (tax., Eritrea without locality), 666 (fig. 16), 667 (fig. 20).

Pithitis tarsata (MORAWITZ, 1872): DALY 1983b: 500 (key), 503 (tax., Eritrea without locality).

Ceratina tarsata MORAWITZ, 1872: EARDLEY & URBAN 2010: 383 (cat. Afr. Reg. (Eritrea, not Ethiopia)).

D i s t r i b u t i o n : no exact locality.

Ceratina tarsata is known from the Afrotropical (Chad, Djibouti, Eritrea, Mali, Senegal, Sudan) and Palaearctic (e.g. Yemen) regions. HIRASHIMA's figure (1969: 655) is cited by DALY (1983a: 59) under *Ceratina citriphila* without comment.

***Compsomelissa boernerii* ALFKEN, 1924**

Compsomelissa borneri (!) ALFKEN, 1924: MICHENER 1975a: 223 (fig. 554), 225 (key), 226 (tax., descr. ♂, Agordat).

Compsomelissa borneri (!) ALFKEN, 1924: EARDLEY & URBAN 2010: 405 (cat. Afr. Reg. (Eritrea, not Ethiopia)).

**Exoneurula zavattarii* n.sp.: MICHENER 1966: 572, 573, 574

Exoneurula zavattarii MICHENER, 1966: MICHENER 1969b: 290, 293

Exoneurula zavattarii MICHENER, 1966: MICHENER 1975: 226 (syn.).

D i s t r i b u t i o n : Agordat.

Compsomelissa boernerii is recorded from the Afrotropical (Burkina Faso (LASSEN et al. 2016), Cameroon, Eritrea, Kenya, Nigeria, Senegal, Sudan) and Palaearctic (Yemen) regions.

***Macrogalea candida* (SMITH, 1879)**

Allodape candida SMITH, 1879: MAGRETTI 1884b: 627 (tax., biol., Keren).

Allodape candida SMITH, 1879: MAGRETTI 1885: 330 (Keren), 352 (cat. expedition).

Allodape candida SMITH, 1879: FRIESE 1915: 270 (tax., Keren).

Macrogalea candida (SMITH, 1879): COCKERELL 1930: 292 (typ. gen., descr. ♂, figs, Saganeiti (Coll. Gribodo)), 293 (biol.).

Macrogalea candida (SMITH, 1879): EARDLEY & URBAN 2010: 410 (cat. Afr. Reg. (Eritrea under Ethiopia)).

D i s t r i b u t i o n : Keren, Saganeiti.

Macrogalea candida, which is widely distributed in the Afrotropical Region, is also known from Ethiopia and Sudan.

***Xylocopa caffra* (LINNÉ, 1767)**

Xylocopa caffra (LINNÉ, 1767): CANNAVIELLO 1900: 297 (tax., Adi Ugri).

Xylocopa caffra LATREILLE (!): MAGRETTI 1906: 12 (Saganeiti).

Xylocopa (Koptorthosoma (!)) caffra (LINNÉ, 1767): DUSMET Y ALONSO 1924: 18 (tax., Asmara).

Xylocopa caffra (LINNÉ, 1767): EARDLEY 1987: 2 (cat. Afr. Reg. (Eritrea under Ethiopia)).

Xylocopa (Koptorthosoma) caffra (LINNÉ, 1767): VICIDOMINI & WHITMAN-MASCHERINI 2003: 113 (Eritrea without locality).

Xylocopa caffra (LINNÉ, 1767): EARDLEY & URBAN 2010: 340 (cat. Afr. Reg. (Eritrea under Ethiopia)).

D i s t r i b u t i o n : Adi Ugri, Asmara, Saganeiti.

Xylocopa caffra, which is widespread in the Afrotropical Region, is also known from Ethiopia.

***Xylocopa combusta* SMITH, 1854**

Xylocopa flavorufa var. *combusta* SMITH, 1854: MAGRETTI 1906: 12 (Debre Bizen).

Xylocopa var. *combusta* SMITH, 1854: FRIESE 1915: 271 (biol., Asmara, Gheleb).

Xylocopa (Koptorthosoma (!)) flavorufa var. *combusta* SMITH, 1854: DUSMET Y ALONSO 1924: 23 (tax., Asmara).

Xylocopa combusta SMITH, 1854: EARDLEY 1987: 4 (cat. Afr. Reg. (Eritrea under Ethiopia)).

Xylocopa combusta SMITH, 1854: EARDLEY & URBAN 2010: 345 (cat. Afr. Reg. (Eritrea under Ethiopia)).

D i s t r i b u t i o n : Asmara, Gheleb, Saganeiti.

Xylocopa combusta is widely distributed in the Afrotropical Region and is also recorded from Ethiopia.

***Xylocopa erythrina* GRIBODO, 1894**

Xylocopa erythrina n.sp.: GRIBODO 1894: 295 (Saganeiti), 296 (descr. ♀).

Xylocopa erythrina GRIBODO, 1894: DE DALLA TORRE 1896: 210 (world cat.).

Xylocopa (Xylocopa) erythrina GRIBODO, 1894: VACHAL 1899: 104 (key ♂, tax., Saganeiti).

Xylocopa erythrina GRIBODO, 1894: FRIESE 1909b: 216 (tax., cat. Africa), 219 (key ♀), 227 (tax., descr. ♀, Saganeiti), 465 (cat. Africa).

Xylocopa erythrina GRIBODO, 1894: FRIESE 1915: 271 (tax., Saganeiti).

Xylocopa (Perixylocopa) erythrina GRIBODO, 1894: HURD & MOURE 1963: 232 (typ. subgen.), 234 (tax., world cat.), 303 (world cat.).

Xylocopa (Xylomelissa) erythrina GRIBODO, 1894: MAA 1968: 106 (tax., lectotype design., Saganeiti).

- Xylocopa erythrina* Gribodo, 1894: EARDLEY 1987: 4 (cat. Afr. Reg. (Eritrea under Ethiopia)).
- Xylocopa (Perixylocopa) erythrina* GRIBODO, 1894: VICIDOMINI & WHITMAN-MASCHERINI 2003: 114 (Saganeiti (leg. Fossati)).
- Xylocopa erythrina* (!) GRIBODO, 1894: VICIDOMINI & WHITMAN-MASCHERINI 2003: 316 (Adi Ugri (leg. Andreini), Saganeiti).
- Xylocopa (Perixylocopa) erythrina* GRIBODO, 1894: VICIDOMINI 2006b: 123 (Adi Ugri (leg. Magretti), Asmara (leg. Cilli), Fort Ronch, Saganeiti), 125 (cat. Genova).
- Xylocopa erythrina* GRIBODO, 1894: EARDLEY & URBAN 2010: 346 (cat. Afr. Reg. (Eritrea under Ethiopia)).
- Xylocopa (Xylomelissa) erythrina erythrina* GRIBODO, 1894: PENATI & MARRIOTTI 2015: 49 (tax., Saganeiti, cat. Hymenoptera described by Gribodo).
- Xylocopa (Xylomelissa) erythrina* GRIBODO, 1894: PAULY et al. 2018: 8 (tax., biol., Tessenei), 9 (figs 7a-d), 10 (figs 8a-d).
- **Xylocopa fraudulenta* n.sp.: GRIBODO 1894: 295 (Saganeiti), 297 (descr. ♂).
- Xylocopa fraudulenta* GRIBODO, 1894: DE DALLA TORRE 1896: 211 (world cat.).
- Xylocopa (Xylocopa) fraudulenta* GRIBODO, 1894: VACHAL 1899: 158 (tax.). not seen
- Xylocopa fraudulenta* GRIBODO, 1894: FRIESE 1909b: 216 (tax., cat. Africa), 217 (key ♂), 220 (key ♀), 228 (tax., descr. ♂, Eritrea under Ethiopia), 465 (cat. Africa).
- Xylocopa fraudulenta* GRIBODO, 1894: FRIESE 1915: 271 (biol., Eritrea without locality under Ethiopia, Somaliland, Ostafrika bis Nyassa-See).
- Xylocopa (Acroxylocopa) fraudulenta* GRIBODO, 1894: HURD & MOURE 1963: 234 (typ. subgen.), 236 (tax., world cat.), 304 (world cat.).
- Xylocopa fraudulenta* GRIBODO, 1894: MAA 1968: 106 (lectotype design., syn.).
- Xylocopa fraudulenta* GRIBODO, 1894: PENATI & MARRIOTTI 2015: 55 (syn., Saganeiti, cat. Hymenoptera described by Gribodo).

D i s t r i b u t i o n : Adi Ugri, Saganeiti, Tessenei.

Xylocopa erythrina is widely distributed in the Afrotropical Region and is also known from Ethiopia.

***Xylocopa flavorufa* (DEGEER, 1778)**

- Xylocopa flavorufa* (DEGEER, 1778): MAGRETTI 1906: 13 (Adi Ugri).
- Xylocopa flavorufa* (DEGEER, 1778): FRIESE 1909b: 216 (tax., cat. Africa), 218 (key ♂), 220 (key ♀), 233 (tax., descr. ♀ ♂, Eritrea without locality), 465 (cat. Africa).
- Xylocopa flavorufa* (DEGEER, 1778): FRIESE 1915: 271 (Eritrea without locality).
- Xylocopa flavorufa* (DEGEER, 1778): EARDLEY 1987: 5 (cat. Afr. Reg. (Eritrea under Ethiopia)).
- Xylocopa flavorufa* (DEGEER, 1778): EARDLEY & URBAN 2010: 348 (cat. Afr. Reg. (Eritrea under Ethiopia)).

D i s t r i b u t i o n : Adi Ugri.

Xylocopa flavorufa is widespread in the Afrotropical Region and is also recorded from Ethiopia.

***Xylocopa gaullei* VACHAL, 1898**

- Xylocopa ? gaullei* VACHAL, 1898: MAGRETTI 1906: 11 (Ghinda).

D i s t r i b u t i o n : Ghinda.

Xylocopa gaullei is also recorded from Burundi, Ethiopia, Tanzania and Uganda. The record from Eritrea is not mentioned in FRIESE (1909b, 1915: 271), EARDLEY (1987: 6) and EARDLEY & URBAN (2010: 350).

***Xylocopa hottentota* SMITH, 1854**

- Xylocopa hottentota* SMITH, 1854: MAGRETTI 1906: 11 (Ghinda), 15 (Keren), 19 (Sabarguma).
Xylocopa hottentota SMITH, 1854: EARDLEY 1987: 6 (cat. Afr. Reg. (Eritrea under Ethiopia)).
Xylocopa (Nodula) hottentota SMITH, 1854: VICIDOMINI 2006b: 122 (Asmara (leg. Cilli), Keren (leg. Derchi, ??), Saati (leg. Magretti), Sabarguma (leg. Magretti), without locality (? Ghinda leg. Magretti)), 125 (cat. Genova).
Xylocopa hottentota SMITH, 1854: EARDLEY & URBAN 2010: 350 (cat. Afr. Reg. (Eritrea under Ethiopia)).
Xylocopa (Xylomelissa) hottentota SMITH, 1854: PAULY et al. 2018: 13 (tax., biol., Tessenei, Eritrea without locality), 14 (figs 12a, b), 15 (figs 12c, d, 13a-c).
 **Xylocopa carinata alpina* n.var.: FRIESE 1923: 6 (descr. ♀ ♂, Eritrea without locality, Ethiopia, Kenya).
Xylocopa (Xylomelissa) carinata alpina FRIESE, 1922 (!): HURD & MOURE 1963: 223 (tax., world cat.), 318 (world cat.).
Xylocopa hottentota alpina FRIESE, 1922 (!): RASMUSSEN & ASCHER 2008: 24 (cat. taxa described by Friese).
Xylocopa hottentota alpina FRIESE, 1922 (!): EARDLEY & URBAN 2010: 352 (cat. Afr. Reg.).
Xylocopa carinata var. *alpina* FRIESE, 1922 (!): PAULY et al. 2018: 13 (syn.).
 **Xylocopa tarsata* SMITH, 1854: MAGRETTI 1906: 13 (Adi Ugri).
Xylocopa tarsata SMITH, 1854: FRIESE 1909b: 216 (tax., cat. Africa), 217 (key ♂), 220 (key ♀) 221 (tax., descr. ♀ (?) ♂, Eritrea without locality), 465 (cat. Africa).
Xylocopa tarsata SMITH, 1854: FRIESE 1915: 270 (Eritrea without locality).
Xylocopa (Nodula) hottentota var. *tarsata* SMITH, 1854: VICIDOMINI 2006b: 123 (Godofelassi (leg. Magretti)), Saganeiti), 125 (cat. Genova).

D i s t r i b u t i o n : Adi Ugri, Asmara, Godofelassi, Keren, Saati, Saganeiti, Tessenei.
Xylocopa hottentota is widely distributed in the Afrotropical Region and is also recorded from Ethiopia and Sudan.

***Xylocopa inconstans* SMITH, 1874**

- Xylocopa inconstans* SMITH, 1874: MAGRETTI 1906: 10 (Sabarguma), 13 (Adi Ugri), 15 (Keren).
Xylocopa inconstans SMITH, 1874: FRIESE 1909b: 216 (tax., cat. Africa), 218 (key ♂), 219 (key ♂), 221 (key ♀), 253 (tax., descr. ♀ ♂, Keren), 466 (cat. Africa).
Xylocopa (Koptorthosoma (!)) inconstans SMITH, 1874: DUSMET Y ALONSO 1924: 25 (tax., Eritrea without locality).
Xylocopa inconstans SMITH, 1874: EARDLEY 1987: 8 (cat. Afr. Reg. (Eritrea under Ethiopia)).
Xylocopa (Xenoxylocopa) inconstans SMITH, 1874: PRIORE 1990: 36 (biol., Keren).
Xylocopa (Xenoxylocopa) inconstans SMITH, 1874: VICIDOMINI & Priore 1999: 130 (Keren).
Xylocopa (Xenoxylocopa) inconstans SMITH, 1874: VICIDOMINI 2002: 135 (Asmara).
Xylocopa (Xenoxylocopa) inconstans SMITH, 1874: VICIDOMINI & WHITMAN-MASCHERINI 2003: 114 (Fort Ranch (leg. Fossati), Saganeiti (leg. Fossati)), 116 (Adi Ugri (leg. Andreini), Mai Daro (? Andreini), Keren (leg. Andreini), Saganeiti (leg. Fossati, coll. Gribodo)).
Xylocopa (Xenoxylocopa) inconstans SMITH, 1874: VICIDOMINI 2005: 365 (Asmara, Betek, Saganeiti).
Xylocopa inconstans SMITH, 1874: EARDLEY & URBAN 2010: 353 (cat. Afr. Reg. (Eritrea under Ethiopia)).
Xylocopa (Xenoxylocopa) inconstans SMITH, 1874: PAULY et al. 2018: 16 (tax., biol., Keren), 17 (figs 14a-d).
 **Xylocopa inconstans flavocincta* n.var.: FRIESE 1909b: 253 (tax., descr. ♀ ♂, Keren).
Xylocopa inconstans var. *flavocincta* FRIESE, 1909: FRIESE 1915: 272 (Keren).
Xylocopa inconstans var. *flavocincta* FRIESE, 1909: LEVEQUE 1928: 21 (syn. of *X. chiyakensis* (COCKERELL, 1908)).

Xylocopa inconstans var. *flavocincta* FRIESE, 1909: HURD & MOURE 1963: 247 (syn. of *X. chiyakensis* (COCKERELL, 1908), world cat.), 304 (world cat.).

Xylocopa inconstans var. *flavocincta* FRIESE, 1909: RASMUSSEN & ASCHER 2008: 50 (cat. taxa described by Friese).

Xylocopa inconstans flavocincta FRIESE, 1909: MAWDSLEY 2017: 136 (syn.).

**Xylocopa chiyakensis* (COCKERELL, 1908): EARDLEY 1987: 3 (cat. Afr. Reg. (Eritrea under Ethiopia)).

Xylocopa chiyakensis (COCKERELL, 1908): EARDLEY & URBAN 2010: 344 (cat. Afr. Reg. (Eritrea under Ethiopia)).

D i s t r i b u t i o n : Adi Ugri, Asmara, Betek, Keren, Sabarguma, Saganeiti.

Xylocopa inconstans is widespread in the Afrotropical Region and is also known from Ethiopia and Sudan.

***Xylocopa lugubris* GERSTAECKER, 1857**

Xylocopa lugubris GERSTAECKER, 1857: FRIESE 1909b: 216 (tax., cat. Africa), 217 (key ♂), 221 (key ♀), 225 (tax., descr. ♂, Eritrea without locality), 465 (cat. Africa).

Xylocopa lugubris GERSTAECKER, 1857: EARDLEY 1987: 9 (cat. Afr. Reg. (Eritrea, not Ethiopia)).

Xylocopa lugubris GERSTAECKER, 1857: EARDLEY & URBAN 2010: 356 (cat. Afr. Reg. (Eritrea, not Ethiopia)).

**Xylocopa angolensis* SMITH, 1874: FRIESE 1915: 271 (tax., Eritrea without locality).

D i s t r i b u t i o n : without exact locality.

Xylocopa lugubris is widely distributed in the Afrotropical Region, but is not recorded from the neighbour countries. The identification should be checked.

***Xylocopa modesta* SMITH, 1854**

Xylocopa modesta SMITH, 1854: MAGRETTI 1906: 11 (Ghinda), 13 (Adi Ugri), 19 (Ailet).

D i s t r i b u t i o n : Adi Ugri, Ailet, Ghinda.

Xylocopa modesta SMITH, 1854 is widely distributed in the Afrotropical Region and is also recorded from Sudan. The record from Eritrea is not mentioned in FRIESE (1909b: 245), EARDLEY (1987: 9) and EARDLEY & URBAN (2010: 357).

***Xylocopa pubescens* SPINOLA, 1838**

Xylocopa pubescens (!) SPINOLA, 1838: EARDLEY 1987: 11 (cat. Afr. Reg. (Eritrea under Ethiopia)).

Xylocopa pubescens SPINOLA, 1838: EARDLEY & URBAN 2010: 361 (cat. Afr. Reg. (Eritrea under Ethiopia)).

**Xylocopa aestuans* (LINNAEUS, 1758): WALKER 1871: 58 (tax., Harkeko).

Xylocopa aestuans (LINNAEUS, 1758): MAGRETTI 1885: 280 (S Khor Langheb), 353 (cat. expedition).

Xylocopa aestuans (LINNAEUS, 1758): MAGRETTI 1906: 10 (biol., Massawa).

Xylocopa aestuans (LINNAEUS, 1758): FRIESE 1909b: 92 (biogeogr.), 216 (tax., cat. Africa), 218 (key ♂), 221 (key ♀), 242 (tax., descr. ♀ ♂, Asmara, Keren), 465 (cat. Africa).

Xylocopa aestuans (LINNAEUS, 1758): FRIESE 1915: 271 (Asmara, Keren).

Xylocopa (Koptorthosoma (!)) aestuans (LINNAEUS, 1758): DUSMET Y ALONSO 1924: 17 (tax., Assab).

Xylocopa aestuans (LINNAEUS, 1758): GUIGLIA 1931: 163 (tax., Beilul, Gahare).

Xylocopa aestuans (LINNAEUS, 1758): GUIGLIA 1932: 132 (tax., Agordat, Beilul, Gahare, Keren).

Xylocopa aestuans (LINNAEUS, 1758): GUIGLIA 1939a: 202 (tax., Keren).

Xylocopa (Mesotrichia) aestuans (LINNAEUS, 1758): GUIGLIA 1959: 317 (tax., Agordat, Gahare, Keren, footnote 1: orig. descr.).

Xylocopa (Koptortosoma) aestuans (LINNAEUS, 1758): VICIDOMINI & WHITMAN-MASCHERINI 2003: 112 (Fort Ranch, Keren, Tessenei).

Xylocopa (Koptortosoma) aestuans (LINNAEUS, 1758): VICIDOMINI 2005: 364 (Massawa).

D i s t r i b u t i o n : Agordat, Beilul, Fort Ranch, Harkeko, Keren, S Khor Langheb, Massawa, Tessenei.

Xylocopa pubescens is also known from Cameroon, Ethiopia, Kenya, Mozambique, Nigeria, Senegal, Somalia, Sudan and Tanzania The record from Harkeko by WALKER (1871) is not mentioned in INNES BEY (1911: 107).

***Xylocopa scioensis* GRIBODO, 1884**

**Xylocopa scioensis nigropunctata* n.var.: VICIDOMINI 2006a: 2 (descr., Saganeiti), 3 (tab. 1: tax.). unavailable name

Xylocopa scioensis var. *nigropunctata* VICIDOMINI, 2006: EARDLEY & URBAN 2010: 362 (cat. Afr. Reg. (Eritrea, not Ethiopia)).

**Xylocopa scioensis rufocollaris* n.var.: VICIDOMINI 2006a: 2 (descr., Debre Bizen), 3 (tab. 1: tax.). unavailable name

Xylocopa scioensis var. *rufocollaris* VICIDOMINI, 2006: EARDLEY & URBAN 2010: 362 (cat. Afr. Reg. (Eritrea, not Ethiopia)).

**Xylocopa flavorufa* var. *combusta* SMITH, 1854: MAGRETTI 1906: 12 (Debre Bizen, Saganeiti).

D i s t r i b u t i o n : Debre Bizen, Saganeiti.

Xylocopa scioensis is widespread in the Afrotropical Region and is also recorded from Ethiopia and Sudan. The new taxa described by VICIDOMINI (2006a) are unavailable names according to ICZN (1999: Art. 45.6.3).

***Xylocopa tessellata* MAA, 1970**

Xylocopa (Ctenoxylocopa) tessellata n.sp: MAA 1970: 724 (tax.), 728 (tax.), 729-30 (key), 730 (figs 4, 8), 734 (fig. 15), 742 (descr. ♀ ♂, Abula ("Abullo"), Gahare, Ghinda, Keren, Massawa), 745 (fig. 27), 747 (fig. 33), 749 (fig. 43).

Xylocopa tessellata MAA, 1970: EARDLEY 1987: 14 (cat. Afr. Reg. (Eritrea, not Ethiopia)).

Xylocopa (Ctenocylocopa) tessellata MAA, 1970: VICIDOMINI 2006b: 122 (Abula (Abullo), Gahare, Keren), 125 (cat. Genova).

Xylocopa tessellata MAA, 1970: EARDLEY & URBAN 2010: 366 (cat. Afr. Reg. (Eritrea, not Ethiopia)).

**Xylocopa oblonga* SMITH, 1874: MAGRETTI 1906: 13 (Adi Ugri).

**Xylocopa violacea* (LINNAEUS, 1758): WALKER 1871: 59 (tax., Harkeko, Massawa).

Xylocopa violacea SCOPOLI (!): INNES BEY 1911: 107 (without locality, cat. Lord Collection).

Xylocopa violacea (LINNAEUS, 1758): GUIGLIA 1931: 163 (tax., Abula, Gahare).

D i s t r i b u t i o n : Abula, Adi Ugri, Gahare, Ghinda, Harkeko, Keren, Massawa.

Xylocopa tessellata is also known from Djibouti and Sudan.

***Xylocopa torrida* (WESTWOOD, 1834)**

Xylocopa (Mesotricha) torrida (WESTWOOD, 1834): VICIDOMINI & WHITMAN-MASCHERINI 2003: 113 (Adi Ugri (leg. Magretti)).

Xylocopa (Mesotricha) torrida (WESTWOOD, 1834): VICIDOMINI 2005: 65 (Asmara (leg. Cilli)).

D i s t r i b u t i o n : Adi Ugri, Asmara.

Xylocopa torrida is widespread in the Afrotropical Region and is also recorded from Ethiopia. The record from Eritrea is not mentioned in EARDLEY & URBAN (2010: 367).

Xylocopa ustulata SMITH, 1854

Xylocopa (Ctenoxylocopa) ustulata SMITH, 1854: MAA 1970: 724 (tax.), 725 (lectotype design.), 727 (biogeogr.), 728 (biol.), 730 (figs 2, 6), 729 (key), 731 (key), 734 (fig. 12), 735 (tax., descr. ♀ ♂, Asmara, Ghinda, Keren, Saati), 745 (fig. 26), 747 (fig. 35), 749 (fig. 40).

Xylocopa ustulata SMITH, 1854: EARDLEY 1987: 14 (cat. Afr. Reg. (Eritrea under Ethiopia)).

Xylocopa ustulata SMITH, 1854: VICIDOMINI 2003: 271 (tax., Keren).

Xylocopa (Ctenoxylocopa) ustulata SMITH, 1854: VICIDOMINI & WHITMAN-MASCHERINI 2003: 115 (Keren (leg. Andreini)).

Xylocopa (Ctenoxylocopa) ustulata SMITH, 1854: VICIDOMINI 2005: 367 (Keren (leg. Andreini)).

Xylocopa ustulata SMITH, 1854: EARDLEY & URBAN 2010: 367 (cat. Afr. Reg.).

**Xylocopa hottentota* SMITH, 1854: FRIESE 1909: 92 (biogeogr.), 216 (cat. Afr. Reg.), 217 (key ♂), 219 (key ♀), 232 (tax., descr. ♀ ♂, Eritrea without locality), 465 (cat. Afr. Reg.).

Xylocopa (Xylocopa) hottentota SMITH, 1854: MAIDL 1912: 257 (fig. 5), 270 (tax., Massawa).

Xylocopa hottentota SMITH, 1854: FRIESE 1915: 271 (Eritrea without locality).

**Xylocopa taschenbergi* VACHAL, 1898: COCKERELL 1917: 460 (tax., Massawa).

D i s t r i b u t i o n : Asmara, Ghinda, Keren, Massawa, Saati.

Xylocopa ustulata is also known from Ethiopia, Gambia, Kenya, Mali, Niger, Nigeria, Somalia, Sudan and Madagascar.

Acknowledgements

I am greatly indebted to Wolfgang Brunnbauer (Naturhistorisches Museum Wien, Austria) and Manuela Vizek (Naturhistorisches Museum Wien, Austria) for their support in compilation of the catalogue.

Zusammenfassung

Derzeit sind in Eritrea 72 Arten der Familie Apidae nachgewiesen. Die Unterfamilie Apinae ist mit 48 Arten vertreten, die Unterfamilie Nomadinae mit vier Arten und die Unterfamilie Xylocopinae mit 22 Arten.

References

- ANDRÉ E. (1899-1903): Les Mutillides. — In: ANDRÉ E. (ed.), Species des Hyménoptères d'Europe & Algérie Tome 8: 1-64 (October 1899); 65-144 (April 1900); 145-224 (April 1901); 225-304 (October 1901); 305-384 (July 1902); 385 till end (January 1903).
- ALFKEN J.D. (1926): Beitrag zur Kenntnis der Bienenfauna von Ägypten. — *Senckenbergiana* **8/2**: 96-128.
- ALFKEN J.D. (1927): Zur Erforschung des Persischen Golfes (Beitrag Nr. 3). Apidae (Hym.). — *Entomologische Mitteilungen* **16/2**: 148-152.
- ALFKEN J.D. (1930): Die Apidae der Hamburgischen Südarabien-Expedition C. Rathjens – H. v. Wissmann. — *Mitteilungen der Münchner Entomologischen Gesellschaft* **20/1**: 44-48.
- ALFKEN J. (1934): Beitrag zur Kenntnis der Crocisa-Arten von Aegypten (Hymenoptera: Apoidea). — *Bulletin de la Société Entomologique d'Égypte* **18/1-2**: 164-176.
- COCKERELL T.D.A. (1919): Natal Bees. — *Annals of the Durban Museum* **2/4**: 189-196.

- BISCHOFF H. (1923): Zur Kenntnis afrikanischer Schmarotzerbienen. — Deutsche Entomologische Zeitschrift **1923/6**: 585-603.
- BROOKS R.W. (1988): Systematics and Phylogeny of the Anthophorine Bees (Hymenoptera: Anthophoridae, Anthophorini). — University of Kansas Science Bulletin **53/9**: 436-575.
- CANNAVIELLO E. (1900): Contributo alla Fauna Entomologica della colonia Eritrea. — Bullettino della Società Entomologica Italiana **32/3**: 289-308.
- COCKERELL T.D.A. (1917): New Records of Bees from Natal. — Annals of the Durban Museum **1/5**: 460-468.
- COCKERELL T.D.A. (1930a): A new African genus of Ceratinidae (Hymenoptera, Apoidea). — Revue de Zoologie et Botanique Africaines **18/3-4**: 291-293.
- COCKERELL T.D.A. (1930b): African bees of the family Xylocopidae principally from Liberia and the Belgian Congo. — Revue de Zoologie et Botanique Africaines **18/3-4**: 294-306.
- COCKERELL T.D.A. (1931): Some African Bees. — The Annals and Magazine of Natural History, 10th Series, **8/46**: 400-405.
- COCKERELL T.D.A. (1932): Descriptions and Records of Bees. — CXLIV. — The Annals and Magazine of Natural History, 10th Series, **9/53**: 447-458.
- COCKERELL T.D.A. (1933a): Descriptions and Records of Bees. — CXLII. — The Annals and Magazine of Natural History, 10th Series, **11/63**: 372-384.
- COCKERELL T.D.A. (1933b): Descriptions and Records of Bees. — CXLIII. — The Annals and Magazine of Natural History, 10th Series, **11/64**: 456-468.
- COCKERELL T.D.A. (1934): Some African Melopinine Bees. — Revue de Zoologie et Botanique Africaines **26/1**: 46-62.
- COCKERELL T.D.A. (1935): Bees from Eritrea, Northwest Africa. — American Museum Novitates **809**: 6 pp.
- COCKERELL T.D.A. (1937c): African bees of the genera *Ceratina*, *Halictus* and *Megachile*. — London (Trustees of the British Museum): XVI+254 pp.
- COCKERELL T.D.A. (1946): African Anthophorid Bees. — The Annals and Magazine of Natural History, 11th Series, **13/99**: 245-249.
- DALY H.V. (1983a): Taxonomy and ecology of Ceratinini of North Africa and the Iberian Peninsula (Hymenoptera: Apoidea). — Systematic Entomology **8/1**: 29-62.
- DALY H.V. (1983b): Bees of the genus *Pithitis* from Northeast Africa and the Arabian Peninsula (Hymenoptera: Apoidea). — Journal of the Kansas Entomological Society **56/4**: 499-505.
- DALY H.V. & G.R. ELSE (1988): Lectotype Designations for African Bees of the genus *Ceratina* Described by T.D.A. Cockerell. — The Pan-Pacific Entomologist **64/1**: 87-91.
- DALLA TORRE DE C.G. (1896): Catalogus hymenopterorum hucusque descriptorum et systematicus et synonymicus. Volumen 10: Apidae (Anthophila). — Leipzig (Wolfgang Engelmann): VIII+643 pp.
- DUSMET y ALONSO J.M. (1924): Las "*Xylocopa*" (Hymen. Apidae) en las Colecciones de Madrid. — Trabajos del Museo Nacional de Ciencias Naturales, Serie Zoológica, **49**: 58 pp.
- EARDLEY C.D. (1983): A taxonomic revision of the genus *Xylocopa* LATREILLE (Hymenoptera: Anthophoridae) in southern Africa. — Entomology Memoir, Department of Agriculture, Republic of South Africa, **58**: 20 pp.
- EARDLEY C.D. (1987): Catalogue of Apoidea (Hymenoptera) in Africa south of the Sahara Part 1: The genus *Xylocopa* LATREILLE (Anthophoridae). — Entomology Memoir, Department of Agriculture and Water Supply, Republic of South Africa, **70**: 67 pp.
- EARDLEY C.D. (1989): The Afrotropical species of *Eucara* FRIESE, *Tetralonia* SPINOLA and *Tetraloniella* ASHMEAD (Hymenoptera: Anthophoridae). — Entomology Memoir, Department of Agriculture and Water Supply, Republic of South Africa, **75**: 62 pp.

- EARDLEY C.D. (1991a): The Melectini in Sub-Saharan Africa (Hymenoptera: Anthophoridae). — Republic of South Africa, Department of Agricultural Development, Entomology Memoir **82**: 49 pp.
- EARDLEY C.D. (1991b): The genus *Epeolus* LATREILLE from sub-Saharan Africa (Hymenoptera: Anthophoridae). — Journal of Natural History **25**/3: 711-731.
- EARDLEY C.D. (2004): Taxonomic revision of the African stingless bees (Apoidea: Apidae: Apinae: Meliponini). — African Plant Protection **10**/2: 63-96.
- EARDLEY C.D. & R.W. BROOKS (1989): The genus *Anthophora* LATREILLE in southern Africa (Hymenoptera: Anthophoridae). — Entomology Memoir, Department of Agriculture and Water Supply, Republic of South Africa, **76**: 55 pp.
- EARDLEY C. & R. URBAN (2010): Catalogue of Afrotropical bees (Hymenoptera: Apoidea: Apiformes). — Zootaxa **2455**: 1-548.
- ENDERLEIN G. (1903): Neue und weniger bekannte africanische Xylocopen, besonders aus der Ausbeute des Freiherrn von Erlanger in Galla und Somali. — Berliner Entomologische Zeitschrift **48**/1-2: 45-64.
- FRIESE H. (1897): Die Bienen Europa's (Apidae europaeae) nach ihren Gattungen, Arten und Varietäten. Theil III. Solitäre Bienen. Gattung *Podalirius*. — Berlin (R. Friedländer & Sohn): VI+316 pp.
- FRIESE H. (1905): Neue *Anthophora*-Arten aus Afrika (Hym.). — Zeitschrift für Systematische Hymenopterologie und Dipterologie **5**/4: 233-242.
- FRIESE H. (1909a): Apidae. — In: SJÖSTEDT Y. (ed.), Wissenschaftliche Ergebnisse der Schwedischen Zoologischen Expedition nach dem Kilimandjaro, dem Meru und den umgebenden Massai-Steppen Deutsch-Ostafrikas 1905-1906 unter Leitung von Prof. Dr. Y. Sjöstedt **2**/8,6: 119-168.
- FRIESE H. (1909b): Die Bienen Afrikas nach dem Stande unserer heutigen Kenntnisse. — In: Zoologische und anthropologische Ergebnisse einer Forschungsreise im westlichen und zentralen Südafrika ausgeführt in den Jahren 1903 – 1905 mit Unterstützung der Kgl. Preussischen Akademie der Wissenschaften zu Berlin **2**/2: 83-476, pls. 9, 10.
- FRIESE H. (1911): Nachtrag zu "Bienen Afrikas". — Zoologische Jahrbücher, Abteilung für Systematik, Geographie und Biologie der Tiere **30**/6: 651-670.
- FRIESE H. (1915): Zur Bienenfauna von Abessinien (Hym.). — Deutsche Entomologische Zeitschrift **1915**/3: 265-298.
- FRIESE H. (1923): III. Nachtrag zu "Bienen Afrikas". — Zoologische Jahrbücher, Abteilung für Systematik, Geographie und Biologie der Tiere **46**/1: 1-42.
- FRIESE H. (1924): Über die Arten der Bienengattung *Allodape* in Afrika (Hym. Apid.). — Deutsche Entomologische Zeitschrift **1924**/1: 65-81.
- GRIBODO G. (1879): Note Imenotterologiche. — Annali del Museo Civico di Storia Naturale di Genova **14**: 325-347.
- GRIBODO G. (1894): Note imenotterologiche: Nota II. Nuovi generi e nuove specie di imenotteri antofili ed osservazione sopra alcune specie già conosciute (Continuazione). — Bullettino della Società Entomologica Italiana **26**/2: 262-314.
- GUIGLIA D. (1931a): Spedizione del Barone Raimondo Franchetti in Dancalia. Imenotteri aculeati. — Annali del Museo Civico di Storia Naturale Giacomo Doria **55**: 157-164.
- GUIGLIA D. (1932): Contributo alla conoscenza della fauna imenotterologica dell'Africa Orientale. — Memorie della Società Entomologica Italiana **10**: 118-136.
- GUIGLIA D. (1939a): Imenotteri aculeati raccolti in Eritrea e in Etiopia. — Memorie della Società Entomologica Italiana **17**/2: 188-202.
- GUIGLIA D. (1939b): Missione al Lago Tana diretta da G. Dainelli (1937) — Hymenoptera Aculeata. — Memorie della Società Entomologica Italiana **18**/1: 80-92.

- GUIGLIA D. (1940): Di alcune specie d'Imenotteri aculeati dell'Africa Orientale Italiana delle Collezioni del Museo Civico di Storia Naturale di Trieste. — Atti del Museo Civico di Storia Naturale Trieste **14/19**: 287-298.
- GUIGLIA D. (1956): Missione del Prof. Giuseppe Scortecci in Migiurtinia col contributo del Consiglio Nazionale delle Ricerche III: Hymenoptera. — Annali del Museo Civico di Storia Naturale "Giacomo Doria" **68**: 306-311, pl. 6.
- GUIGLIA D. (1959): Missione del Prof. Giuseppe Scortecci in Migiurtinia (Somalia Sett.) Hymenoptera: Vespidae, Pompilidae, Sphecidae, Apidae). — Atti della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale di Milano **98**, 4: 310-319.
- HEDICKE H. (1929): Bemerkungen über einige paläarktische und äthiopische *Anthophora*-Arten (Hym. Apid.). — Deutsche Entomologische Zeitschrift **1929/1**: 65-71.
- HIRASHIMA Y. (1969): Synopsis of the genus *Pithitis* KLUG of the world (Hymenoptera: Anthophoridae). — Pacific Insects **11/3-4**: 649-669.
- HURD P.D. Jr. & J.S. MOURE (1963): A Classification of the large Carpenter Bees (Xylocopini) (Hymenoptera: Apoidea). — University of California Publications in Entomology **29**: VI+365pp.
- INNES BEY W. (1911): Une liste d'Insectes recueillis probablement par J. LORD en Égypte et déterminés par F. WALKER. — Bulletin de la Société Entomologique d'Égypte **4/1**: 97-115.
- ICZN (1999): International Code of Zoological Nomenclature. — London (International Trust for Zoological Nomenclature): XXIX+306 pp.
- KOHL F.F. (1894): Zur Hymenopterenfauna Afrikas. — Annalen des k.k. Naturhistorischen Hofmuseums in Wien **9/3**: 279-350.
- LEVEQUE N. (1928): Carpenter Bees of the Genus *Mesotricha* obtained by the American Museum Congo Expedition, 1909 – 1915. — American Museum Novitates **300**: 1-23.
- LIEFTINCK M.A. (1968): A review of the Old World Species of *Thyreus* PANZER (= *Crocisa* JURINE) (Hym., Apoidea, Anthophoridae) Part 4: Palearctic Species. — Zoologische Verhandlungen **98**: 139 pp. 4 pls.
- MAA T.C. (1968): A revision of type specimens of Xylopine Bees in the Genova Museum collection. — Annali del Museo Civico di Storia Naturale "Giacomo Doria" **77**: 103-114.
- MADL M. (2018a): Notes on type-material of Meliponini (Hymenoptera: Apidae: Apinae) from the Horn of Africa deposited in the Natural History Museum Vienna, Austria. — Entomofauna **39/2** (3): 533-536.
- MADL M. (2018): A preliminary catalogue of the Hymenoptera (Insecta) of the Republic of Djibouti. — Linzer biologische Beiträge **50/2**: 907-967.
- MAGRETTI P. (1884a): Raccolte imenotterologiche nell'Africa orientale. Relazione preventiva. — Bullettino della Società Entomologica Italiana **15/4**: 241-253.
- MAGRETTI P. (1884b): Risultati di raccolte imenotterologiche nell'Africa orientale. — Annali del Museo Civico di Storia Naturale di Genova **21**: 523-636.
- MAGRETTI P. (1885): Nel Sudan Orientale. Ricordi d'un viaggio in Africa per studii zoologici. — Atti della Società Italiana di Scienze Naturali **26/3-4**: 257-355.
- MAGRETTI P. (1895): Esplorazioni del Giuba e dei suoi affluenti compiuta dal Cap. V. Bottego durante gli anni 1892-93 sotto gli auspicii della Società Geografica Italiana. Risultati Zoologici IX: Hymenoptera. — Annali del Museo Civico di Storia Naturale di Genova **35**: 149-173.
- MAGRETTI P. (1906): Materiali per la conoscenza della Fauna Eritrea. Imenotteri. Fam. Mutillidi. Con appunti del viaggio in Colonia. — Bullettino de la Società Entomologica Italiana **37/1**: 3-96.
- MAIDL F. (1912): Die Xylocopen (Holzbienen) des Wiener Hofmuseums. Ein Beitrag zu einer Monographie dieser Gattung. — Annalen des k.k. Naturhistorischen Hofmuseums in Wien **26/3-4**: 249-330, pls 3, 4.

- MCGINLEY R.J. (1989): A Catalog and Review of Immature Apoidea (Hymenoptera). — *Smithsonian Contribution to Zoology* **494**: 1-24.
- MEYER R. (1921): Apidae – Nomadinae I. Gattung *Crocisa* JUR. — *Archiv für Naturgeschichte* **86/Abteilung A**, 1: 67-178.
- MEYER R. (1922): Nachtrag I. Zur Bienen-Gattung *Crocisa* JUR. Die *Crocisa*-Arten des Ungarischen National-Museums. — *Annales Historico-Naturales Musei Nationalis Hungarici* **19**: 182-186.
- MICHENER C.D. (1971a): Biologies of African Allodapine Bees. — *Bulletin of the American Museum of Natural History* **145/3**: 219-302.
- MICHENER C.D. (1966): A New Genus of Ceratinine Bees from the Red Sea Area. — *Journal of the Kansas Entomological Society* **39/4**: 572-576.
- MICHENER C.D. (1969): African genera of allodapine bees (Hymenoptera: Anthophoridae: Ceratinini). — *Journal of the Kansas Entomological Society* **42/3**: 289-293.
- MICHENER, C.D. 1971b: The genus *Macrogalea*. — *Entomological Essays to Commemorate the Retirement of Professor K. Yasumatsu*: 61-71.
- MICHENER C.D. (1975): A Taxonomic Study of African Allodapine Bees (Hymenoptera, Anthophoridae, Ceratinini). — *Bulletin of the American Museum of Natural History* **155/2**: 67-240.
- MICHENER C.D. (1997): Genus-Group names of Bees and Supplemental Family-Group Names. — *Scientific Papers, Natural History Museum, The University of Kansas*, **1**: 1-81.
- MINCKLEY R.L. (1998): A Cladistic Analysis and Classification of the Subgenera and Genera of the Large Carpenter Bees, Tribe Xylocopini (Hymenoptera: Apidae). — *Scientific Papers, Natural History Museum, The University of Kansas*, **9**: 1-47.
- MOURE J.S. (1961): A Preliminary Supra-specific Classification of the Old World Meliponine Bees (Hym., Apoidea). — *Studia Entomologica* **4/1-4**: 181-242.
- PAULY A. & Z.A. HORA (2013): Apini and Meliponini from Ethiopia (Hymenoptera: Apoidea: Apidae: Apinae). — *Belgian Journal of Entomology* **16**: 1-35.
- PAULY A., HORA Z.A., WAYESSA D.N. & K. AMBERBIR (2018): The *Xylocopa* LATREILLE, 1802, of Ethiopia (Hymenoptera: Apidae). — *Belgian Journal of Entomology* **62**: 1-30.
- PENATI F. & A. MARIOTTI (2015): Catalog of Hymenoptera described by G. GRIBODO (1846-1924) (Insecta). — *Zootaxa* **3929**: 1-183.
- PRIESNER H. (1957): A Review of the Anthophora-Species of Egypt (Hymenoptera: Apidae). — *Bulletin de La Société Entomologique d'Égypte* **61/1**: 1-115.
- PRIORE R. (1990): La collezione degli Apoidea (Hymenoptera) dell'Istituto di Entomologia agraria di Portici – X – *Coelioxys* LATR. (Megachilidae), *Xylocopa* LATR., *Ceratina* LATR. (Xylocopidae). — *Bollettino del Laboratorio di Entomologia Agraria "Filippo Silvestri"* **46**: 31-44.
- RASMUSSEN C. & J.S. ASCHER (2008): Heinrich FRIESE (1860-1948): names proposed and notes on a pioneer melittologist (Hymenoptera, Anthophila). — *Zootaxa* **1833**: 1-118.
- RIGHTMYER M.G. & M. ENGEL (2003): A New Palearctic Genus of Melectine Bees (Hymenoptera: Apidae). — *American Museum Novitates* **3392**: 1-22.
- SANDHOUSE G.A. (1943): The Type Species of the Genera and Subgenera of Bees. — *Proceedings of the United States National Museum* **92/3156**: 519-619.
- VACHAL M.J. (1899): Essai d'une révision synoptique des espèces européennes et africaines du genre *Xylocopa* LATR. — *Miscellanea Entomologica* **7/78**: 89-112; 10-11: 145-160.
- VACHAL M.J. (1900): Essai d'une révision synoptique des espèces européennes et africaines du genre *Xylocopa* LATR. (fin). — *Miscellanea Entomologica* **8/8-9**: 106-108.
- VACHAL M.J. (1910): Collections recueillis par M. le Baron Maurice de Rothschild dans l'Afrique Orientale. — *Bulletin du Muséum d'Histoire Naturelle* **15/8**: 529-534.

- VICIDOMINI S. (2002): Xylocopini (Hymenoptera, Apidae, Xylocopinae) presenti nelle collezioni entomologiche italiane: il Museo Civico di Storia Naturale di Trieste. II. — Atti del Museo Civico di Storia Naturale di Trieste **49**: 133-136.
- VICIDOMINI S. (2003): Systematic and distribution of Xylocopini (Hymenoptera: Apidae: Xylocopinae): new and rare records for Africa (Part III) and description of two new taxa. — Atti del Museo Civico di Storia Naturale di Trieste **50**: 267-283.
- VICIDOMINI S. (2005): Xylocopini (Hymenoptera: Apidae: Xylocopinae) presenti nelle collezioni entomologiche italiane: l'Istituto di Zoologia (= Ex-I.N.E.) ed il Museo Zoologico di Roma (Parte III); il Servizio Fitosanitario Regionale del Piemonte, Torino; il Museo Civico di Storia Naturale, Montebelluna; il Museo di Scienze Naturali, Storia ed Arte, L'Aquila). — Annali del Museo Civico de Rovereto, Sezione Archeologia, Storia, Scienze Naturali **20**: 357-373.
- VICIDOMINI S. (2006a): Sistematica e distribuzione degli Xylocopini (Hymenoptera: Apidae): nuove e rare segnalazioni per l'Africa (parte V). Rassegna delle forme cromatiche di *Xylocopa caffra*, *X. scioensis* e *X. nigrita*. — Il Naturalista Campano **2006/2**: 8 pp.
- VICIDOMINI S. (2006b): Xylocopini (Hymenoptera: Apidae: Xylocopinae) presenti nelle collezioni entomologiche italiane: il Museo civico di Storia naturale G. Doria (Genova). — Pianura – Science e Storia dell'Ambiente Padano **20**: 121-125.
- VICIDOMINI S. & R. PRIORE (1999): Xylocopini presenti nelle collezioni entomologiche italiane: l'Istituto di Entomologia Agraria Filippo Silvestri, Portici. — Bollettino del Laboratorio di Entomologia Agraria "Filippo Silvestri" **55**: 127-134.
- VICIDOMINI S. & S. WHITHMAN-MASCHERINI (2003): Xylocopini (Hymenoptera: Apidae: Xylocopinae) presenti nelle collezioni entomologiche italiane: il Museo civico di Storia naturale, Milano (II) e il Museo zoologico La Specola, Firenze (III). — Pianura – Science e Storia dell'Ambiente Padano **17**: 111-118.
- WALKER F. (1871): A List of Hymenoptera collected by J.K. Lord, Esq. in Egypt, in the neighbourhood of the Red Sea, and in Arabia. With descriptions of the new species. — London (E.W. Janson): IV+59 pp.

Author's address: Michael MADL
 Naturhistorisches Museum Wien
 2. Zoologische Abteilung
 Burgring 7
 A-1010 Vienna, Austria
 E-Mail: michael.madl@nhm-wien.ac.at