

Linzer biol. Beitr.	53/1	101-110	August 2021
---------------------	------	---------	-------------

Notes on the Sphecidae (Hymenoptera) of Gabon

Michael MADL

A b s t r a c t : In Gabon the family Sphecidae is represented by 15 species and one subspecies belonging to the genera *Chlorion* LATREILLE, 1802 (one species), *Chalybion* DAHLBOM, 1843 (three species), *Isodontia* PATTON, 1880 (two species), *Prionyx* VANDER LINDEN, 1827 (two species), *Sceliphron* KLUG, 1801 (two species and one subspecies) and *Sphex* LINNAEUS, 1758 (five species). New records are reported from Gabon and Republic of Congo.

K e y w o r d s : Sphecidae, Chlorioninae, Sceliphrinae, Sphecinae, new records, Gabon, Republic of Congo

Introduction

FAIRMAIRE (1858) published the hymenopterological results of the entomological expedition to Gabon carried out by H.C. Deyrolle in 1856 and 1857 (THOMSON 1858) and listed four species of the family Sphecidae including a new one. COSTA (1866), KOHL (1890, 1894) and KIRBY (1896) added further data to the Sphecid fauna of Gabon. At the end of the 19th century six species were known from Gabon. In the 20th century, the knowledge of the Gabunese Sphecidae was increased by the papers of SCHULZ (1911), KOHL (1918), BERLAND (1926a, b, 1927, 1928, 1956), LECLERCQ (1955, 1969), VECHT & BREUGEL (1968) and HENSEN (1988).

Until now 15 species and one subspecies of the family Sphecidae have been recorded from Gabon. They belong to the genera *Chlorion* LATREILLE, 1802 (one species), *Chalybion* DAHLBOM, 1843 (three species), *Isodontia* PATTON, 1880 (two species), *Prionyx* VANDER LINDEN, 1827 (two species), *Sceliphron* KLUG, 1801 (two species and one subspecies) and *Sphex* LINNAEUS, 1758 (five species). Currently the genera *Ammophila* KIRBY, 1798 and *Podalonia* SPINOLA, 1853, which are widely distributed in the Afrotropical region, are not recorded from Gabon.

This paper is based on the material of the Natural History Museum Vienna, which was mainly collected at the end of the 19th century. Few specimens have been collected by the author during a field trip to Gabon in 2019. The material contains specimens of *Chalybion bocandei* (SPINOLA, 1851), *Chalybion clypearatum* (KOHL, 1918), *Sceliphron fossuliferum complex* KOHL, 1918, *Sceliphron spirifex* (LINNAEUS, 1758) and *Sphex nigrohirtus* KOHL, 1895.

Annotated catalogue and material

In the 19th century the geographic term "Gabon" was used to describe only the land area of the Bay of Gabon (French: Estuaire du Gabon), which is part of the province of Estuaire in today's Republic of Gabon. The specimens were probably collected in the surroundings of Libreville. Anyway, it is necessary to check the life history of the collector.

Many specimens are labelled "Chûtes de Samlia, Riv. N. Gamio, leg. Mocquereys", of which several misspellings have been published. The locality has been assigned to Gabon or Sierra Leone (e.g. HENSEN 1988) in the scientific literature. This geographical and orthographical problem was first solved by RÉGIMBART (1895: 126) and later mentioned by SCHOUTEDEN (1918: 127) and RASMUSSEN & ASCHER (2015: 299). The correct spelling is "Chûtes de Samba, Rivière Ngounié" and the locality is situated near Sindara in the province of Ngounié (Gabon).

Subfamily Chloriontinae

Chlorion maxillosum ciliatum (FABRICIUS, 1787)

Chlorion maxillosum ciliatum (FABRICIUS, 1787): BOHART & MENKE 1976: 90 (world catalogue).

**Chlorion xanthoceros maxillare* (PALISOT DE BEAUVOIS, 1805): LECLERCQ 1955: 42 (taxonomy (note below fig. 1B)), 44 (footnote 1: Lambaréné).

Sphex (Chlorion) xanthocerus var. *maxillaris* PALISOT DE BEAUVOIS, 1805: BERLAND 1956: 1163 (taxonomy, description ♂, Libreville), 1165 (key to varieties of *Sphex xanthocerus*).

**Chlorion xanthoceros instabile* (SMITH, 1856): LECLERCQ 1955: 45 (footnote 1: Lambaréné).

**Sphex (Chlorion) xanthocerus* var. *subcyaneum* GERSTAECKER, 1857: BERLAND 1956: 1164 (taxonomy, description ♂ ♀, Libreville), 1165 (key to varieties of *Sphex xanthocerus*).

D i s t r i b u t i o n : Estuaire: Libreville – Moyen-Ogooué: Lambaréné.

Chlorion maxillosum ciliatum, which is widely distributed in the Afrotropical region, is also recorded from the Republic of Congo.

Subfamily Sceliphrinae

Chalybion bocandei (SPINOLA, 1851)

Chalybion bocandei (SPINOLA, 1851): HENSEN 1988: 37 (key), 40 (figs. 109, 110), 43 (key to subspecies, taxonomy, description ♂ ♀, "Chûtes de Samlia R., N. Camie" = Chûtes de Samba (Rivière Ngounié) near Sindara (Gabon – not Sierra Leone), Fernand-Vaz (Gabon – not Congo), Gabon without exact locality).

**Sceliphron (Chalybion) fuscipenne* (SMITH, 1856): KOHL 1918: 21 (key ♀), 26-7 (key ♂), 72 (taxonomy, description ♂ ♀, Gabon without exact locality), 139 (biogeography, world catalogue).

Sceliphron (Chalybion) fuscipenne (SMITH, 1856): ARNOLD 1928a: 235 (keys ♀, ♂), 241 (taxonomy, description ♂ ♀, Gabon without exact locality).

Chalybion fuscipenne (SMITH, 1856): LECLERCQ 1955: 46 (Chûtes de Samlia = Chûtes de Samba (Rivière Ngounié)).

M a t e r i a l e x a m i n e d : Gabon without exact locality 1♂ 1♀.

D i s t r i b u t i o n : Ogooué-Maritime: Fernand-Vaz – Ngounié: Chûtes de Samba (Rivière Ngounié) near Sindara – no exact locality (KOHL 1918, ARNOLD 1928a, HENSEN 1988, this paper).

Chalybion bocandei is also known from the neighbour countries Equatorial Guinea (Mbini) and Cameroon.

***Chalybion clypeatum* (FAIRMAIRE, 1858)**

Pelopoeus clypeatus n.sp.: FAIRMAIRE 1858: 264 (description, Gabon without exact locality).
Sceliphron clypeatus (FAIRMAIRE, 1858): DALLA TORRE 1897: 382 (taxonomy, world catalogue).
Pelopoeus clypeatus FAIRMAIRE, 1858: KOHL 1918: 80 (synonym of *Sceliphron (Hemicalybion) eckloni* (DAHLBOM, 1845) based only on description).
Chalybion (Hemicalybion) clypeatum (KOHL, 1918): BOHART & MENKE 1976: 103 (taxonomy, world catalogue).
**Sceliphron (Hemicalybion) brachystylus* KOHL, 1888: KOHL 1918: 83 (taxonomy, description ♂ ♀, Gabon without data), 139 (biogeography, world catalogue).

M a t e r i a l e x a m i n e d : Gabon without data 1♂.

D i s t r i b u t i o n : Estuaire: no exact locality (FAIRMAIRE 1858) – without exact locality (KOHL 1918, this paper).

Chalybion clypeatum, which is widely distributed in the Afrotropical region, is also recorded from the Republic of Congo.

***Chalybion schulthessirechbergi* (KOHL, 1918)**

Chalybion schulthessirechbergi (KOHL, 1918): HENSEN 1988: 37 (key), 44 (figs. 120-122), 46 (taxonomy, description ♂ ♀, Lambaréne, between Lambaréne and sea).

D i s t r i b u t i o n : Moyen-Ogooué: Lambaréne – between Lambaréne and sea (exact geographic location unknown).

Chalybion schulthessirechbergi is also known from Cameroon, Central African Republic, Democratic Republic of Congo, Ivory Coast, Liberia, Namibia and Uganda.

***Sceliphron fossuliferum* (GRIBODO, 1895)**

Sceliphron (Sceliphron) fossuliferum (GRIBODO, 1895): BOHART & MENKE 1976: 105 (world catalogue).

**Sceliphron (Sceliphron) quartinae quartinae* GRIBODO, 1884: LECLERCQ 1955: 59 (taxonomy, "Chûtes de Samlia, riv. N'Gamie" = Chûtes de Samba (Rivière Ngounié) near Sindara, Gabon without exact locality (1♂ 19. 3. 1892 without collector (note by Madl: leg. H. Brauns – see dates in KOHL 1894: 342 *Philanthus triangulum*)).

M a t e r i a l e x a m i n e d : Ogooué-Ivindo Province: Monts de Belinga, Mbondo (0°51'30"N, 13°11'32"E), ca. 540 m, 1♀ 02.02.2019 leg. M. Madl.

D i s t r i b u t i o n : Estuaire: no exact locality (LECLERCQ 1955) – Ngounié: Chûtes de Samba (Rivière Ngounié) near Sindara – Ogooué-Ivindo: Mbondo.

Sceliphron fossuliferum, which is widely distributed in the Afrotropical region, is also recorded from the Republic of Congo.

***Sceliphron fossuliferum complex* KOHL, 1918**

Sceliphron (Pelopoeus) complex n.sp.: KOHL 1918: 25-5 (key ♀, Gabon – not West Africa), 105 (taxonomy), 107 (description ♀, "Choutes de Samlia River, N'Gamio" = Chûtes de Samba (Rivière Ngounié) near Sindara), 139 (biogeography, world catalogue).

Sceliphron (Pelopoeus) complex KOHL, 1918: ARNOLD 1928a: 236 (key), 246 (taxonomy, not examined).

Sceliphron complex KOHL, 1918: ARNOLD 1930: 16 (catalogue Afrotropical Region).

Sceliphron (Pelopoeus) complex KOHL, 1918: LECLERCQ 1955: 59 (synonym of *Sceliphron (Pelopoeus) quartinae voeltzkovii* (!) KOHL, 1909, footnote: type not in Institut Royal des Sciences Naturelles de Belgique, probably in Naturhistorisches Museum Wien).

Sceliphron (Sceliphron) fossuliferum complex KOHL, 1918: VECHT & BREUGEL 1968: 215 (key to subspecies), 216 (taxonomy, Chûtes de Samlia, Riv. N. Gamio = Chûtes de Samba (Rivière Ngounié) near Sindara).

Sceliphron (Sceliphron) fossuliferum ssp. *complex* KOHL, 1918: BOHART & MENKE 1976: 105 (world catalogue).

Sceliphron fossuliferum complex KOHL, 1918: DOLFFUSS 1989: 12 (type catalogue Natural History Museum Vienna), 17 (type catalogue Natural History Museum Vienna).

M a t e r i a l e x a m i n e d : Ngounié: Chûtes de Samlia, Riv. N. Gamio 1♀ leg. Mocquereys (Holotype) – Ngounié Province: RP 403 Lebamba – Mbigou, Dibwangu (2°06'43"S, 11°34'49"E), ca. 400, 1♀, 31.02.2019, leg. M. Madl.

D i s t r i b u t i o n : Ngounié: Chûtes de Samba (Rivière Ngounié) near Sindara, Dibwangu.

Sceliphron fossuliferum complex is also known from Equatorial Guinea (Mbini), Republic of Guinea and Rwanda.

***Sceliphron spirifex* (LINNAEUS, 1758)**

Pelopoeus spirifer (!) (LINNAEUS, 1758): FAIRMAIRE 1858: 265 (Gabon without exact locality).

Pelopoeus spirifex (LINNAEUS, 1758): COSTA 1866: 24 (Gaban = Gabon without further data).

Sceliphron spirifex (LINNAEUS, 1758): KOHL 1894: 342 (Gabon without exact locality).

Pelopoeus spirifex (LINNAEUS, 1758): KIRBY 1896: 265 (Gabon without exact locality (only Ogooué river)).

Sceliphron (Pelopoeus) spirifex (LINNAEUS, 1758): KOHL 1918: 5 (fig. 9), 7 (taxonomy), 8 (taxonomy), 9 (figs. 12, 13), 12 (biogeography), 15 (fig. 20), 16 (key ♀), 19 (key ♀), 25 (key ♂), 86 (taxonomy, description ♂ ♀, Gabon without exact locality), 87 (fig. 46), 139 (biogeography, world catalogue), 144 (biology), 145 (biology), 146 (biology), 147 (biology), 148 (biology), 149 (biology), 150 (biology), 162 (biology), 163 (fig. 65: larva), 167 (biology), pl. 2 (figs. 11 (?), 13).

Sceliphron (Sceliphron) spirifex (LINNAEUS, 1758): LECLERCQ 1955: 60 (taxonomy, "Chûtes de Samlia, riv. N'Gamie" = Chûtes de Samba (Rivière Ngounié) near Sindara), 63 (footnote: Libreville).

Sceliphron (Sceliphron) spirifex (LINNAEUS, 1758): VECHT & BREUGEL 1968: 231 (map), 232 (key), 234 (fig. 20), 235 (fig. 21), 236 (fig. 22), 216 (taxonomy, localities see KOHL (1918) and LECLERCQ (1955)), 240 (fig. 23).

Sceliphron spirifex (LINNAEUS, 1758): LECLERCQ 1969: 1048 (taxonomy, Mbila – Gabon, not Republic of Congo).

Sceliphron (Sceliphron) spirifex (LINNAEUS, 1758): BOHART & MENKE 1976: 105 (world catalogue).

M a t e r i a l e x a m i n e d : Moyen-Ogooué: Ndjolé, Hotel Kedyann, 1♀, 04.02.2019, leg. M. Madl.

D i s t r i b u t i o n : Estuaire: Libreville, no exact locality (FAIRMAIRE 1858, COSTA 1866, KOHL 1894, 1918) – Moyen-Ogooué: Ndjolé – Ngounié: Chûtes de Samba (Rivière Ngounié) near Sindara, Mbila – no exact locality (KIRBY 1896).

Sceliphron spirifex, which is widely distributed in the Afrotropical and southern Palearctic regions, is also known from Cameroon and the Republic of Congo.

Subfamily Sphecinae

***Isodontia pelopoeiformis* (DAHLBOM, 1845)**

Sphex (Isodontia) pelopoeiformis (!) DAHLBOM, 1845: BERLAND 1926c: 282 (taxonomy, Lambaréné).

Sphex (Isodontia) pelopoeiformis DAHLBOM, 1845: BERLAND 1956: 1171 (taxonomy, description ♂ ♀, Lambaréné, Talogoriga = Talagouga), 1180 (key).

Isodontia pelopoeiformis (DAHLBOM, 1845): BOHART & MENKE 1976: 121 (tab. 7: prey), 124 (world catalogue), 125 (fig. Q).

Distribution: Moyen-Ogooué: Lambaréné, Talagouga.

Isodontia pelopoeiformis, which is widely distributed in the Afrotropical and southern Palaearctic regions, is also recorded from Cameroon and the Republic of Congo.

***Isodontia stanleyi* (KOHL, 1890)**

Sphex (Isodontia) stanleyi (KOHL, 1890): BERLAND 1926b: 282 (Lambaréné).

Sphex (Isodontia) stanleyi (KOHL, 1890): BERLAND 1956: 1172 (taxonomy, description ♂ ♀, Lambaréné, Gabon without exact locality), 1180 (key).

Isodontia stanleyi (KOHL, 1890): BOHART & MENKE 1976: 124 (world catalogue).

Distribution: Estuaire: no exact locality (BERLAND 1956) – Moyen-Ogooué: Lambaréné.

Isodontia stanleyi, which is widely distributed in the Afrotropical region, is not reported from the neighbour countries.

***Prionyx kirbii marginatus* (SMITH, 1856)**

Prionyx kirbyi (!) ssp. *marginatus* (SMITH, 1856): BOHART & MENKE 1976: 133 (taxonomy, world catalogue).

**Sphex albiseptus* LEPELETIER DE SAINT-FARGEAU & SERVILLE, 1828: KOHL 1894: 342 (Gabon without exact locality).

Sphex (Parasphe) albiseptus LEPELETIER DE SAINT-FARGEAU & SERVILLE, 1828: BERLAND 1926a: 168 (Gabon without further data).

**Sphex albiseptus congoensis* n.var.: BERLAND 1926a: 169 (description ♀, Libreville).

Sphex (Parasphe) albiseptus race *marginatum* var. *congoensis* BERLAND, 1926: ARNOLD 1928b: 352 (taxonomy (synonym), not seen).

Chlorion (Parasphe) albiseptus var. *congoensis* (BERLAND, 1926): ARNOLD 1930: 17 (taxonomy, catalogue Afrotropical region).

Sphex albiseptus var. *congoensis* BERLAND, 1926: LECLERCQ 1955: 35 (synonym of *Sphex (Prionyx) albiseptus marginatus sjöstedti* CAMERON, 1910).

Sphex (Parasphe) albiseptus var. *congoensis* BERLAND, 1926: BERLAND 1956: 1169 (taxonomy, Gabon within French Congo).

Sphex albiseptus var. *congoensis* BERLAND, 1927 (!): BOHART & MENKE 1976: 133 (lectotype designation by Menke (♀ Libreville)).

**Sphex (Prionyx) albiseptus marginatus sjöstedti* CAMERON, 1910: LECLERCQ 1955: 33 (key to subspecies and varieties), 35 (taxonomy, "Chûtes de Samlia, riv. Nord de Gamie" = Chûtes de Samba (Rivière Ngounié) near Sindara).

Distribution: Estuaire: Libreville, no exact locality (KOHL 1894) – Ngounié: Chûtes de Samba (Rivière Ngounié) near Sindara – no exact locality (BERLAND 1926a).

Prionyx kirbii marginatus, which is widely distributed in Afrotropical region and on the Arabian Peninsula, is also known from the Republic of Congo.

***Prionyx viduatus* (CHRIST, 1791)**

Sphex (Parasphe) viduatus CHRIST, 1791: BERLAND 1956: 1167 (taxonomy, description ♂ ♀, au-delà d'Abembé = Alembé), 1179 (key).

Sphex (Prionyx) viduatus CHRIST, 1791: LECLERCQ 1955: 36 (taxonomy, Chûtes de Samlia, riv. N'Gamie = Chûtes de Samba (Rivière Ngounié) near Sindara).

Priomyx viduatus (CHRIST, 1791): BOHART & MENKE 1976: 125 (fig. 28F), 134 (world catalogue).
 **Sphex pubescens* FABRICIUS, 1793: FAIRMAIRE 1858: 264 (Gabon without exact locality).

Distribution: Estuaire: no exact locality (Fairmaire 1858) – Moyen-Ogooué: Alembé – Ngounié: Chûtes de Samba (Rivière Ngounié) near Sindara.

Prionyx viduatus, which is widely distributed in the Afrotropical and Palaearctic regions, is also recorded from Cameroon.

***Prionyx* sp.**

Enodia fervens (FABRICIUS, 1775): FAIRMAIRE 1858: 264 (Gabon without exact locality – misidentification).

Distribution: Estuaire: no exact locality (FAIRMAIRE 1858).

This is probably the record of *Sphex (Parasphex) albiseptus* LEPELETIER DE SAINT-FARGEAU & SERVILLE, 1828 = *Prionyx kirbii marginatus* (SMITH, 1856) in BERLAND 1926a (p. 168).

***Sphex caeruleanus* DRURY, 1773**

Sphex (Sphex) caeruleanus DRURY, 1773: BOHART & MENKE 1976: 114 (taxonomy, world catalogue).

**Sphex nigripes* var. *pulchripennis* MOCSÁRY, 1883: KOHL 1890: 147 (key, Gabon without exact locality).

Sphex pulchripennis MOCSÁRY, 1883: KOHL 1895: 63 (taxonomy, description ♂, Gabon without exact locality).

Sphex pulchripennis MOCSÁRY, 1883: DALLA TORRE 1897: 365 (world catalogue).

Sphex (Sphex) pulchripennis MOCSÁRY, 1883: BERLAND 1927: 153 (taxonomy, Gabon without exact locality).

Sphex haemorrhoidalis var. *pulchripennis* MOCSÁRY, 1883: LECLERCQ 1955: 14 (key), 16 ("Chûtes de Samlia, riv. Nord Gamie" = Chûtes de Samba (Rivière Ngounié) near Sindara).

Sphex (Sphex) haemorrhoidalis var. *pulchripennis* MOCSÁRY, 1883: BERLAND 1956: 1177 (taxonomy, description ♂ ♀, Ogooué), 1180 (key).

Sphex haemorrhoidalis f. *pulchripennis* MOCSÁRY, 1883: LECLERCQ 1969: 1049 (taxonomy, Mbila – Gabon, not Republic of Congo).

Material examined: Gabon: no exact locality, 1♀ – Republic of Congo (mislabelled: Gabon): Kuila = Quilou, 4♂♂ 1♀, 1894.

Distribution: Ngounié: Chûtes de Samba (Rivière Ngounié) near Sindara, Mbila – no exact locality (KOHL 1890, 1895, BERLAND 1927, 1956, this paper).

Sphex caeruleanus is also recorded from Benin, Cameroon, Democratic Republic of Congo, Ghana, Ivory, Republic of Congo, Republic of Guinea, Sierra Leone and South Africa.

***Sphex fumicatus* CHRIST, 1791**

Sphex (Sphex) fumicatus CHRIST, 1791: BOHART & MENKE 1976: 114 (world catalogue).

**Sphex umbrosus* var. *metallicus* TASCHENBERG, 1869: KOHL 1894: 342 (Cap Lopez, Gabon without exact locality).

Sphex (Sphex) umbrosus var. *metallicus* TASCHENBERG, 1869: LECLERCQ 1955: 21 ("Chutes de Samlia, riv. Nord Gamie" = Chûtes de Samba (Rivière Ngounié) near Sindara).

Sphex (Sphex) umbrosus var. *metallicus* TASCHENBERG, 1869: BERLAND 1956: 1174 (taxonomy, description ♂ ♀, biology, Gabon without exact locality), 1175 (fig. 4).

M a t e r i a l e x a m i n e d : Gabon without exact locality, 4 ♀♀, 1896, leg. Herman.

D i s t r i b u t i o n : Estuaire: no exact locality (KOHL 1894, this paper) – Ngounié: (Chûtes de Samba (Rivière Ngounié) near Sindara – Ogooué-Maritime: Cap Lopez – no exact locality (BERLAND 1956).

Sphex fumicatus is known from the Afrotropical and Palaearctic regions. There are no records from the neighbour countries.

***Sphex haemorrhoidalis* FABRICIUS, 1781**

Sphex (Sphex) haemorrhoidalis FABRICIUS, 1781: BOHART & MENKE 1976: 115 (world catalogue).

**Sphex nigripes volubilis* n.var.: KOHL 1895: 64 (description ♂ ♀, "Chûtes de Samlia, Riv. N. Gamio" = Chûtes de Samba (Rivière Ngounié) near Sindara), pls. 4 (fig. 14), 5 (figs. 35, 35).

Sphex nigripes var. *volubilis* KOHL, 1895: DALLA TORRE 1897: 434 (world catalogue, Gabon – not West Africa).

Sphex haemorrhoidalis subsp. *nigripes* var. *volubilis* KOHL, 1895: TURNER 1912: 369 (taxonomy, *nigripes* is subspecies of *haemorrhoidalis*, but var. *volubilis* not mentioned).

Sphex nigripes var. *volubilis* KOHL, 1895: BERLAND 1928: 329 (Franceville, Lambaréné).

Chlorion (Proterosphe) *haemorrhoidalis* var. *volubilis* (KOHL, 1895): ARNOLD 1930: 18 (taxonomy, catalogue Afrotropical Region).

Sphex haemorrhoidalis var. *volubilis* KOHL, 1895: LECLERCQ 1955: 15 (key).

Sphex (Sphex) haemorrhoidalis var. *volubilis* KOHL, 1895: BERLAND 1956: 1177 (taxonomy, description ♂ ♀, Franceville, Lambaréné).

Sphex haemorrhoidalis volubilis KOHL, 1895: BOHART & MENKE 1976: 115 (synonym).

D i s t r i b u t i o n : Haut-Ogooué: Franceville – Moyen-Ogooué: Lamabaréné – Ngounié: Chûtes de Samba (Rivière Ngounié) near Sindara.

Sphex haemorrhoidalis, which is widely distributed in the Afrotropical region, is also recorded from Cameroon and the Republic of Congo.

***Sphex nigrohirtus* KOHL, 1895**

Sphex (Sphex) nigrohirtus KOHL, 1895: BERLAND 1956: 1178 (taxonomy, description ♂ ♀, Lambaréné).

Sphex (Sphex) nigrohirtus (KOHL, 1895): BOHART & MENKE 1976: 115 (taxonomy, world catalogue).

**Sphex (Sphex) conradii* nov.sp.: BERLAND 1927: 154 (description ♀, Lambaréné), 155 (fig. 16).

Sphex conradii BERLAND, 1927: ARNOLD 1951: 145 (synonym).

Sphex (Sphex) conradii BERLAND, 1927: BERLAND 1953: 275 (synonym).

Sphex conradii BERLAND, 1927: BOHART & MENKE 1976: 115 (lectotype designation by Menke (♀ Bioko)).

M a t e r i a l e x a m i n e d : Gabon: Moyen-Ogooué Province: Ndjolé, Hotel Kedyann, 1♀, 05.02.2019, leg. M. Madl – Ogooué-Ivindo Province: RN 4 Koundameyong – Lalara, river W Djidji (0°14'19"N 11°46'41"E), ca. 410 m, 1♀, 04.02.2019, leg. M. Madl – Republic of Congo (mislabelled: Gabon): Kuila = Quilou, 1♀, 1894.

D i s t r i b u t i o n : Moyen-Ogooué: Lambaréné, Ndjolé – Ogooué-Ivindo: Djidji.

Sphex nigrohirtus, which is widely distributed in the Afrotropical region, is also known from Equatorial Guinea (Mbini), Cameroon and the Republic of Congo.

***Sphex tomentosus* FABRICIUS, 1787**

Sphex (Sphex) tomentosus FABRICIUS, 1787: BOHART & MENKE 1976: 117 (world catalogue).

**Sphex tuberculatus* SMITH, 1873: SCHULZ 1912: 167 (Gabon without exact locality).

Sphex (Sphex) tuberculatus SMITH, 1873: BERLAND 1927: 152 (Gabon without exact locality).
Sphex (Sphex) tuberculatus SMITH, 1873: BERLAND 1956: 1176 (taxonomy, description ♂ ♀, LIBREVILLE), 1180 (key).

D i s t r i b u t i o n : Estuaire Province: Libreville, no exact locality (SCHULZ 1912, BERLAND 1927).

Sphex tomentosus, which is widely distributed in the Afrotropical region, is also recorded from Cameroon.

Acknowledgements

I am indebted to Hans K. Mipounga (Institut de Recherche Agronomiques et Forestières, Libreville, Gabon) for his generous help during the collecting trip and discussions about the geographic positions of historical collecting sites. I would like to thank Katja Furlanić and Marvyn O. Ndong Sima (Libreville, Gabon) for their hospitality in Gabon. I am very grateful to Manuela Vizek (Natural History Museum Vienna, Austria) for her assistance in the Hymenoptera Collection and the former librarian Wolfgang Brunnbauer (Natural History Museum Vienna, Austria), who provided many references. I don't know how to express my thanks to Manfred A. Jäch (Natural History Museum Vienna, Austria) for his remarkable understanding for collecting insects outside freshwater during our expedition to Gabon.

Zusammenfassung

Die Familie Sphecidae ist in Gabun mit 15 Arten und eine Unterart aus den Gattungen *Chlorion* LATREILLE, 1802 (eine Art), *Chalybion* DAHLBOM, 1843 (drei Arten), *Isodontia* PATTON, 1880 (zwei Arten), *Prionyx* VANDER LINDEN, 1827 (zwei Arten), *Sceliphron* KLUG, 1801 (zwei Arten und eine Unterart) und *Sphex* LINNAEUS, 1758 (fünf Arten) nachgewiesen. Neue Funddaten aus Gabun und der Republik Kongo werden mitgeteilt.

References

- ARNOLD G. (1928a): The Sphegidae of South Africa Part X. — Annals of the Transvaal Museum **12/3**: 233-279, pls. 9-11.
- ARNOLD G. (1928b): The Sphegidae of South Africa Part XI. — Annals of the Transvaal Museum **12/4**: 338-375.
- ARNOLD G. (1930): A Check-List of the Sphecidae of the Ethiopian Region. — Transvaal Museum (University Press Cambridge, England): 21 pp.
- BERLAND L. (1926a): Les Sphegidae (Hyménoptères) du Muséum National de Paris. — Bulletin du Muséum National d'Histoire Naturelle Paris **32/3**: 163-170.
- BERLAND L. (1926b): Les Sphegidae (Hyménoptères) du Muséum National de Paris (3e Note). — Bulletin du Muséum National d'Histoire Naturelle Paris **32/5**: 282-285.
- BERLAND L. (1927): Les Sphegidae (Hyménoptères) du Muséum National de Paris (4e Note). — Bulletin du Muséum National d'Histoire Naturelle Paris **33/2**: 150-156.
- BERLAND L. (1928): Les Sphegidae (Hyménoptères) du Muséum National de Paris (5e Note). — Bulletin du Muséum National d'Histoire Naturelle Paris **34/5**: 329-331.
- BERLAND L. (1953): XII – Hyménoptères vespiformes. — Mémoires de l'Institut Française d'Afrique Noire **19** (La Réserve Naturelle Intégrale du Mt. Nimba): 271-276.
- BERLAND L. (1956): Les *Sphex* africains (Hymenoptera). — Bulletin de l'Institut Français d'Afrique Noire, Série A, Sciences Naturelles, **18/4**: 1161-1181.

- BOHART R.M. & A.S. MENKE (1976): Sphecid Wasps of the World. A Generic Revision. — Berkeley, Los Angeles, London (University of California Press): IX + 695 pp.
- COSTA A. (1866): Aquisiti fatti durante l'Anno 1863. — Annuario del Museo Zoologico della R. Università di Napoli 3/1: 13-41.
- DALLA TORRE C.G. de (1897): Catalogus hymenopterorum hucusque descriptorum systematicus et synonymicus. Volumen VIII: Fossores (Sphecidae). — Leipzig (Wolfgang Engelmann): VIII + 749 pp.
- DOLFFUSS H. (1989): Verzeichnis der Grabwespentypen am Naturhistorischen Museum in Wien (Hymenoptera, Sphecidae). — Kataloge der wissenschaftlichen Sammlungen des Naturhistorischen Museums in Wien 7/Entomologie 4: 26 pp.
- FAIRMAIRE L. (1858): IV. Ordre Hyménoptères. — In: THOMSON J. (ed.): Archives Entomologiques 2 (Voyage au Gabon. Histoire naturelle des insectes et des arachnides recueillis pendant un voyage fait au Gabon en 1856 et en 1857 par M. Henry C. Deyrolle sous les auspices de MM. Le Comte de Mniszech et James Thomson précédée de histoire du voyage par M. James Thomson): 263-267.
- HENSEN R.V. (1988): Revision of the nominate subgenus *Chalybion* DAHLBOM (Hymenoptera, Sphecidae). — Tijdschrift voor Entomologie 131/1: 13-64.
- KIRBY W.F. (1896): A List of the Orthoptera, Hymenoptera, and Hemiptera collected by Miss Kingsley on the river Ogové, with Descriptions of some new Genera and Species. — The Annals and Magazine of Natural History, 6. Series, 18/105: 257-269, pl. 12.
- KOHL F.F. (1890): Die Hymenopterengruppe der Sphecinen. I. Monographie der natürlichen Gattung *Sphex* LINNÉ (sens. lat.). — Annalen des k.k. Naturhistorischen Hofmuseums in Wien 5/2: 77-194, pls. 8-10.
- KOHL F.F. (1894): Zur Hymenopterenfauna Afrikas. — Annalen des k.k. Naturhistorischen Hofmuseums in Wien 9/3-4: 279-350, pls. 13-17.
- KOHL F.F. (1895): Zur Monographie der natürlichen Gattung *Sphex* LINNÉ. — Annalen des k.k. Naturhistorischen Hofmuseums in Wien 10/1: 42-74.
- KOHL F.F. (1918): Die Hautflüglergruppe Sphecinae. IV. Teil. Die natürliche Gattung *Sceliphron* KLUG (*Pelopoeus* LATR.). — Annalen des k.k. Naturhistorischen Hofmuseums in Wien 32: 1-171.
- LECLERCQ J. (1955): Hymenoptera Sphecoidea (Sphecidae I. Subfam. Sphecinae) avec un catalogue synonymique des espèces africaines et des considerations sur la distribution des espèces au Congo Belge. — In: Exploration du Parc National de l'Upemba Mission I. G.F. De Witte en collaboration avec W. Adam, A. Janssens, L. Van Meel et R.R. Verheyen (1946-1949) 34: 137 pp.
- LECLERCQ J. (1969): Contribution à la faune du Congo (Brazzaville). Mission A. Villiers et A. Descarpentries XC. Hyménoptères Sphecidae. — Bulletin de l'Institut Fondamental d'Afrique Noire, série A, 31/3: 1046-1054.
- RASMUSSEN C. & J.S. ASCHER (2015): Addendum to: Joseph Vachal (1838-1911): French entomologist and politician (Zootaxa, 3442, 1-52). — Zootaxa 3931/2: 298-300.
- RÉGIMBART M. (1895): Revision des Dytiscidae et Gyrinidae d'Afrique, Madagascar et Iles voisines. En Contribution a la faune entomologique du Congo. — Mémoires de la Société Entomologique de Belgique 4: 244 pp.
- SCHOOTEDEN H. (1918): Les Aradides du Congo Belge. — Revue de Zoologique Africaine 6: 127-137.
- SCHULZ W.A. (1911): Zweihundert alte Hymenopteren. — Zoologische Annalen 4/1-2: 1-220.
- THOMSON J. (1858): Histoire de Voyage. — In: THOMSON J. (ed.): Archives Entomologiques 2 (Voyage au Gabon. Histoire naturelle des insectes et des arachnides recueillis pendant un voyage fait au Gabon en 1856 et en 1857 par M. Henry C. Deyrolle sous les auspices de MM. Le Comte de Mniszech et James Thomson précédée de histoire du voyage par M. James Thomson): 9-27.

TURNER R.E. (1912): Notes on Fossiliferous Hymenoptera. — X. — The Annals and Magazine of Natural History, 8th Series, **10**/58: 361-377.

VECHT J. VAN DER & F.M.A. VAN BREUGEL (1968): Revision of the nominate subgenus *Sceliphron* LATREILLE (Hymenoptera, Sphecidae (Studies on the Sceliphronini, part I). — Tijdschrift voor Entomologie **111**/6: 185-255.

Author's address:

Michael MADL
Naturhistorisches Museum Wien
2. Zoologische Abteilung
Burgring 7
A-1010 Vienna, Austria
E-Mail: michael.madl@nhm-wien.ac.at

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Linzer biologische Beiträge](#)

Jahr/Year: 2021

Band/Volume: [0053_1](#)

Autor(en)/Author(s): Madl Michael

Artikel/Article: [Notes on the Sphecidae \(Hymenoptera\) of Gabon 101-110](#)