

Linzer biol. Beitr.	54/2	399-402	Februar 2023
---------------------	------	---------	--------------

On the taxonomy of *Ischnoglossa* KRAATZ, 1856: two new synonymies and a new combination (Coleoptera, Staphylinidae, Aleocharinae, Oxypodini)

Volker ASSING†

A b s t r a c t : Two synonymies and a new generic assignment are proposed: *Xenodota* PACE, 1982 (subgenus of *Ischnoglossa* KRAATZ, 1856) = *Philoleptusa* PACE, 1983, nov.syn.; *Ischnoglossa minor* (AUBÉ, 1863) = *Leptusa syriaca* EPPELSHEIM, 1889, nov.syn. (type species of *Philoleptusa*); *Ischnoglossa picipennis* (SOLSKY, 1874), nov.comb. (ex *Philoleptusa*). Additional records of *Ischnoglossa minor* are reported, among them the first record from continental Spain.

K e y w o r d s : Coleoptera, Staphylinidae, Aleocharinae, Oxypodini, *Ischnoglossa*, new synonymies, new combination, additional records.

Introduction

According to the latest edition of the Palaearctic catalogue (SCHÜLKE & SMETANA 2015), the genus *Ischnoglossa* KRAATZ, 1856 is represented in the Palaearctic region by seven species in two subgenera: the nominal subgenus with six species (three West Palaearctic, one trans-Palaearctic, and two East Palaearctic) and *Xenodota* PACE, 1982 with one species, *Ischnoglossa minor* (AUBÉ, 1863), distributed in the Mediterranean region and the Canary Islands.

Originally described in the genus *Homalota* MANNERHEIM, 1839, *Ischnoglossa minor* was subsequently attributed to the subgenus *Microdota* MULSANT & REY, 1873 (BRUNDIN 1948). In an article dealing with Northwest African and Iberian *Microdota* species, PACE (1982) discovered that *I. minor* in fact belonged to *Ischnoglossa* of the Oxypodini, figured the mouthparts, habitus, other external characters, and the primary sexual characters, including the highly distinctive spermatheca, and assigned the species to the newly described monotypical subgenus *Xenodota*. According to the description and the key provided by PACE (1982), *Xenodota* is distinguished from the nominal subgenus by smaller size, a shorter antennomere III, a stout and rather short maxillary palpomere III, the absence of an anterior impression on the abdominal tergite VI, impunctate anterior impressions on tergites III-V, and a spermatheca with less numerous coils.

In the following year, in a revision of material of *Leptusa* KRAATZ, 1856 in the Scheerpeltz collection, PACE (1983) discovered that *Leptusa syriaca* EPPELSHEIM, 1889, a species described from Syria, did not belong to *Leptusa* and assigned it to the newly described monotypical genus *Philoleptusa*, stating that it was closely allied to *Ischnoglossa*. Strangely, however, *Philoleptusa* is erroneously listed in the tribe Homalotini in the

Palaearctic Catalogue (SCHÜLKE & SMETANA 2015). In his comprehensive revision of *Leptusa*, PACE (1989) subsequently moved *Leptusa picipennis* SOLSKY, 1874 from Uzbekistan to *Philoleptusa*, without stating his reasons and without illustrating characters.

Some years ago, the late Marc Tronquet (Molitg-les-Bains) sent me photos of a female specimen from France that he had identified as *Philoleptusa syriaca* based on the illustrations provided by PACE (1983). The shape of the spermatheca eventually aroused the suspicion that *P. syriaca* was in fact identical to *Ischnoglossa minor*.

Material and methods

The material reported in this study is deposited in the following collections:

- cAss..... author's private collection
- cTro..... private collection Marc Tronquet (Molitg-les-Bains)
- cVog..... private collection Jürgen Vogel, Görlitz
- cWun..... private collection Paul Wunderle

Results

A study of the descriptions and illustrations provided by PACE (1982, 1983) leave no doubt that *Ischnoglossa minor* (type species of the subgenus *Xenodota*) and *Philoleptusa syriaca* (type species of *Philoleptusa*) are in fact conspecific. The drawings of the median lobe of the aedeagus (PACE 1982: figures 30-32; PACE 1983: figures 5-6) differ somewhat regarding the shape of the ventral process, but these differences are most likely a result of artefacts resulting from the dissection and preparation of the genitalia. The highly distinctive spermatheca of *Philoleptusa syriaca* (PACE 1983: figure 4) is clearly that of *Ischnoglossa minor*. These findings result in the following new synonymies and new combination.

Sub genus *Xenodota* PACE, 1982

Xenodota PACE, 1982: 81 ff. (type species: *Homalota minor* AUBÉ, 1863).

Philoleptusa PACE, 1983: 57 f. (type species: *Leptusa syriaca* EPPELSHEIM, 1889); nov.syn.

***Ischnoglossa (Xenodota) minor* (AUBÉ, 1863)**

Homalota minor AUBÉ, 1863: 380.

Ocyusa postica MULSANT & REY, 1875: 434.

Leptusa syriaca EPPELSHEIM, 1889: 161; nov.syn.

Ischnoglossa pulchella ISRAELSON, 1969: 151.

M a t e r i a l e x a m i n e d : CANARY ISLANDS: Tenerife: 1♂, Teide National Park, El Portillo, 2050 m, pine forest, litter sifted, 11.IV.1992, leg. Assing (cAss); 2 exs. [identified by J. Vogel], Aguamansa, under pine bark, 12.VI.2011, leg. Peschel (cVog). La Palma: 1♂, 4♀♀, 5 exs., NW Roque de los Muchachos, 1700 m, pine forest, under pine bark, 3.IV.1999, leg. Assing & Wunderle (cAss, cWun). El Hierro: 2♂♂, 4♀♀, 1 ex., El Pinar, leg. Franz (cAss, cWun). SPAIN: 2 exs. [identified by J. Vogel], Alicante, Sierra de Aitana, Castell de Guadalest, 25.III.1971 (cVog). FRANCE: 1♀ [identified based photos], Var, Collobrières env., 11.II.1952, leg. Joffre (cTro); 1♂ [identified based a photo], Bouche-du-Rhône, Col de Sormiou, decomposing dry trunk with mold, 8.II.2010, leg. Gompel (cTro). GREECE: Peloponnese: 1♀, Olympia, under bark of pine, 1.IV.1986, leg. Assing (cAss); 1♂, Ahaia, Kalogria, 24-25.IV.1999, leg. Angelini

(cAss); 1♀, Ahaia, E Kato Klitoria, Agios Nikolaos, 25.IV.1999, leg. Brachat (cAss); 1♀ [identified by P. Wunderle], Erimanthos, 500 m, 24.III.1992, leg. Frisch (cWun). C o r f u : 1♀, [identified by P. Wunderle], Kerkyra, old castle, 13.X.1987 (cWun); 2 exs. [identified by J. Vogel], Moraitika, V.2010, leg. Sieber (cVog). S a m o s : 1♀, N Psili Amos, 37°43'N, 26°59'E, 90 m, under bark near stream, 19.IV.2003, leg. Brachat & Meybohm (cAss).

C o m m e n t : *Ischnoglossa minor* is found under bark and rather widespread in the Mediterranean region and in the Canary Islands. In the Mediterranean it is a rare species and has been recorded from Northwest Africa (Tunisia, Algeria, Morocco), Spain, France, Greece, Cyprus, and Syria (material examined; SCHÜLKE & SMETANA 2015). The above specimens include the first records from mainland Spain and from Greece. In the Canary Islands, *I. minor* appears to be less rare (see material examined).

***Ischnoglossa (Xenodota) picipennis* (SOLSKY, 1874), nov.comb.**

Leptusa picipennis SOLSKY, 1874: 159.

Philoleptusa picipennis (SOLSKY, 1874): PACE (1989).

The identity of this species, which has been recorded only from Uzbekistan, is currently unclear. The proposed combination results from the synonymy of *Philoleptusa* with *Xenodota*, but requires confirmation based on a revision of type material.

Acknowledgements

The present paper was initiated by photos and a request sent to me by the late Marc Tronquet. Marc Tronquet, Jürgen Vogel, and Paul Wunderle communicated additional records of *Ischnoglossa minor*. Benedikt Feldmann (Münster) proof-read the manuscript.

Zusammenfassung

Zwei Namen werden synonymisiert: *Xenodota* PACE, 1982 (Subgenus von *Ischnoglossa* KRAATZ, 1856) = *Philoleptusa* PACE, 1983, nov.syn.; *Ischnoglossa minor* (AUBÉ, 1863) = *Leptusa syriaca* EPPELSHEIM, 1889, nov.syn. (Typusart von *Philoleptusa*). Aus ersterer Synonymie ergibt sich das Binomen *Ischnoglossa picipennis* (SOLSKY, 1874), nov.comb. (ex *Philoleptusa*). *Ischnoglossa minor* wird erstmals vom spanischen Festland und aus Griechenland nachgewiesen.

References

- AUBE C. (1863): [new taxa]. — In: GRENIER A., Catalogue des coléoptères de France, par M. le Dr. A. Grenier, et matériaux pour servir à la faune des coléoptères français. — A. Grenier, Paris: 3-79, 1-135.
- BRUNDIN L. (1948): *Microdota*-Studien. — Entomologisk Tidskrift **69**: 8-66.
- EPPELSHEIM E. (1889): Neue Staphylinen Europa's und der angrenzenden Länder. — Deutsche Entomologische Zeitschrift **33**: 161-183.
- ISRAELSON G. (1969): Some additions to the coleopterous fauna of the Canary Islands. — Eos, Revista Española de Entomología **44** [1968]: 149-157.
- MULSANT E. & C. REY (1875): Pp. 163-365. — In: Histoire naturelle des coléoptères de France. Brévipennes. Aléochariens (Suite). Aléocharaïres. — Deyrolle, Paris: 565 pp.

- PACE R. (1982): Studio su alcune specie iberiche e magrebine di *Atheta* del sottogenere *Microdota* MULS. & REY (Coleoptera Staphylinidae). — Giornale Italiano di Entomologia, Cremona **1**: 73-83.
- PACE R. (1983): Risultati dello studio delle specie del genere *Leptusa* KRAATZ della collezione Scheerpeltz al Naturhistorisches Museum di Vienna (Coleoptera, Staphylinidae). — Annalen des naturhistorischen Museum in Wien (B) **85**: 53-102.
- PACE R. (1989): Monografia del genere *Leptusa* KRAATZ (Coleoptera Staphylinidae). — Memorie del Museo Civico di Storia Naturale di Verona (II° Serie), Sezione Scienze della Vita (A: Biologica) **8**: 1-307.
- SCHÜLKE M. & A. SMETANA (2015): Staphylinidae, pp. 304-1134. — In: LÖBL I. & D. LÖBL (eds), Catalogue of Palaearctic Coleoptera. Volume 2. Hydrophiloidea – Staphyloinoidea. Revised and updated edition. Leiden: Brill: xxvi + 1702 pp. —
- SOLSKY S.M. (1874): Zhestkokrylye (Coleoptera), Tetrad 1 — In: FEDSCHENKO A.P. (ed.), Puteshestvie v Turkestan. Tom III, Zoogeograficheskiya izsledovaniya, Chast V, Otdel Shestoy. — Imperatorskoe obshschestvo lyubiteley estestvoznaniya, antropologii i etnografii, St. Petersburg and Moskva: iv + 222 pp. —

Previous address of the author: Dr. Volker ASSING†
 Gabelsbergerstr. 2
 D-30163 Hannover, Germany
 E-mail: vassing.hann@t-online.de

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Linzer biologische Beiträge](#)

Jahr/Year: 2023

Band/Volume: [0054_2](#)

Autor(en)/Author(s): Assing Volker

Artikel/Article: [On the taxonomy of Ischnoglossa KRAATZ, 1856: two new synonymies and a new combination \(Coleoptera, Staphylinidae, Aleocharinae, Oxypodini\) 399-402](#)