

Lauterbornia 40: 39-41, D-86424 Dinkelscherben, 2001-02-28

The zoobenthos of the Koprinka reservoir, Central Bulgaria

Stefan Stoichev

With 1 Table

Schlagwörter: Nematoda, Oligochaeta, Chironomidae, Diptera, Insecta, Koprinka See, Bulgarien, Stausee, Faunistik

Keywords: Nematoda, Oligochaeta, Chironomidae, Diptera, Insecta, Koprinka lake, Bulgaria, reservoir, faunistics

The study gives recent information on the zoobenthos of the Koprinka reservoir. For the first time data are reported on Nematoda. In total 22 benthic taxa were found: 9 nematodes, 3 oligochaets and 10 chironomids.

1 Short description of the lake

The Koprinka reservoir is located in the Kazanlak Valley, 7 km west of the town of Kazanlak and 35 km northwest of Stara Zagora. During the first years of exploitation additional hydro-technical equipment was installed by which the water level of the reservoir was raised 4-5 m, and the water surface increased from 8 400 000 to about 11 000 000 m²; the volume now is 138 084 000 m³. (DIMITROV & LYUDSKANOVA 1967). The lowest level of the reservoir is 356 m above sea level, the maximum is 391 m. The maximum depth near the impounding dam 35 m. Koprinka reservoir has three branches – Koprinski, Dunavski and Gyurlya, oriented on a north-south axis.

Previous data on the zoobenthos of the reservoir were given by DIMITROV (1960) and DIMITROV & LYUDSKANOVA (1967).

2 Material and methods

The zoobenthic material was collected by the author in May, July, August, September and December 1995-2000 using a Birge-Eckman bottom sampler. The samples were rinsed on sieves, mesh width 500 μ m and 150 μ m. To get the nematodes extended, they were heated up to 60 °C in a water basin and then fixed in 4 % formaline solution; preparation and identification according to GAGARIN (1981) on the basis of the formula of DE MAN (1886). For identifying the Chironomidae larvae we followed PANKRATOVA (1970, 1977, 1983).

The samples were collected from 6 sites in the reservoir: 1. East; 2. Middle; 3. West; 4. Koprinski Rakav; 5. Dunavski Rakav; 6. Rakav Gyurlya.

The boecenotic analysis (frequency of occurrence pF %, frequency of dominance DF %, and range of dominance Dt %) was made by the method of DE VRIES (1937).

Tab. 1: Species composition and dominance analysis of the zoobenthos of Koprinka reservoir

	Sites						Dominance analysis		
	1	2	3	4	5	6	pF %	DF %	Dt %
NEMATODA									
<i>Dorylaimus stagnalis</i> DUJARDIN	x	x	x	x	x	x	92.0	76.0	82.6
<i>Monhystera stagnalis</i> BASTIAN					x	x	16.0		
<i>Aphanolaimus aquaticus</i> DE DAY						x	4.0		
<i>Enoploides fluviatilis</i> MICOLETZKY		x			x		8.0		
<i>Enoploides</i> sp.				x			4.0		
<i>Tripyla glomerans</i> BASTIAN		x	x		x	x	68.0	56.0	82.3
<i>Tripyla</i> sp.					x		4.0		
<i>Tobrilus gracilis</i> (BASTIAN)	x		x	x	x		80.0	4.0	5.0
<i>Rhabditida</i> CHITWOOD									
<i>Diplogasteridae</i> MICOLETZKY									
<i>Paraigolaimella anomala</i> GAGARIN						x	4.0		
OLIGOCHAETA									
<i>Limnodrilus hoffmeisteri</i> CLAPAREDE					x	x	8.0		
<i>Limnodrilus claparedeanus</i> ROTZEL					x		4.0		
DIPTERA: CHIRONOMIDAE									
<i>Chironomus</i> gr. <i>plumosus</i> KIEFFER	x	x	x	x	x	x	96.0	88.0	91.6
<i>Chironomus</i> sp.					x		4.0		
<i>Chironomus riparius</i> MEIGEN			x	x		x	32.0	4.0	12.5
<i>Cryptochironomus</i> gr. <i>defectus</i> KIEFFER	x		x		x	x	60.0	8.0	13.0
<i>Cryptochironomus</i> sp.					x		4.0		
<i>Cricotopus</i> gr. <i>sylvestris</i> (FABRICIUS)			x	x	x		56.0	14.0	7.1
<i>Cricotopus sylvestris</i> (FABRICIUS)				x	x		12.0	4.0	33.3
<i>Polypedilum</i> gr. <i>nubeculosum</i> (MEIGEN)						x	8.0		
<i>Tanytarsus</i> gr. <i>gregarius</i> KIEFFER		x		x	x		20.0	8.0	40.0
<i>Tanytarsus mancus</i> WALKER			x				4.0		
<i>Limnophyes</i> gr. <i>prolongatus</i> KIEFFER					x		4.0		

3 Species composition and dominance analysis (Tab. 1)

Dorylaimus stagnalis and *Chironomus* gr. *plumosus* were found in the whole reservoir. Beside the species of high pF % and high Dt % values (*Dorylaimus stagnalis*, *Chironomus* gr. *plumosus* and *Tripyla glomerans*), species with high presence frequency and low values of the range of dominance (*Cricotopus* gr. *sylvestris*, *Cryptochironomus* gr. *defectus*, *Tobrilus gracilis*) were found. The chironomids *Cricotopus sylvestris* and *Tanytarsus* gr. *gregarius* are of low frequency, but of higher importance as dominants.

Acknowledgements

I like to thank Vera Staykova for her help in collecting the material, and Nikolai Chernev for technical support.

References

- DIMITROV, M. (1960): Hydrologische und hydrobiologische Charakteristik der Talsperre "Koprinka".- In: Arbeiten des zentralen Forschungsinstitutes fuer Fischzucht und Fischerei, Abteilung Suesswasserfischerei - Plowdiw (Bulgarien), Ed. Of the State: 87-122, Varna
- DIMITROV, M. & ZH. LYUDSKANOVA (1967): Feeds and feeding correlation of commercial fish in the "Koprinka" dam.- In: Proceedings of the Research Institute of Fisheries and Oceanography, 8: 339-358, Varna
- GAGARIN, V. (1981): Opredelitel - presnovodnie nematody evropejskoj chasti SSSR, Ed. Nauka.- Monogr.:1-248, St. Petersburg
- MAN, J. DE (1886): Anatomische Untersuchungen Über Freilebende Nordsee Nematoden.- 82 pp., (Paul Froberg) Leipzig
- PANKRATOVA, V. (1970): Lichinki i kukolki komarov podsemeistva Orthoclaadiinae fauny SSSR (Diptera, Chironomidae=Tendipedidae).- Opredelitel Faune SSSR 102: 1-343, Leningrad (in Russian)
- PANKRATOVA, V. (1977): Lichinki i kukolki komarov podsemeistv Podonominae i Tanypodinae fauny SSSR (Diptera, Chironomidae=Tendipedidae).- Opredelitel Faune SSSR 112: 1-155, Leningrad (in Russian)
- PANKRATOVA, V. (1983): Lichinki i kukolki komarov podsemeistva Chironominae fauny SSSR (Diptera, Chironomidae=Tendipedidae). -Opredelitel Faune SSSR 134: 1-296, Leningrad (in Russian)
- VRIES, M. DE (1937): Methods used in plant sociology and agricultural botanical grassland research.- Herbage Review 5: 76-82, Aberystwyth

Author's adress: Dr. Stefan Stoichev, Bulgarian Academy of Sciences, Institute of zoology, Boul. Tsar Osvoboditel 1, BG-1000 Sofia, Bulgaria

Received: 2000-09-23

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Lauterbornia](#)

Jahr/Year: 2001

Band/Volume: [2001_40](#)

Autor(en)/Author(s): Stoichev Stefan

Artikel/Article: [The zoobenthos of the Koprinka reservoir, Central Bulgaria. 39-41](#)