

The Fossil Planipennia – a Review

**By Thomas SCHLÜTER, Dar es Salaam
Department of Geology, University**

ABSTRACT

Systematics, palaeontology and stratigraphy of the orders of the Neuropteroidea are briefly discussed and previous work is summarized. From different authors 33 families have been established (19 existent and 14 extinct) within the order Planipennia and are presented in a phylogenetic tree which is mainly based on the classification outline of WITHYCOMBE (1924). Many questions on the phylogenetic relationships remain open since only the usefulness of morphologically recognizable characters provides palaeontological evidence in a stratigraphic context.

1. THE PHYLOGENETIC RELATIONSHIPS OF THE NEUROPTEROIDEA

The Neuropteroidea are doubtlessly the most archaic group of the Holometabola, but up to now it is not clearly evidenced if the 3 recent orders Megaloptera, Raphidioptera and Planipennia (i. e. Neuroptera s. str.) really represent higher categories with a common ancestor, i. e. if they are of monophyletic origin. Phylogenetic interpretations mainly based on morphological characters of the living species have been given by ACHELIG (1975 and 1981), HENNIG (1969), ASPÖCK et. al. (1980), important former authors are HANDLIRSCH (1906 - 1908), TILLYARD (1926), WITHYCOMBE (1924), CARPENTER (1943 and 1954) and O. MARTYNOVA (1952 and 1962). Generally the Megaloptera, Raphidioptera and Planipennia are well accepted as monophyletic groups and treated as orders within the Neuropteroidea, but only a few - often cryptic - characters indicate so.

However, the phylogenetic relationships of these orders became even more confused when in 1980 RODENDORF & RASNITSYN published their book on "The historical development of the class insecta" (in Russian) and the extinct group Glosselytrodea (synonymous with Jurinida) was added. The Glosselytrodea represent a taxon stratigraphically well distributed mainly in palaeozoic times which was formerly assigned to the Paurometabola. The opinion of RODENDORF & RASNITSYN on the phylogenetic relationships of the Neuropteroidea is here given in figure 1 and should be valid as an arbitrary scheme pending further discussions.

2. INTERPRETATION OF FOSSILS

Fossil Neuropteroidea are mainly preserved in sedimentary rocks as wing or, rarely, as almost complete specimen impressions or, in an often much better state of preservation, in fossiliferous resins; the former evidenced the stratigraphic record from Upper Permian upwards, the latter since early Cretaceous. Unfortunately not any derived groundplan characters are recognizable in the wing venation, hence it follows that especially Palaeozoic fossils that agree in the wing venation with recent Neuropteroidea should not automatically be assigned to one of their subgroups. For instance the determination of fossil Planipennia by their comb-like proliferation of branches of the radial sector is only tentative, since ACHTELIG (1980) gave evidence of this character in species of the extinct Palaeodictyoptera and Megasecoptera. "On the other hand, the non-pectinate radial sector, the fusion of the bases of M and Cu, and the distal fusion of the anal veins are not characters of the Neuropteroidea, let alone of the Planipennia" (ACHTELIG 1980).

3. STEMGROUP - PLANIPPENNIA

There are some fossils from late Palaeozoic (Upper Permian) which presumably belong to the Planipennia, but "the most striking similarity between the wings of several of "... these ..." Permian Planipennia and a number of recent families is the presence of trichosors (small thickenings of the wing margin between the tips of the veins and veinlets)" (ACHTELIG 1980). Different families (Palaeomerobiidae, Sialidopseidae, Permithonidae, Archeosmylidae and some synonymous names exist) from the Upper Permian of Australia, Siberia, South Africa and South America have been established, but actually we do not know whether they belong to the stemgroup or to more advanced recent subgroups of the Planipennia. These families are formally classified due to the amount of branches in the radial sector, the running position of the subcosta either into the costa or into the radius, and the size of the forewings. Probably all of these characters do not have significant value in a phylogenetic interpretation of the whole order, but they are sufficient for a palaeontological (more typological) classification outline, which, naturally, is based only on morphological peculiarities.

Possibly the Palaeomerobiidae and the Sialidopseidae are closer related, forming a superfamily Hemerobiidea (sensu MARTYNOVA 1962), whereas the Permithonidae and the Archeosmylidae, which survived up to the Triassic, are forming another superfamily Polystoechotidea (sensu MARTYNOVA 1962). CARPENTER (1954) considered the Lower Permian Permoberothidae to be the oldest Planipennia, but these are now generally assigned to the Glosselytroidea (RODENDORF & RASNITSYN 1980).

4. PROPOSALS FOR PHYLOGENETIC TREES OF THE PLANIPPENNIA

Another arbitrary or perhaps a working hypothesis is the assumption that all the categories in the rank of the recent families are monophyletic since we do not know the exact sister-group relationships between these families. Detailed classification outlines have been given already by HANDLIRSCH (1906 - 1908) and TILLYARD (1962), but WITHYCOMBE's phylogenetic tree published in 1924, in which he showed the presumed sister-group relationships within the order Planipennia, has been adopted in its main emphasis by later authors as FRIEDRICH (1953), and slightly modified by ASPÖCK et al. (1980) and SCHLÜTER (1984). Some objections against it came from the Russian school of Palaeoentomology, especially by Olga MARTYNOVA (1952) and more specified in 1962 (see RODENDORF et al. 1962). W. HENNIG compared and discussed both opinions in his famous book "Die Stammesgeschichte der Insekten" (1969) and figured the two different phylogenetic trees (fig. 2) proposed by MARTYNOVA and WITHYCOMBE. Later some additional extinct taxa in the rank of new families have been established and their relationships




Fig. 1: The phylogenetic relationships of the orders of the Neuropteroidea, as presumed and presented by RODENDORF & RASNITSYN (1980). Slightly modified especially in the nomenclature.

discussed. However, a main agreement in the classification of the superfamilies as shown here in table 1 is clearly recognizable. Obviously the comparison leads to the following still open questions:

1. The position of the Mantiscoidea:
 - a. They are derived from early Hemerobioidea after MARTYNOVA.
 - b. They are derived or at least closely related to the Osmyloidea after WITHYCOMBE.
2. The position of the Sisyridae:
 - a. They belong to the Hemerobioidea after MARTYNOVA.
 - b. They belong to the Osmyloidea after WITHYCOMBE, RODENDORF & RASNITSYN and other authors.

5. THE STRATIGRAPHIC RECORD OF THE FAMILIES OF THE PLANIPENNIA

The framework of the phylogenetic tree given in figure 3 is generally a modified version of that already published in WITHYCOMBE (1924). Virtually all the systematic categories in the rank of families established by later authors are incorporated herein. Yet this scheme is not wholly satisfactory, but for the moment the scheme here given, in which the order Planipennia is subdivided into 33 families (19 living and 14 extinct), appears to be reasonable in the light of consideration of both living and extinct groups in a joint lineage diagram.

MARTYNOVA (1962)**Myrmeleontidea**

Solenoptilidae
 Nymphitidae
 not mentioned
 Myrmeleonidae
 not mentioned
 Ascalaphidae
 Nemopteridae

Hemerobiidea

Psychopidae
 not mentioned
 Kalligrammatidae
 Brongniartiellidae
 Palaeomerobiidae
 Sialidopseidae
 Hemerobiidae
 Sisyridae
 Chrysopidae
 Mesochrysopidae

not mentioned

not mentioned
 Berothidae
 not mentioned
 Prohemerobiidae
 Proberothidae
 Osmylopsychosidae

Polystoechotidea

Permithonidae
 Archeosmylidae
 Osmylidae
 Osmylitidae
 belonging to Hemerobiidea
 Mesopolystoechotidae
 not mentioned
 not mentioned
 belonging to Hemerobiidea

Coniopterygidea

Coniopterygidae

not mentioned

not mentioned
 not mentioned

WITHYCOMBE (1924) and later authors**Myrmeleontoidea**

Solenoptilidae
 Nymphitidae
 Nymphidae
 Myrmeleonidae
 Stilbopterygidae
 Ascalaphidae
 Nemopteridae

Hemerobioidae

Psychopsidae
 Bruchaiseridae
 Kalligrammatidae
 ancestors of Rapimatidae after RIEK
 Palaeomerobiidae
 Sialidopseidae
 Hemerobiidae
 belonging to Osmyloidea
 Chrysopidae
 belonging to Chrysopidae

Mantispoidea

Dilaridae
 Berothidae
 Mantispidae
 not mentioned
 not mentioned
 belonging to Osmyloidea

Osmyloidea

Permithonidae
 Archeosmylidae
 Osmylidae
 Osmylitidae
 Osmylopsychopsidae
 Mesopolystoechotidae
 Polystoechotidae
 Neurorthidae
 Sisyridae

Coniopterygoidea

Coniopterygidae

Ithonoidea

Ithonidae
 Rapismatidae (incl. Brongniartiellidae)

Table 1: Comparison of the systematic classification outlines presented by MARTYNOVA (1962) and WITHYCOMBE (1924) for the families of the order Planipennia.




Fig. 2: Proposals of phylogenetic trees of the families of the Planipennia as given in MARTYNOVA (1952) and WITHYCOMBE (1924). Circles indicate the following taxa: 1: Permithonidae (incl. Permegalomidae) and Archeosmylidae from the Upper Permian of Australia and the USSR; 2: Palaemerobiidae from the Upper Permian of the USSR; 3: Petrushevskia borisi MARTYNOVA from the Upper Triassic of Issyk-Kul; 4: Triassopsychops superba TILLYARD from the Upper Triassic of Ipswich (Australia); 5: Sogjuta speciosa MARTYNOVA from the Upper Triassic of Issyk-Kul; 6: Mesypochrysa latipennis MARTYNOVA from the Upper Jurassic of Karatau (Kasachstan).

But similarly as presented in the book of HARLAND et al. (1967), another object of figure 3 lies essentially in the documentation of published records of fossil Planipennia in their stratigraphic context. First- and last-known fossil records of the respective families were to be related as precisely as possible to the standard scheme of the stratigraphic divisions and units. Such a compilation should be a valuable information for a variety of uses, e. g. such as to draw attention to the deficiencies of our present knowledge.

Particularly, more research on the known ranges of the living families is necessary, as opposed to the hypothetical ranges often ascribed to taxa in phylogenetic and other interpretations. Another aim of this paper is to provide a documentation of time ranges of the families of the Planipennia easily accessible to the non-palaeontologically informed and experienced neuropterologist.

REFERENCES

- ACHTELIG, M., 1975: Die Abdomenbasis der Neuropteroidea (Insecta, Holometabola). Eine vergleichend anatomische Untersuchung des Skeletts und der Muskulatur.-Zoomorphologie, 82: 201-242.




Fig. 3: The stratigraphic range of the families of the order Planipennia. 19 existent (in brackets the number of living species) and 14 extinct families are given. Wide beams give a more or less continuous fossil record for the respective family, thin lines indicate fossil gaps but presumable ranges. Lines connected by brackets give possible relationships.

- ACHTELIG, M., 1981: Revisionary notes.- In: *Insect Phylogeny*, by W. HENNIG: Chichester, New York, Brisbane, Toronto: John Wiley & Sons.
- ADAMS, P. A., 1967: A review of the Mesochrysinæ and Nothochrysinæ (Neuroptera: Chrysopidae).- *Bull. Mus. Compar. Zool.*, 135: 215-238.
- ASPÖCK, H., ASPÖCK, U., HÖLZEL, H., 1980: *Die Neuropteren Europas*. Krefeld: Goecke & Evers.
- CARPENTER, F. M., 1929: A Jurassic Neuroptera from the lithographic limestone of Bavaria.- *Psyche*, 36: 190-194.
- CARPENTER, F. M., 1943: The Lower Permian insects of Kansas, Part 9. The orders Neuroptera, Raphidioidea, Caloneuroidea, and Protorthoptera (Probnisidea), with additional Protodonata and Megasecoptera.- *Proc. amer. Acad. Arts Sci.*, 75: 55-84.
- CARPENTER, F. M., 1959: Fossil Nemopteridae (Neuroptera).- *Psyche*, 66: 20-24.
- CARPENTER, F. M., 1935: Tertiary insects of the family Chrysopidae.- *J. Paleont.*, 9: 259-271.
- CROWSON, R. A., 1967: Order Neuroptera.- In: HARLAND et al. (Edit.) *The Fossil Record*. London: Geological Society.
- FRIEDRICH, H., 1953: Neuroptera.- In: BRONN, H. G. (Edit.) *Klassen und Ordnungen des Tierreiches*, 5: 1-148. Leipzig.
- HAGEN, H., 1854: Ueber die Neuropteren der Bernsteinfauna.- *Verh. Zool. Bot. Ges. Wien*, 4: 221-232.
- HARLAND, W. B., COX, A. V., LLEWELLYN, P. G., PICKTON, C. A. G., SMITH, A. G., WALTERS, R., 1982: *A geologic time scale*. Cambridge: University Press.
- HARLAND, W. B., HOLLAND, C. H., HOUSE, M. R., HUGHES, N. F., REYNOLDS, A. B., RUDWICK, M. J. S., SATTERTHWAITTE, G. E., TARLO, L. B., WILLEY, E. C., 1967: *The Fossil Record*. London: Geological Society.
- HENNIG, W., 1969: *Die Stammesgeschichte der Insekten*. Frankfurt a. Main: Waldemar Kramer.
- HENNIG, W., 1981: *Insect Phylogeny*. Chichester, New York, Brisbane, Toronto: John Wiley & Sons.
- HENRIKSEN, K. L., 1922: Eocene Insects from Denmark.- *Danmarks geologiske Undersøgelse*, 37: 1-36.
- HURD, P. D., SMITH, R. F., DURHAM, J. W., 1961: The fossiliferous amber of Chiapas, Mexico.- *Ciencia Communicat. orig.*, 21: 107-118.
- ILLIES, H., 1941: Das Insekten führende Alttertiär von Havighorst.- *Z. Geschiebeforschung Flachlandgeologie*, 17: 16-28.
- JARZEMBOWSKI, E. A., 1980: Fossil insects from the Bembridge Marls, Palaeogene of the Isle of Wight, Southern England.- *Bull. Brit. Museum (Nat. Hist.) Geol. Ser.*, 33: 237-293.
- KRÜGER, L., 1923: Neuroptera succinica baltica.- *Stettiner Entomol. Zeitung*, 84: 68-92.
- LARSSON, S. G., 1975: Palaeobiology and mode of burial of the insects of the Lower Eocene Mo-clay of Denmark.- *Bull. geol. Soc. Denmark*, 24: 193-209.
- LARSSON, S. G., 1978: Baltic Amber - a Palaeobiological Study.- *Entomonograph*, 1: 1-192.
- MacLEOD, E. G., 1970: The Neuroptera of the Baltic Amber. I. Ascalaphidae, Nymphidae, and Psychopsidae.- *Psyche*, 77: 146-180.
- MARTYNOVA, O. M., 1952: Permian lacewings from the USSR.- *Trudy paleont. Inst.*, 40: 187-237 (in Russian).
- MARTYNOVA, O. M., 1962: Order Neuroptera.- In: RODENDORF, B. B. (Edit.) *Arthropoda - Tracheata and Chelicerata*. In: *Textbook of Palaeontology*. 9: 272-282.
- McALPINE, J. F., MARTIN, J. E. M., 1969: Canadian Amber - A palaeontological treasure-chest.- *Canad. Entomologist*, 101: 819-838.
- MEINANDER, M., 1975: Fossil Coniopterygidae (Neuroptera), Notulae Entom., 55: 53-37.
- PANFILOV, D., 1968: Kalligrammatides (Neuroptera, Kalligrammatidae) from Jurassic deposits of Karatau.- In: RODENDORF, B. B. (Edit.) *Jurassic insects from Karatau*: 166-174 (in Russian).
- PANFILOV, D., 1980: New representatives of lacewings (Neuroptera) from the Jurassic of Karatau.- In: DOLIN, V., PANFILOV, D., PONOMARENKO, A., PRITYKINA, L. (Edit.) *Mesozoic Fossil Insects*. Naukova Dumka, Kiev: 82-111 (in Russian).

- PINTO, I. D., 1972: Late Paleozoic insects and crustaceans from Paraná basin and their bearing on chronology and continental drift.- *An. Acad. bras. Ciênc. Supp.*, 44: 247-254.
- PONGRÁCZ, A., 1935: Die eozäne Insektenfauna des Geiseltales.- *Nova Acta Leopoldina (N. F.)*, 2: 485-572.
- RIEK, E. F., 1955: Fossil insects from the Triassic beds at Mt. Crosby, Queensland.- *Australian J. Zool.*, 3: 654-691.
- RIEK, E. F., 1968: Undescribed fossil insects from the Upper Permian of Belimont, New South Wales.- *Rec. Australian Mus.*, 27: 303-310.
- RIEK, E. F., 1974: The Australian Moth-Lacewings (Neuroptera: Ithonidae).- *J. Australian Ent. Soc.*, 13: 37-54.
- RIEK, E. F., 1975: On the Phylogenetic Position of *Brucheiser argentinus* Navás 1927 and Description of a Second Species from Chile (Insecta: Neuroptera).- *Stud. Neotrop. Fauna*, 10: 117-126.
- RIEK, E. F., 1976: New Upper Permian insects from Natal, South Africa.- *Ann. Natal Mus.*, 22: 755-789.
- RIEK, E. F., 1976: The family Stilbopterygidae (Neuroptera) in Australia.- *J. Australian Ent. Soc.*, 15: 297-302.
- RODENDORF, B. B., BEKKER-MIGDISOVA, E. G., MARTYNOVA, O. M., SHAROV, A. G., 1961: Palaeozoic insects from the Kuznez basin.- *Trudy paleont. Inst.*, 85: 1-705 (in Russian).
- RODENDORF, B. B., RASNITSYN, A. P., 1980: The historical development of the class insecta.- *Trudy paleont. Inst.*, 175: 1-269.
- SCHLÜTER, T., 1978: Zur Systematik und Palökologie harzkonservierter Arthropoda einer Taphozönose aus dem Cenomanium von NW-Frankreich.- *Berliner geowiss. Abh. A*, 9: 1-150.
- SCHLÜTER, T., 1982: *Cimbrochrysa moleriensis* n. g. n. sp. und *Hypochrysa hercyniensis* n. sp., zwei fossile Chrysopidae-Arten (Insecta: Planipennia) aus dem europäischen Tertiär.- *N. Jb. Geol. Paläont. Mh.*, 1982: 257-264.
- SCHLÜTER, T., 1984: Palaeontology and evolutionary relationships.- In: CANARD, M., SÉMÉRIA, Y., NEW, T. (Edit.) *Biology of Chrysopidae*. The Hague, Boston, Lancaster: Junk.
- SCHLÜTER, T., STÜRMER, W., 1984: Die Identifikation einer fossilen Rhachiberothinae-Art (Planipennia: Berothidae oder Mantispidae) aus mittelkretazischem Bernstein NW-Frankreichs mit Hilfe röntgenographischer Methoden.- *Progr. World's Neuropterology. Proc. 1st Internat. Sympos. Neuropterology Graz (Austria)*: 49-55.
- TILLYARD, R. J., 1926: Upper Permian insects of New South Wales, Part I. Introduction and the order Hemiptera. Part II. The orders Mecoptera, Paramecoptera and Neuroptera.- *Proc. linn. Soc. New South Wales*, 51: 1-30; 265-282.
- TILLYARD, R. J., 1932: The Lower Permian insects of Kansas. XIV. XV. The orders Neuroptera and Plecoptera.- *Americ. J. Sci.*, 23: 1-30; 97-134; 237-272.
- WEIDNER, H., 1958: Einige interessante Insektenlarven aus der Bernsteininklusen-Sammlung des Geologischen Staatsinstituts Hamburg (Odonata, Coleoptera, Megaloptera, Planipennia).- *Mitt. Geol. Staatsinst. Hamburg*, 27: 50-68.
- WHALLEY, P. E. S., 1980: Neuroptera (Insecta) in amber from the Lower Cretaceous of Lebanon.- *Bull. Brit. Mus. nat. Hist. (Geol.)*, 33: 157-164.
- WHALLEY, P. E. S., 1983: *Fera venatrix* gen. and sp. n. (Neuroptera, Mantispidae) from amber in Britain.- *Neuroptera International*, 2: 229-233.
- WITHYCOMBE, C. L., 1924: Some aspects of the biology and morphology of the Neuroptera. With special reference to immature stages and their possible phylogenetic significance. - *Transacta ent. Soc.*, 1924: 303-412.
- ZALESKY, M. D., 1928: Sur un nouvel insecte neuroptéroïde du Permien du bassin de Kama.- *Bull. Soc. entom. France*, 28: 381-385.
- ZHERIKIN, V. V., SUKACHEVA, I. D., 1973: On the Cretaceous insectbearing "ambers" (resins) of northern Siberia.- *Dokl. ezheg. Chten. N. A. Kholodkovskogo*, 24: 3-48 (in Russian).

ZHERIKIN, V. V., 1980: Class Insecta.- In: Development and changes in mesozoic and cenozoic faunal complexes. Isdatelstvo "Nauka", 40-97.

Address of the author: Dr. Thomas S c h l ü t e r
Department of Geology
University of Dar es Salaam
P.O. Box 35052, Tanzania