

New records of lecideoid lichens from the Southern Hemisphere

by

H. HERTEL

SUMMARY

1. Species are reported from various Southern Hemisphere regions for the first time:
new to the Southern Hemisphere: *Farnoldia similligens*, *Lecidea surculata*, *Lecides umbonata* and *Lecidea verruca*;
new to Uganda: *Tremolecia strata*;
new to South Africa: *Protoparmelia petraecoides*;
new to Bouvetøya: *Carboneas assentiana*;
new to Prince Edward Islands: *Lecidella elaeochroma*;
new to Kerguelen Island: *Lecidella elaeochroma* and *Stephanocyclos hennsenianus*;
new to Heard Island: *Fuscidea astholodes* and *Carboneas assentiana*;
new to New Guinea: *Porpidea albocerulescens* and *Trapelia coarctata*;
new to Norfolk Island: *Lecanora subcoarctata*;
new to New Zealand: *Carboneas assentiana* (Auckland Is.), *Claduzadeana macula*,
Immersaria athroocarpa, *Lecidea capensis*, *Lecidea diducens*, *Schaefferia fuscocinerea*, *Trapelia involuta*;
new to Tasmania: *Carboneas vorticosa*, *Lecanora subcoarctata*, *Lecidea surculata*, *Lecidella stigmatica* and *Poeltiaria turgescens*;
new to Macquarie Island: *Lecidella elaeochroma*;
new to South America: *Bellmeria subcordiza*, *Farnoldia similligens*, *Immersaria athroocarpa*, *Lecidea surculata*, *Lecides umbonata*, *Lecanora subcoarctata* (Galapagos Is.) and *Lecidea verruca*;
new to Ecuador (Galapagos Islands): *Lecanora subcoarctata*;
new to Colombia: *Tremolecia strata*;
new to Peru: *Lecidella stigmatica*;
new to Chile: *Immersaria athroocarpa*;
new to Argentina: *Farnoldia similligens*, *Lecidella stigmatica*, *Lecidea surculata*, *Lecides kalbi*, *Lecidea umbonata*, *Lecidea verruca*, *Sporastatia testudinea* and *Xenolecia spadicommum*;
new to Juan Fernandez Islands: *Poeltiaria corralensis*;
new to the Falkland Islands: *Lecidea surculata*;
new to South Georgia: *Carboneas assentiana* and *C. vorticosa*;

- new to South Orkney Islands: *Carbonea assentiana*;
new to South Shetland Islands: *Carbonea assentiana*, *Lecidella stigmatis*;
new to Antarctica (Antarctic Peninsula): *Carbonea assentiana*, *Fuscoidia dissipabilis*,
Lecidella stigmatis and *Sporastatia polypora*.
2. *Lecidea sciatrapha* Hue is identified as a synonym of *Carbonea assentiana* (xanthone-
deficient chemotype).
3. A neotype is chosen for *Lecidea sciatrapha* Hue.

As part of my research on sedicolous lecideoid lichens of cool and cold climates (i.e. of alpine and polar regions), I began in 1982 to monograph the species of the Antarctic floral zone, defined by CABRERA (1971) to include such areas as Juan Fernandez, Patagonia, New Zealand and Tasmania, as well as Antarctica and the subantarctic islands. A preliminary survey, with a key to some 80 species (HERTEL 1984) was followed by floristic reports on species from New Zealand (HERTEL 1985) and from other parts of the Southern Hemisphere (HERTEL 1987b). Further data were included in some taxonomic accounts of specific groups (HERTEL & KNOPH 1985, HERTEL & RAMBOLD 1990). Additional floristic records, including many new and surprising ones, are presented in this paper, based mainly on the revision of usually undetermined herbarium specimens borrowed from various public herbaria (AAS, AD, AK, ANUC, BG, BM, CANU, CBO, CHR, COLO, GZU, H, HO, LPS, M, MEL, NSW, OTA, PC, S, STU, UPS, W, WIS) as well as of private collections (K. Kalb, G. Kantvilas, H. Mayrhofer, V. Wirth). I wish to thank all the owners and keepers of this herbaria for their support.

I wish to thank Prof. Dr Josef POELT (Graz) and Dr Gerhard RAMBOLD (München) for stimulating discussions, DEUTSCHE FORSCHUNGSOEINENSCHAFT (DFO) for financial support of my research in Antarctic lecideoid lichens, Prof. Dr Christian LEUCKERT (Berlin) for his help in identifying xanthones and Mrs Oedrun SCHWARZE, who carefully did all the TLC work. Last but not least I wish to express my sincere thanks to Dr Gintaras KANTVILAS (Hobart) who followed my studies on Tasmanian lichens with great interest and support and kindly corrected the manuscript linguistically.

(1) *Bellidemora subscorediza* (Lynge) R. Sant.

SANTESSON in MOSERO, Lich. sel. exs. Uppl., no. 30 (1988). - *Aspicilia subscorediza* (Lynge) R. Sant. in VEZDA, Lich. sel. exs., no. 1434 (1976); GALLOWAY, Flora New Zealand Lich. : 23 (1985) - *Lecidea subscorediza* Lynge in DAHL, LYNGE & SCHOLANDER, Skrif. Svalbard Ishavet 70: 22 (1937).

Argentina: Tierra del Fuego: Sierra Alvear, the southern slope, above Las Cotorras (about 20 km ENE of Ushuaia), on exposed rocks in the alpine region, 550 m, 7 February 1940, R. SANTESSON 7228 (B, with *Tremolecia strata*). - Montes Martial bei Ushuaia, oberhalb Acroilla, Silikatfelsen im Krummholzbereich, S- und SE-Hänge, 700-1000 m, 17. Januar 1989, J. POELT (GZU).

This sorediate species, known from alpine and arctic areas in the Northern Hemisphere, is almost constantly sterile and so probably often overlooked. All other Southern Hemisphere records are from New Zealand only (GALLOWAY 1985).

(2) *Carbonea assentiana* (Nyl.) Hertel

HERTEL, Lecideaceae exs., no. 103 (1984); HERTEL, Beih. Nova Hedwigia 79: 442-443 (1984); HERTEL, Mitt. Bot. München 23: 323 (1987). - *Lecidea assentiana* Nyl. in CRUMBIE, Journ. Bot. (London) 13: 334 (1873).

First described from Kerguelen Island, this whitish to pale yellowish lichen is widespread and locally common in subantarctic and maritime Antarctic regions. All individuals contain astranadin (usually with chlorastranarin) and zeorin. However, at least some of the collections from Kerguelen Island and the Prince Edward Islands (where the species is rather common) usually have a more intensely yellowish thallus and also contain two unidentified xanthones (O -methylidichloronorlichexanthone and a O -methylmonochloronorlichexanthone according to Ch. LEUCKERT; see HERTEL 1984: 473). These additional xanthones are lacking constantly in the populations of the maritime Antarctic islands (Bouvet Island, Heard Island, South Orkney Islands, South Shetland Islands) and of the Antarctic Peninsula, where this taxon seems to be one of the most common lecideoid crustose lichen on non-calcareous rocks.

Carbonea assentiana and *C. phaeostoma* seem not to belong to the genus *Carbonea* as circumscribed recently by RAMBOLD (1989). However, because I hesitate to include these two species both in the genus *Lecanora* or in the genus *Adelolecia* (the apothecia of *A. pilati* (Hepp) Hertel & Hafellner as well as those of *C. phaeostoma* contain the anthraquinone 2-chloroemodin), or to create a new genus without additional studies on related groups, I provisionally retain these two species within the genus *Carbonea*.

The distribution of the chemotypes in *Carbonea assentiana* follows the usual pattern for various Northern Hemisphere lichens, where the least complex (most depauperated) chemotypes are restricted to regions with the coldest climates (Arctic, alpine belts of mountain ranges). These xanthone-deficient populations of *Carbonea assentiana* are regarded as such a lichen acid deficient race ("0-strains", LEUCKERT & POELT 1978), which we do not accept as a distinct taxon.

(a) the xanthone-containing chemotype

Prince Edward Islands: Prince Edward Island: Vicinity of Kent Crater, $46^{\circ} 37' - 46^{\circ} 38' S$, $37^{\circ} 53' - 55' E$, loose pebbles on moist ground, c. 130 m, 1 May 1982, H. HERTEL 24309 (M), 24348 (M), 24350 (M), 24356 (M).

Marion Island: Stora Ridge, $46^{\circ} 52' S$, $37^{\circ} 51' E$, 60-70 m, 26 April 1982, H. HERTEL 24221 (M), 24226 (M). - Transvaal Cove, NW of Albatross Lakes, $46^{\circ} 52' S$, $37^{\circ} 51' E$, 50 m, 9 May 1982, H. HERTEL 24542b (M). At Albatross Lakes, 50 m, 9 May 1982, H. HERTEL 24532 (M), 24542a (M). - Long Ridge, between Long Ridge South (lake) and Sea Elephant Bay, c. $46^{\circ} 52' S$, $37^{\circ} 48' E$, 430 m, 23 April 1982, HERTEL 24136 (M). - Western coast, Swartkop Point, $46^{\circ} 56' S$, $37^{\circ} 35' E$, 70 m, 17 May 1982, H. HERTEL 24719 (M). - Along Van den Boogaard River, from foot of Tafelberg to mouth of Rockhopper Bay, $46^{\circ} 53' S$, $37^{\circ} 50' E$, 60 m, 3 May 1982, H. HERTEL 24396a (M).

(b) the xanthone-deficient chemotype

Synonym: *Lecidea sclerophila* Hue, Deux Expédi. Antarct. Franç.: 110-112 (1915); ZAHLBRUCKNER, Catal. Lich. Univ. 3: 686 no. 6711 (1925) ("sclerophila").

Type: Antarctic Peninsula: "îlot Goudier, Port Lockroy, près de l'île Wiencke, chenal de Rosen, sur les roches (diorites), dans les endroits un peu abrités, III^e excursion, no. 86, le 28 décembre 1908" (no material was detectable in PC).

Neotype: Antarctic Peninsula: Wiencke Island, Port Lockroy, Goudier Islet, $64^{\circ} 50' S$, $63^{\circ} 31' W$, halfway up island, 6 m alt., on N facing vertical grandiorite rock (confluent patches diffusely covering an extensive area, 14 December 1944, I.M. LAMB, Operation Tabarin, no. 1883 (BM)).

Auckland Islands: Frenchs Island: Off Crozier Point, in grey zone at h.w.m., 29 December 1962, P.W. JAMES 763 (BM).

South Georgia: Bird Island: SW slopes of Roché Peak (GR 032 155), on dry schistose rock facing SW, exposed, 12 November 1971, D.C. LINDSAY 3062 (AAS). - Right-Whale Bay, behind N end of Binder Bay (GR 054 154), on dry E-facing cliff, c. 30 m, 4 January 1972, D.C. LINDSAY 3801 (AAS). - S end of Binder Bay (GR 054 154), on E-facing nitrogenous outcrop, 5 m, 3 January 1972, D.C. LINDSAY 3771 (AAS). - Royal Bay, near waterfall on SW side of Whale Valley (GR 158 098), on dry non-nitrogenous boulder, 30 m, 17 January 1972, D.C. LINDSAY 3948 (AAS). - South shore of Larson Harbour (GR 163 062), on bird-perching rock by shore, 3 m, 30 December 1971, D.C. LINDSAY 3733 (AAS). - North side of Wink Bay (GR 173 073), on rock on dry south-facing non-nitrogenous scree slope, 100 m, 27 December 1971, D.C. LINDSAY 3669 (AAS).

Heard Island: Steep hill between Rogers Head, Corinthian Bay, West Beach and the foot of the glacier from Big Ben (Kaiser Wilhelm Peak), $53^{\circ} 05' S$, $73^{\circ} 24' E$, 28 November -2 December 1929, R.A.N.Z.A.R.E. no. B 140-15 (M, fragm. ex hb. C.W. Dodge; det. C.W. DODGE as "Lecidea subarctica de Less."); no. B 140-23 (AD, det. C.W. DODGE as "Lecidea subassentans Nyl."). - Cape Laurens, Laurens Peninsula, scree slope facing Atlas Cove, c. 100 m SE of waterfall close to Vanhoffen Bluff, 60-120 m, very protected face, constant moisture from waterfall, January 1963, A. GILCHRIST 19 (MEL 1026848). - Mt Olsen, Laurens Peninsula, on terrace of black sand and rock, 210-250 m, January 1963, A. GILCHRIST 33 (MEL 1026861). - Mt Andrew, in hollows between mounds of Azorella setigera, sheltered area, 100-250 m, January 1963, A. GILCHRIST 2 (MEL 1026852), A. GILCHRIST 5-7 (MEL 1026844). - Mt Drygaliski, NE face, 120-150 m, wind protected, constantly moist area, January 1963, A. GILCHRIST 32 (MEL 1026846).

Bouvetanya: Nunatak ridge E of Rustadkollen, dry basalt slab scree facing E, $54^{\circ} 25' S$, $3^{\circ} 18' E$, 1 January 1979, T. ENOELSKJÖLN 340 (BG, M).

South Orkney Islands: Laurie Island: A small plateau on the N.W. seaward edge of Cape Geddes, $60^{\circ} 42' S$, $44^{\circ} 35' W$, steep slopes down to Brown's Bay, 50 m, 8 April 1946, Falkland Islands Dependencies Survey, no. C3-1001 p.p.t.e. (BM).

Coronation Island: Saunders Point, 0-9 m, 8 October 1950, W.J.L. SLADEN H612/40 (BM).

Lynch Island: On rock fragment near gull rockery, $60^{\circ} 40' S$, $45^{\circ} 38' W$, 10 m, 16 March 1966, D.C. LINDSAY 978 (AAS).

Sigry Island: 90 m S of hill, 30 m alt., $60^{\circ} 42' S$, $45^{\circ} 37' W$, 13 January 1948, P.E. BIOOS, Falkland Islands Dependencies Survey, no. H81-1 (BM). - Marble Knolls, Three Lakes Valley, $60^{\circ} 42' S$, $45^{\circ} 37' W$, 29 October 1966, D.C. LINDSAY 1345 (AAS).

South Shetland Islands: King George Island: Level top of cliff N. of Base Hutt, Martel Inlet, $62^{\circ} 03' S$, $58^{\circ} 23' W$, 7 m, 1 March 1947, D. NICHOLSON & A. REECE, Falkland Islands Dependencies Survey, no. O36-2 (BM). - Admiralty Bay area, creeping slopes above Paradise Cove, on scree on southern slope, in mass, 80 m, volcanic rock, 26 January 1980, R. OCHYRA 1228/80 (M). - Admiralty Bay area, Smok Hill in Martel Inlet, 5 m, on slightly moist andesite rocks, 18 February 1980, R. OCHYRA 2062/80 (M). - Admiralty Bay area, Agat Point S of Sphinx Hill, $62^{\circ} 11' 30'' S$, $58^{\circ} 26' W$, 15 m, on basalt rocks enriched with nitrogen in nestage of sea gulls, 8 January 1980, R. OCHYRA 365/80 (H). - Admiralty Bay area, Demay Point, on boulders on scree, northern slope, 40 m, 27 January 1980, R. OCHYRA 1364/80 (M). - N side of Barton Peninsula, $62^{\circ} 13' S$, $58^{\circ} 44' W$, 3 m, on pebbles of raised beach, 22 January 1986, D.C. LINDSAY 746 (AAS), 751 (AAS).

Penguin Island: Northern tip of the island, $62^{\circ} 05' S$, $57^{\circ} 55' W$, on N facing volcanic rock, 10 m, 27 January 1966, D.C. LINDSAY 804 (AAS), 807 (AAS).

Robert Island: Coppermine Cove, $62^{\circ} 22' S$, $59^{\circ} 45' W$, 60 m, dry N.E. slopes with good shelter, 31 December 1934, "Discovery" Expedition 1934-35, no. 1485d (BM). - Edwards Point, $62^{\circ} 29' S$, $59^{\circ} 30' W$, 10 m, ridge of S-facing raised beach, 5 January 1966, D.C. LINDSAY 613 (AAS).

Deception Island: From area of small ash with a few stones, 6 January 1936, British Graham Land ("Penola") Expedition 1934-37, no. 1351a (BM - no lichen products). - 16 January 1936, British Graham Land ("Penola") Expedition 1934-37, no. 1381 (BM). - 5 February 1945, E.H. BUCK, Operation Tabarin, no. 2358 (BM). -

Antarctica: East Graham Land: Trinity Peninsula, Hope Bay, summit of Pyramid Peak, $63^{\circ} 26' S$, $57^{\circ} 01' W$, 575 m, 30 April 1945, I.M. LAMB, Operation Tabarin, no. D-2420 (BM, neben *Buekia illustrabilis* Lamb, *Acarospora chlorophana* (Wahlenb.) Massal., *Miniquidicus* spec.).

West Graham Land: Goudier Islet (see neotype). - Doumer Island, $64^{\circ} 51' S$, $63^{\circ} 35' W$, rocky outcrop on N. side, c. 30 m, N. exposure, on granodiorite rocks, 12 December 1944, I.M. LAMB, Operation Tabarin, no. 1904 (BM). - Argentine Islands, Galindez Island, $63^{\circ} 15' S$, $64^{\circ} 15' W$, from nearly vertical low sea cliff, 2-8 m, 31 March 1935, British Graham Land ("Penola") Expedition 1934-37, no. 1116nn (BM). - Berthelot Islands, Green Island, $65^{\circ} 19' S$, $68^{\circ} 15' W$, from N facing inland cliff, 18 March 1935, British Graham Land ("Penola") Expedition 1934-37, no. 1061w (BM). -

(3) *Carbonea phaeocystoma* (Nyl.) Hertel

HERTEL, Beih. Nova Hedwigia 79: 443-444 (1984); HERTEL, Mitt. Bot. München 23: 324-325 (1987). - *Leclides phaeocystoma* Nyl. in CROMBIE, Journ. Bot. (London) 13: 334 (1875).

Heard Island: Mt Andree, in hollows between mounds of *Azorella selago*, in a sheltered area, 100-250 m, January 1963, A. GILCHRIST 3-7 (MEL 1026844, a few apothecia beside *Carbonea arvensis*).

Macquarie Island: Bauer Bay, coastal rocks, 2-3 m, 10 January 1972, R. HNATIUK 11712 (MEL 1027283), 11714 (MEL 1027297), 11725 (MEL 1027300). - Approx. half way along the W side of Lake Strahan (Gratitude Lake), 180 m, pebbles embedded in moss, 4 February 1964, R. FILSON (5935) & J. PHILLIPS (MEL 1041279). - North Head of Macquarie Island, on exposed rock in a westerly aspect, 15 January 1964, I.M. LAMB, hb. Filson no. 5724 (MEL 1048265). - W of Aurora Cave, coastal rock, 21 January 1972, R. HNATIUK 11812 (MEL 1027262, III treated specimen with almost pure black discs).

All the specimens contain stranorin and zeorin in the thallus, and 2-chloroemodin (by TLC) in those apothecia in which the reddish pruina of the epiphymenium is well developed.

(4) *Carbonea vitellinaria* (Nyl.) Hertel

HERTEL, Mitt. Bot. München 19: 442 (1983) - *Leclides vitellinaria* Nyl., Bot. Notiser : 177 (1852) - *Napolechia vitellinaria* (Nyl.) Rehm in RABENHORST, Krypt. Deutschland, 2. Aufl. 1 (3): 318 (1890).

New Zealand: Westland: Tasman Valley, ca. $43^{\circ} 07' S$, $170^{\circ} 13' E$, on rock outcrop on ridge near Bell Hut, 8 April 1966, B.A. FINERAN 2325 (CANU) and 2330 (CANU).

(5) *Carbonea vorticosa* (Flk.) Hertel

HERTEL, Mitt. Bot. München 19: 442 (1983) and 21: 306-308 (1985); HERTEL, Beih. Nova Hedwigia 79: 444-445 (1985); RAMBOLD, Biblioth. Lichenol. 34: 83 (1989) - *Leclides vorticosa* (Flk.) Koerb., Syst. Lich. Germ.: 251-252 (1855).

New Zealand: Westland: Mt Haast, Col between West and Middle Peaks, 2355 m, 10 January 1967, R.G. CUNNINGHAME (CANU, hb. FINERAN 2482, cryptothalline thallus with a few apothecia only).

Otago: St Bathans Range, 1675 m, 10 January 1972, C.D. MEURK (BM).

Tasmania: Mt Wellington, on dolerite, alpine rocks, 1260 m, 10 February 1984, P. JAMES & O. KANTVILAS 301/84 (hb. Kantvilas, a few apothecia associated with *Tremolecia strata*).

South Georgia: About 10-15 m below "Summit 706 m" due E of Swinhoe Peak above Hamburg Glacier, on block scree on S-facing slope, probably never receiving direct sunshine, 6 November 1976, R.I. LEWIS SMITH 2526 (AAS).

Antarctica: Dronning Maud Land: Skigården, h. 1651, granite, 1 February 1985, T. ENOEKSKJÖN s.n. (BO).

(6) *Clauzadea mucula* (Th. Teyl.) Coppins & Rambold

in RAMBOLD, Biblioth. Lichenol. 34: 85 (1989) - *Lecidea mucula* Th. Teyl. in MACKAY, Flora Hibernica 2: 115 (1836).

New Zealand: Canterbury: Phipps Peak, summit, 1905, on rock, near Arthur Pass, ca. 42° 54' S, 171° E, 5 March 1966, B.A. FINERAN 2275 (CANU, associated with *Lecidea lapicida* (Ach.) Ach.).

(7) *Farnoldia* cf. *micropora* (Massal.) Hertel

HERTEL, Mitt. Bot. München 19: 443 (1983) - *Lecidea micropora* Massal., Atti I.R. Istr. Veneto, ser. 3, 2: 368 (1856). - *Tremolecia nivalis* (Anzi) Hertel, Khumbu Himal 6 (3): 334-336 (1977) - *Lecidea rhætica* Hepp ex Th. Fr., Lich. Actiol 209 (1860); HERTEL, Beih. Nova Hedwigia 24: 92-96 (1967).

South Shetland Islands: Deception Island: From area of small ash with a few stones, rock with traces of CaCO₃ (HCl weakly positive), 6 January 1936, British Graham Land ("Penola") Expedition 1934-37, no. 1251a (BM), 13511 (BM). This specimen is most probably conspecific with one from Greenwich Island which was determined by HERTEL (1984 p. 430) as *Farnoldia dissipabilis* (Nyl.) Hertel. *Farnoldia micropora* and *F. dissipabilis* are very closely related. They are identical in ascocarp characters and they differ in their highly variable shape and the colour of the thallus areoles. More material from Antarctic regions is necessary for a better understanding of the variability of this species.

Half Moon Island: Hill south of the hut, 62° 35' S, 59° 55' W, on basaltic rock facing WNW, 30 m, 5 January 1966, D.C. LINDSAY 600 (AAS, with *Lecidella stigmatis*). This specimen is provisionally included under *Farnoldia micropora* for I am not absolutely sure whether it represents a distinct taxon. Its epilithic thallus is poorly developed (small isolated white areoles, 0.1-0.3 mm in diam., 0.1 mm thick) and is I-negative. The very glossy black apothecia are small (up to 0.55 mm in diam.) with a distinct margin only when young. Larger apothecia have a shallow sunken zone in the centre of the discs. The hymenium is 70-80 µm, spores 16-19-23 x 8-9-10 µm. The substrate give no CO₂-reaction with HCl, however basalt often bears a slightly calciphilous lichen flora.

Antarctica: Antarctic Peninsula: West Graham Land, Port Lockroy, Couder Inlet, 64° 50' S, 63° 31' W, on limpet shells among loose stones on ground until recently snow covered, 8 m, 22 December 1944, L.M. LAMP, Operation Tabarin, no. 2080 (BM). TLC: no lichen acids. A very remarkable record of this pioneer lichen, hitherto known only from holartic regions (mapped in HERTEL, Mitt. Bot. München 12: 135, 1975). This calciphilous lichen occurs in a habitat which is not covered by snow for only a few weeks each year. It colonizes shells laying on the ground in an area completely formed by acid

rock. This record demonstrates the efficiency of long range dispersal by ascospores.

(8) *Farmakidix similigena* (Nyl.) Hertel

HERTEL, Mitt. Bot. München 19: 433 (1983) - *Lecidea similigena* Nyl., Flora 64: 451 (1881); HERTEL, Herzogia 1: 328 (1969) - *Lecidea subRACTICIA* Arnold ex LETTALU, Feddes Report. 56: 214 (1954); HERTEL, Beih. Nova Hedwigia 24: 97-98 (1967).

Argentina: Tierra del Fuego: Dept. Ushuaia, Montes Martial bei Ushuaia oberhalb Acrosilla, Siliatfelsen im Krummholzbereich, S- und SE-Hänge, 700-1000 m, über kalkhaltigem Schiefer, 17. Januar 1989, J. POELT (OZU, associated with *Lecidea umbonata*).

This well-developed specimen has a whitish thallus composed of dispersed, plane, more or less isolated areolas, 0.2-0.7 mm in diam. The medulla reacts I+ intensely violet; apothecia are up to 0.7 mm in diam., black, plane, with a distinct margin; hymenium is 55-65 µm; subhymenium is very pale blue-green; hypothecium is brown black. Spores 11-15.0-19 x 5.0-5.6-7.0 µm.

The small apothecia, the very dark hypothecium and rather low hymenium are remarkable.

(9) *Furcidea asboloides* (Nyl.) Hertel & V. Wirth

in HERTEL, Beih. Nova Hedwigia 79: 454 (1984); HERTEL, Mitt. Bot. München 21: 308-309 (1985) et 23: 325 (1987) - *Lecidea asboloides* Nyl. in CROMBIE, Journ. Bot. (London) 14: 21 (1876).

Heard Island: South Barrier Plateau, on decayed lava flow, facing E, sheltered from W, 3 February 1965, J. CRICK (MEL 1020728, beside *Carbonea phaeostoma* and *Tephromela atrata* agg.).

Auckland Islands: Auckland Island: Summit of Mt Eden, 420 m, 31 December 1962, P.W. JAMES 833 (BM).

Macquarie Island: Top of Gadgets Gully, in moist position near creek, 11 April 1972, R. WATERHOUSE A38 (MEL 1020915, beside *Lecidea charadriodes*). - Hill on E side of Lake Prior, on rocks in an easterly aspect at an altitude of 275 m at station 6, 23 January 1964, N. BARRET & R. FILSON 3791 (MEL 1048264). - Mouth of Flat Creek on side of rock stack about 20-40 m NE of beach, 6 m, 18 November 1966, K. SIMPSON E25a (MEL 1000260). - NE side of Lake Major, in westerly aspect at Station 10, 240 m, 19 February 1964, R. FILSON 6165 (MEL 1048289).

(10) *Immersaria athroocarpa* (Ach.) Rambold & Pietachmann

in RAMBOLD, Biblioth. Lichenol. 34: 239 (1989) - *Lecidea athroocarpa* (Ach.) Ach. Method. Lich. 41-42 (1803); HERTEL, Khumbu Himal 6(3): 228-230 (1977) - *Porpidea athroocarpa* (Ach.) Hertel & Rambold in Hertel, Lecideac. exs., fasc. VIII, 8, no. 156 (1983) - *Amypgdalaria athroocarpa* (Ach.) BANNAULT & ROUX, Anal. Jard. Bot. Madrid 44(2): 221-225 (1987) [comb. inval.].

Chile: Prov. Malleco: Cordillera de Nahuelbuta, Los Alpes, Pundo Solano (near southern boundary of the province), on granite rocks in forest, 1200 m, 13 January 1958, W.J. EVERDAM 10764 (COLO L-44418).

New Zealand: Canterbury: Kirkliston Range, 1670–1830 m, 25 March 1978, D.J. GALLOWAY s.n. (BM). – Foggy Peak, 64 km WNW of Christchurch, 43° 17' S, 171° 45' E, 1972, O.C. BRATT 72/1963 (HO 38880). – Foggy Peak, 1200–1500 m, 25 January 1985, H. MAYRHOFER 7473 (GZU). – Mt Peel, summit, 43° 51' S, 171° 09' E, 1700–1700 m, 16 January 1985, H. HERTEL 29521 (M).

Otago: Flagstaff Hill near Dunedin, rock outcrop (basalt), near summit, 45° 50' S, 170° 28' E, ± 650 m, 1 February 1985, H. HERTEL 30105 (M). – Old Man Range, summit plateau, between Hyde Rock and Obelisk, at foot of "tor 5475", with *Ceratodon purpureus*, 1660 m, 3 February 1985, H. HERTEL 30122 (M).

(11) *Lecanora physciella* (Darb.) Hertel

HERTEL, Beih. Nova Hedwigia 79: 446–447 (1984); HERTEL, Mitt. Bot. München 23: 326 (1987). – *Leclodes physciella* Darb., Brit. Antarct. (Terra Nova) Exped. Nat. Hist. Rep., Botany I: 33–35 (1923).

Antarctica: Victoria-Land: Cape Hallett, on volcanic rock, 72° 18' S, 170° 18' E, 1969, O.L. LANGE (M). – Geike Land (S. Victoria Land), Spring 1900, Southern Cross Expedition, C.E. BROCHGOREVINK (BM).

East Graham Land: Trinity Peninsula, Hope Bay, summit of Pyramid Peak, 63° 26' S, 57° 01' W, 575 m, 30 April 1945, I.M. LAMB, Operation Tabarin, no. 2413 (BM), no. D-2413 (BM), no. D-2416 (BM), no. D-2422 (BM). – Hope Bay, low hill on E. side of bay, 100 m, on loose stone, 16 April 1945, I.M. LAMB, Operation Tabarin, no. 2389 (BM). – Hope Bay, from coast E of Mt Bransfield, 190 m, 15 October 1946, A. REECE, Falkland Islands Dependencies Survey no. D361-2 (BM).

West Graham Land: Port Lockroy, Goudier Islet, 64° 50' S, 63° 31' W, not far from summit of island, 7 m, 17 March 1944, I.M. LAMB, Operation Tabarin, no. 1163 (BM). – Wiencke Island, N buttress of Noble Peak, 64° 48' S, 63° 23' W, 455 m, on loose stones at base of N facing rock face, 20 November 1944, I.M. LAMB, Operation Tabarin, no. 1799 (BM). – Wiencke Island, Snow Comb Hill, 370 m, on N facing rocky slope, 8 October 1944, I.M. LAMB, Operation Tabarin, no. 1328 (BM). – Marguerite Bay, Debenham Islands, Barbara Island, 65° 08' S, 67° 07' W, 3 January 1937, British Graham Land ("Penola") Expedition 1934–37, no. 1542e (BM), no. 1542g (BM). – Debenham Islands: Barry Island, 65° 08' S, 67° 07' W, 26 November 1936, British Graham Land ("Penola") Expedition 1934–37, no. 1498a (BM). – Doumer Island, 64° 31' S, 63° 35' W, rocky outcrop on N. side, c. 30 m, N. exposure, on granodiorite rocks, 12 December 1944, I.M. LAMB, Operation Tabarin, no. 1912 (BM).

Most of the specimens were tested by TLC. They uniformly contain usnic acid, sometimes accompanied by an unidentified fatty acid.

Lecanora physciella seems to be truly endemic to the Antarctic continent, confined to the Continental Antarctic region, reaching and extending only into the southern and eastern parts of the Antarctic Peninsula.

(12) *Lecanora subcoarcata* (Knight) Hertel

HERTEL, Beih. Nova Hedwigia 79: 448 (1984); HERTEL, Mitt. Bot. München 23: 319–321 (1985); RAMBOLD, Biblioth. Lichenol. 34: 319 (1989). – *Leclodes subcoarcata* Knight, Transact. Proceed. New Zeal. Inst. 8: 316–317 (1876); GALLOWAY, Flora New Zealand Lich.: 239 (1985).

Ecuador, Galapagos Islands: Isla Santa Cruz (Indefatigable I.); vicinity of Academy Bay, aerohaline belt along shore E of CDRS, no date, W.A. WEBER (COLO L-41000).

Norfolk Island: [without exact locality], $28^{\circ} 00' S$, $168^{\circ} 00' E$, 24–27 August 1975.

J. & T. GILBERT, hb. BRATT 73/934 (HO 48186).

New Zealand: North Auckland: Thames Co., Slipper Island, Watchman Rock, on maritime rocks, August 1973, B.W. & G.C. HAYWARD s.n. (AK 161262). – 17.5 km ENE of Warkworth, Tawharanui Peninsula, 3.5 km W of Tokatu Pt., $36^{\circ} 22' S$, $174^{\circ} 50' E$, on sea-shore rocks, 25 May 1981, L. TIBELL 11191 (UPS, beside *Tylophallus pachysanthes*).

Tasmania: Furneaux Group, Bass Strait: Babel Island, SE point of the island, on granite outcrop, 12 m, 1 February 1967, J. WHINRAY (MEL 28275). – Little Goose Island, South Coast, c. 60 m from the tip of the southeastern point, 1.8 m, on coastal granite and small boulders, c. 10 m in from high water level, 18 February 1978, J. WHINRAY 972 (MEL 1025639).

Main Island: Sister Bay, c. 11 km S of Dover, $43^{\circ} 27' S$, $147^{\circ} 01' E$, on dolerite, exposed but out of normal wave area, 3 m, 6 July 1968, G.C. BRATT 68/318 (HO 63753). – De Witt Island, on rocks, littoral in salt spray zone, 28 January 1977, J.A. ELIX 3342 (ANUC).

Lecanora subcoeruleata was hitherto known only from New Zealand.

(13) *Leclides astromorio* Knight

HERTEL, Mitt. Bot. München, 21: 321–322 (1985), RAMBOLD, Biblioth. Lichenol. 34: 165 (1989).

New Zealand: Canterbury: Lees Valley road, 7 km N of Oxford, on greywacke rocks in open bush, 460 m, 28 May 1980, J.A. ELIX 8306 (ANUC). – Rangitata Valley, Boundary Stream, $43^{\circ} 47' S$, $171^{\circ} 11' 30'' E$, little river gorge, c. 550 m, 17 January 1985, H. HERTEL 29650 (M). – Cass, on boulders of greywacke in *Dicarlis*–*Festuca* tussock country above the Biological Station, 8 January 1927, G.E. & Greta DU RIETZ 1454:6 (UPS), 1454:7 (UPS). – Banks Peninsula, Kennedy's Bush, $43^{\circ} 39' S$, $172^{\circ} 16' E$, c. 280 m, hill with siliceous outcrops, 1 November 1981, V. WIRTH (10282) & L. GREENFIELD (STU). – Mt Peel, Lynn Stream Valley, Long Speer above Acland's hut, $43^{\circ} 50' 30'' S$, $171^{\circ} 11' E$, 530 m, 15 January 1985, H. HERTEL 29437 (M); 770 m, H. HERTEL 29455 (M). – Lake Pukaki Hotel, on non-calcareous boulders in *Festuca novae-zelandiae*–tussock at the road side, 12 March 1927, G.E. & G. DU RIETZ 2094b (UPS, beside *Rimularia pezophila*).

Otago: Flagstaff Hill near Dunedin, $45^{\circ} 50' S$, $170^{\circ} 28' E$, basaltic rocks near summit, 650 m, with *Lecanora blanda*, 1 February 1985, H. HERTEL (30117) & H. MAYRHOFER (M). – Abbott's Hill, Dunedin, J.S. THOMSON no. 978 (OTA 029455, CHR 160438, both det. ZAHLBRUCKNER as "Leclides fuscoatra f. famosa (Hoffm.) Nyf."). – Old Man Range, summit plateau near Hyde Rock, $49^{\circ} 22' S$, $169^{\circ} 12' E$, 1610–70 m, small rocks lying on ground in scattered alpine tundra, 3 February 1985, H. HERTEL 30368 (M). – Mt Wakkin, on rock, J.S. THOMSON 1551 (CHR 399191).

Tasmania: Mt Barrow Chalet, $41^{\circ} 23' S$, $147^{\circ} 25' E$, 600 m, granite, 30 January 1966, G.C. BRATT 3233 (HO 38855). – Coastal rocks between Shelly Point and Buxton Point S of Swansea (N of Mayfield), $42^{\circ} 14' S$, $148^{\circ} 01' E$, dolerite, 18 February 1985, H. MAYRHOFER (no. 7327) & H. HERTEL (OZU).

All specimens uniformly contain gyrophoric acid.

(14) *Leclides surculatus* Th.Pr.

Th. FRIES, Lich. Arctoi : 213 (1860); HERTEL, Khumbu Himal 6 (3): 233–234 (1977); HERTEL, Mitt. Bot. München 13: 339–342 (1977); M. INOUR, Journ. Sci. Hiroshima Univ., ser. B, div. 2, 18: 26–28 (1982); A. SCHWAB, Mitt. Bot. Mün-

itage Library, <http://www.biodiversitylibrary.org/>; www.b
chen 22: 307-312 (1986).

Argentina: Prov. Mendoza: In vicinis montis Aconcagua, Puerta del Juan (7), 16 February 1903, G.O.A. MALME s.n., Itin. Regnell. Secund. (S).

Falkland Islands (Islas Malvinas): Port Stanley, the quartzite ridge, 8 April 1902, C. SKOTTSBERG (S).

Tasmania: Crater Peak, alpine moorland, on quartzite, 1200 m, 16 February 1984, P. JAMES & G. KANTVILAS 425/84 (hb. Kanvilas, M).

Lecidella surculata has its centre of distribution in arctic and subarctic regions where it is locally quite common. It is closely related to *Lecidella diducens* Nyl. The main difference between these species is the presence of 2'-O-methylanizide acid (causing the C-red reaction of the excipulum) in *Lecidella diducens*. Although both chemotypes are sympatric over wide parts of their range, they do not appear to be identical in their ecology and chorology.

(15) *Lecidella capensis* Zahlbr.

ZAHLBRUCKNER, Catal. Lich. Univ. 3: 532 no. 6345 (1925) nom. nov.; RAMBOLD, Biblioth. Lichenol. 34: 168 (1989) - *Lecidella ochroidea* Stiz. Ber. Thätigk. St. Gall. Naturw. Ges. 1889-90: 159-160 (1891) nom. illegit. (sed non *Lecidella ochroidea* Ach. 1803).

New Zealand: Nelson: Kahoka Coast, coastal forest near Kahoka Lakes, 40° 33' S, 172° 36' E, 12 December 1982, J.K. BARTLETT 22862 (M, containing 2'-O-methylperistolic acid).

The species usually has a pale yellowish brown thallus and superficially resembles *Lecidella fuscoatra* (L.) Ach., it was hitherto known from South Africa (type) and Australia (RAMBOLD 1989) and is a new record for New Zealand.

(16) *Lecidella diducens* Nyl.

NYLANDER, Flora 48: 148 (1865); HERTEL, Herzogia, 2: 43-44 (1970); HERTEL, Khumbu Himal 6 (3): 243-244 (1977); HERTEL, Mitt. Bot. München 13: 340, 343-344 (1977); SCHWAB, Mitt. Bot. München 22: 322-324 (1986); RAMBOLD, Biblioth. Lichenol. 34: 208 (1989).

New Zealand: Canterbury: South-western slopes of Foggy Peak, Torlesse Range, on exposed greywacke rocks in alpine fell field, 1040 m, 23 March 1980, J.A. ELIX 7665 (ANUC). - Torlesse Range, Porter Pass W of Springfield, 43° 17' S, 171° 44' E, c. 1000 m, 22 January 1985, H. MAYRHOFER 7462 (OZU), 7463 (OZU). - Mt Peel, Lynn Stream Valley, Long Speer above Acland's hut, 43° 59'30" S, 171° 11' E, trail to summit of Mt Peel, c. 650 m, 15 January 1985, H. HERTEL 29421, 29452 (M).

Otago: Mt Maungatua, 45° 54' S, 170° 06' E, rock outcrop at swampy summit plateau, c. 890 m (subalpine/alpine situation), 31 January 1985, H. HERTEL 30058 (M), 30063 (M), 30079 (M).

This bipolar species is well known from Holarctic regions (see HERTEL 1977b: 340). It was first reported from the Southern Hemisphere (Australia) by RAMBOLD (1989) and is a new record for New Zealand. All specimens contain 2'-O-methylanizide acid in the excipula of the saccocarps, causing a bright C-red reaction.

(17) *Lecidea fuscoastrula* Nyf.

NYLANDER, Lich. Novae Zelandiae : 106 (1888); HERTEL, Beih. Nova Hedwigia 79: 418, 482 (1984); RAMBOLD, Biblioth. Lichenol. 34: 173 (1989).

New Zealand: Canterbury: Lowry Peaks Range, Mt Palm, 42° 41' S, 173° 06' E, 6 km ESE of Waiata, ± 300 m, open tussock grassland area, on sandstone outcrop, 4 December 1980, L. TIBELL 9875 (UPS). - Banks Peninsula, on outcropping siliceous rocks facing SE, 1 January 1981, L. TIBELL 10165 (UPS). - Banks Peninsula, Kennedy's Bush, 43° 39' S, 172° 36' E, hill with siliceous outcrop, ± 200 m, 1 November 1981, V. WIRTH (11152) & L. GREENFIELD (STU), and V. WIRTH (11157) & L. GREENFIELD (STU). - Banks Peninsula Scientific Reserve, Canterbury Plains between Selwyn and Rakaia River, c. 6 km SSW of Dunsandel, 65 m, loose pebbles on ground, 14 January 1985, H. HERTEL 29211 (M). - Rangitata River Co., Stew Point, 43° 43' S, 171° 05' E, rock outcrop on steep, sun exposed, heavily grazed slope, 17 January 1985, H. HERTEL 29241 (M), 29642 (M), 29633 (M), 29644 (M). - Summit of Mt Peel, 43° 51' S, 171° 09' E, 1730 m, schist outcrop, 16 January 1985, H. HERTEL 29522 (M), 29587 (M). - Lake Ohau, Lake Ohau Lodge, associated with *Lecanora subgiscuoides* (Krempehl.) Zahlbr., February 1938, MASON 387a (OTA, Murray Collection).

Otago: Flagstaff Hill near Dunedin, 45° 50' S, 170° 28' E, rock outcrop (basalt) near summit, ± 650 m, 1 February 1983, H. HERTEL 30102 (M). - Summit of Silver Peak (Pulpit Rock), 45° 45' S, 170° 26'30" E, W of Waitati, horizontal schist layers near ground, 760 m, 29 January 1985, H. HERTEL 32345 (M). - Old Man Range, summit plateau near Hyde Rock, 45° 21' S, 169° 12' E, stones on ground in alpine tundra vegetation, ± 1650 m, 3. February 1985, H. HERTEL 30363 (M). - Old Man Range, Fruitlands, schist rocks, 465 m, 6 February 1981, C.D. MEURK & V. WIRTH (ib. Meurk).

(18) *Lecidea kahli* Hertel

HERTEL, Beih. Nova Hedwigia 79: 418-419 (1984).

Chile: Tierra del Fuego: Porvenir, on stones on the seashore NW of the town, in the *Catoplaea sublobulata*-belt, 28 December 1940, R. SANTESSON 6586 (S).

Argentina: Tierra del Fuego: Dept. Rio Grande, *Nothofagus antarctica*-Wald und windgefeigter Rücken etwa 3 km E Estancia Despedida (WSW Rio Grande), 6. Januar 1989, J. POELT (OZU).

Until now, *Lecidea kahli* was known only from the type collection from Chile (Ultima Esperanza, Torres del Paine). The species resembles *Lecidea astrobrunnea* in its general appearance, but differs in its larger spores and C+ red cortex and excipulum (due to the presence of gyrophoric acid). The hypothecium is completely unpigmented and the medulla I negative. The interior part of the excipulum is also unpigmented but appears opaque pale brownish when the sections are observed in water. This is because the hyphae are densely impregnated with minute, pale brownish crystals containing gyrophoric acid which dissolve completely in KOH and turn reddish in hypochlorite solution. The hymenium is 45-50 µm tall, with a dark greyish olive epihymenium. Spores are often poorly developed and are 8-10-13 x 4-5.7-6.5 µm.

(19) *Lecidea lapicida* (Ach.) Ach.

HERTEL, Beih. Nova Hedwigia 79: 419-420 (1984); HERTEL, Mitt. Bot. München 21: 323-325 (1985), 22: 326-327 (1987); O. SCHWARZ, Mitt. Bot. München 22: 324-342 (1986) - RAMBOLD, Biblioth. Lichenol. 34: 210 (1989).

Although intermediate specimens containing both stictic and norstictic acids in significant quantities occur occasionally (SCHWAB 1986) and both chemical strains are sympatric over wide regions there is a remarkable geographical separation in cool temperate regions. Therefore, I tend to follow WIRTH (1987 p.511), who treats these two chemotypes as varieties of *Lecidella lippiae*: the stictic acid chemotype as var. *lippiae*, the norstictic acid chemotype as var. *pantherina*.

var. *lippiae*

Argentina: Staten Island: Port Cook, 1888, C. SPEGAZZINI no. 63 (LPS 35146, det. C.W. DODGE 1970 as "Lecidella impolita MUL. Arg.").

Tierra del Fuego: Porvenir, Morro Piedra, on boulder in *Chiliotrichum* heath, 300 m, 31 December 1949, R. SANTESSON 5355 (S), 5356 (S, beside *Rimularia psephotis*). - Sierra Alvear, the southern slope, above Las Cotorras (about 20 km ENE of Ushuaia), on exposed rocks in the alpine region, c. 350 m, 7 February 1940, R. SANTESSON 939 (S). - Sierra Alvear, Berg SE Monte Alvear, 220-1200 m, N vom Restaurant Tierra Mayor, 14. Januar 1989, J. POELT (GZU). - Parque Nacional de Tierra del Fuego, Juárez Ende der Bahía Lapataia, felsige Küste, 0-5 m, 13. Januar 1989, J. POELT (GZU). - Dept. Río Grande, N-Rand der Anden, Cerro Chenen E Hacienda Chenen, felsiger NE-Hang, 200-230 m, 8. Januar 1989, J. POELT (GZU, 2 Proben).

New Zealand: Canterbury: Phipps Peak, rock at summit, 1905 m, near Arthur Pass, ca. 42° 54' S, 171° E, 5 March 1966, B.A. FINERAN 2273 (CANU, det. C.W. DODGE as "Lecidella dumetina Zahlbr."). - Phipps Peak, ridge to summit, 1830 m, on rock, 5 March 1966, B.A. FINERAN 2263 (CANU, det. C.W. DODGE as "Lecidella dumetina Zahlbr."). - Banks's Peninsula, Devil's Cap (between Lake Forsyth and Akaroa Harbour), 43° 49' 30" S, 172° 49' 30" E, ± E exposed basaltic rocks near summit, 670-710 m, 19 January 1985, H. HERTEL 29693, 29694 (M). - Summit of Mt Peel, 43° 51' S, 171° 09' E, 1740 m, exposed rock surface, 16 January 1985, H. HERTEL 29586 (M).

Otago: Rock and Pillar Range, schist rock top, 1190 m, 18 September 1981, C.D. MEURK ML16 (hb. Meurk). - Maungahau Summit, rock, J.S. THOMSON 1374, ZA 289 (CHR 347036). - Old Man Range, Obelisk, 45° 19' S, 169° 13' W, near Fruitlands, 1690 m, alpine *Celmisia*-*Racoulea* vegetation, siliceous rocks, 6 November 1981, V. WIRTH (11317) & C.D. MEURK (STU). - Old Man Range, Obelisk Range, 1670 m, 3 February 1985, H. HERTEL 30342 (M). - The Remarkables, Double Cone above Lake Alta, 45° 04' S, 168° 48' E, 2250-2280 m, exposed rocks near summit, 5 February 1985, H. HERTEL 30440 (M), 30343 (M), 30435 (M), 30439 (M).

Southland: West Dome, on summit rocks, 1230 m, 20 June 1982, C.D. MEURK ML21 (hb. Meurk).

Tasmania: Hounslow Heath, schist rock outcrops in alpine heath, 1120 m, 17 February 1984, P. JAMES & O. KANTVILAS 344/84 (hb. Kantvilar).

South Georgia: About 10-15 m below "Summit 706 m" due E of Swinhoe Peak above Hamburg Glacier, on block scree on S-facing slope, probably never receiving direct sunshine, 6 November 1976, R.I. LEWIS SMITH 2331 (AAS). - Bird Island: slopes to east above Jordan Cove (GR 032 154), on schistose rock, dry but exposed 25° slope, aspect 230°, 150 m, 11 November 1971, D.C. LINDSAY 3005 (AAS), 3011 (AAS), 3015 (AAS). - Right Whale Bay, c. 1 km SW of Nameless Point (GR 054 137), on moss in pockets of boulder on scree, 200 m, 5 January 1972, D.C. LINDSAY 3823 (AAS). - Cumberland East Bay, Brown Mountain, rocks to west of summit (GR 132 123), on dry exposed greywacke rocks facing west, 300 m, 26 November 1971, D.C. LINDSAY 3233 (AAS). - Cumberland East Bay, Hope Point (GR 133 124), on moderately sheltered dry, non-nitrogenous greywacke outcrop facing south, 50 m, 24 November 1971, D.C. LINDSAY 3187 (AAS), 3193 (AAS). - Cumberland East Bay, c. 1 km W of Mt Duse (GR 133 127), 350 m, exposed dry greywacke outcrop facing W, 23 November 1971, D.C. LINDSAY 3133 (AAS), 3134 (AAS), 3141 (AAS, beside *Tephromela* str.). - Royal Bay, near waterfall on SW side of Whale Valley (GR 158 098), on dry non-nitrogenous boulder, 30 m, 17 January 1972, D.C. LIND-

SAY 3947 (AAS), 3948 (AAS, beside *Carbonaria assentiana*). - 1 km SE of Brocken (GR 162 101), 180 m, on dry non-nitrogenous E-facing outcrops, 23 January 1972, D.C. LINDSAY 4028 (AAS). - 0.5 km W of Cape Vahsel (GR 177 071), on west-facing dry slightly nitrogenous outcrop, 150 m, 28 December 1971, D.C. LINDSAY 3704a (AAS).

Antarctica: West Graham Land: Argentine Islands: Galindez Island, 65° 13' S, 64° 15' W, from typical boulder area, just uncovered by snow, 27 December 1935, British Graham Land ("Pencos") Expedition 1934-37, no. 1328m (BM).

var. *pantherina* Ach.

ACHARIUS, Kongl. Svensk. Akad. Nya Handl.: 232 (1808) - *Leucidea pantherina* (Ach.) Th.Fr., Lich. Scand. : 491-493 (1874).

- *Leucidea lactea* Flk. ex Schaefer., Lich. Helvet. Specii: 127-128 (1828) - *Leucidea lapicida* var. *lactea* (Schaefer.) V. Wirth, Die Flechten Baden-Württembergs: 511 (1987).

New Zealand: Nelson: Upper Cobb Valley, headwaters of Burgoon Stream, 1180 m, 16 December 1983, J.K. BARTLETT 27024 (M).

Canterbury: Phipps Peak, ridge to summit, 1830 m, 5 March 1966, B.A. FINE-RAN 2264 (CANU, det. C.W. DODGE as "Leucidea subtenuibrosa Nyl."). - Mt Peel, summit, 43° 51' S, 171° 09' E, 1600-1700 m, 16 January 1985, H. HERTEL 29505 (M).

Otago: North Old Man Range, summit, 1650 m, 6 November 1981, C.D. MEURK ML15 (hb. Meurk). - Rock and Pillar Range, summit rocks, 1430 m, together with *Buellia* sp., on schist rocks near to ground, 28 January 1985, H. HERTEL 32256 (M), 32267b (M).

(20) *Leucidea hygromma* Nyl.

HERTEL, Mitt. Bot. München 23: 327-329 (1987); RAMBOLD, Biblioth. Lichenol. 34: 214 (1989).

New Zealand: Nelson: W of Matakitaki River Bridge, on schist outcrop, 300 m, 24 May 1970, P. CHILD 361 (Herbar P. Child). - Tasman Mts, Lonely Lake, ± 1270 m, 20 December 1983, J.K. BARTLETT (M). - Upper Cobb Valley, headwaters of Burgoon Stream, 1180 m, 16 December 1983, J.K. BARTLETT 27011 (M).

Westland: on moraine boulders of Franz Josef Glacier, 12 March 1966, W. MARTIN (CHR 407423).

Canterbury: Arthur's Pass National Park, Temple Basin, 42° 54' S, 171° 35' E, 1400 m, 1500 m, niedrige, wasserüberdeckte Sandsteinfelsen am Wegrand, 23 January 1985, H. HERTEL 29833 (M), 29849 (M). - Banks Peninsula, on outcropping siliceous rocks facing SE, 1 January 1981, L. TIBELL 10172 (UPS).

Otago: Mt Maungatua SW of Dunedin, 45° 54' S, 170° 08' E, rock outcrops at swampy summit plateau, 880 m, 31 January 1985, H. HERTEL 30046 (M). - Pulpit Rock near Dunedin, 400 m, 16 November 1968, D.J. CALLOWAY (BM). - Old Man Range, SW of Alexandra, summit plateau near Obelisk, 43° 20' S, 169° 12' E, 1670 m, pebbles lying on ground in scattered alpine vegetation, 3 February 1985, H. HERTEL 30338 (M). - 5 to 6 miles S of Sutton, Mongrel-Middlemarch road, schist tor, c. 430 m, 18 September 1981, C.D. MEURK ML18 (hb. Meurk).

Southland: Kiriwao Summit (S. of Clinton), 640 m, 16 March 1983, C.D. MEURK (hb. Meurk). - West Dome, summit ridge, East Peak, c. 1100 m, 20 June 1982, C.D. MEURK ML36 (hb. Meurk).

Auckland Islands: Auckland Island: Summit of Mt Eden, 420 m, 31 December 1962, P.W. JAMES 832b (BM), 832d (BM).

Campbell Islands: Summit rock of Mt Lyall, 400 m, 22 January 1981, C.D. MEURK

56 (BM). - CBF crest E of Mt Dumas, 190 m, exposed stones on peaty fellfield, 31 December 1980, C.D. MEURK s.n. (BM). - Mt Honey, summit, 370 m, 10 January 1981, C.D. MEURK (BM) and 3 February 1981, C.D. MEURK 102 (BM), 104 (BM).

Macquarie Island: Cliffs on the western side of the island a $\frac{1}{4}$ mile S of Double Point, on rocks in a northerly aspect, 180 m, 3 February 1964, R. FILSON 5914 (MEL 1048287). - NW of Mt Owynn, on rock, 240 m, 30.12.1971, R. HNATIUK 11568 (MEL 1027329).

(21) *Lecidella placodiformis* Hue

HUE, Deux Expéd. Antarct. Franç. (1908-1920), Sci. Nat., Docum. Sci., Lichens: 99-101 (1915); HERTEL, Herzogia 5: 455-456 (1981); HERTEL, Beih. Nova Hedwigia 79: 420-421 (1984).

Antarctica: West Graham Land: Wiencke Island, Port Lockroy, Goudier Islet, $64^{\circ} 30' S$, $63^{\circ} 31' W$, near summit of island, 8 m, on igneous rock, 2 April 1944, I.M. LAMB, Operation Tabarin, A1201 (BM) and (same locality): 11 January 1945, I.M. LAMB, Operation Tabarin, no. 1195 (BM), 2237 (BM). - Doumer Island, $64^{\circ} 31' S$, $63^{\circ} 35' W$, rocky outcrop on N side, c. 30 m, N exposure, on granodiorite rocks, 12 December 1944, I.M. LAMB, Operation Tabarin, no. 1912 (BM). - Argentine Islands, SE corner of Winter Island, overlooking Stella Creek, $63^{\circ} 15' S$, $64^{\circ} 16' W$, rock exposures, 15 March 1960, B.J. TAYLOR 153 (AAS). - Galindez Island, $63^{\circ} 15' S$, $64^{\circ} 15' W$, from nearly vertical low sea cliff, 2-8 m, 31 March 1935, British Graham Land ("Penola") Expedition 1934-37, no. 1116a (BM), 1116d (BM), 1116m (BM), 1116p (BM), 1116t (BM), 1116x (BM). - Marguerite Bay, SE corner of Bourgeois Fjord, $67^{\circ} 40' S$, $67^{\circ} 05' W$, 15 December 1936, British Graham Land ("Penola") Expedition 1934-37, no. 1515b (BM). - Lagotellerie Island, Fallières Coast, $67^{\circ} 53' S$, $67^{\circ} 24' W$, N facing rock slope, 35-80 m, 22 February 1977, R.L. LEWIS SMITH 2202 (AAS).

All specimens listed here contain confluentic acid, 2'-O-methylperlatolic acid, and often 2'-O-methylmicrophyllinic acid as the major constituents. In this respect, this Antarctic population differs from *Lecidea strobrunnea*, a morphologically and anatomically indistinguishable species which contains only 2'-O-methylperlatolic acid.

It is remarkable that in this case, in contrast to most other lichens, the chemically richer chemotype is from an area with a polar climate while that with the "impoverished", simpler chemistry is found in temperate climates, e.g. Australia (see RAMBOLD 1989). It is also remarkable that I have not found *Lecidella placodiformis* represented in the collections from the South Shetland Islands, South Orkney Islands or from South Georgia, three relatively well-collected areas of the Antarctic zone, at least with respect to the more conspicuous species.

(22) *Lecidella sarcogynaoides* Koer.

KOERBER, Syst. Lich. Germ.: 252 (1855); HERTEL, Decheniana 127: 56, 72 (1975); HERTEL, Mitt. Bot. München 21: 422 (1984); HERTEL, Beih. Nova Hedwigia 79: 422 (1984); HERTEL, Mitt. Bot. München 23: 329 (1987); RAMBOLD, Biblioth. Lichenol. 34: 185 (1989).

New Zealand: Canterbury: Banks's Peninsula, Campy Bay, in littoral zone of NE facing sea shore rocks, 31 December 1980, L. TIBELL 10147 (UPS).

Otago: Dry hills N of Alexandra, 280 m, $45^{\circ} 14' S$, $169^{\circ} 25' E$, 280 m, siliceous rocks, 6 November 1981, V. WIRTH (no. 11296) & C.D. MEURK (STU).

(23) *Lecidea sphinctocidarum* Hertel

HERTEL, Beih. Nova Hedwigia 79: 422-423 (1964); HERTEL, Mitt. Bot. München 23: 329 (1987).

South Orkney Islands: *Lynch Island*: On rock fragments near gull rookery, 10 m, 60° 40' S, 45° 38' W, 16 March 1966, D.C. LINDSAY 979 (AAS).

Sigry Island: Borge Bay: Largest of the Thule Islands, 60° 42' S, 45° 37' W, on rock fragments near gull nests, 15 m, 13 October 1966, D.C. LINDSAY 1299 (AAS - a specimen absolutely identical with the type specimen in all anatomical characters, but instead of the typical very pale brownish thallus, with a reddish, rusty coloured one, thus resembling somewhat *Lecidea silacea* (Web.) Ach., and therefore determined as "*Lecidea dictyonif*" by LINDSAY).

This is a remarkable species (keyed in HERTEL 1964) characterized by its relatively tall hymenium (70-100 µm) and its slender and large spores.

Spore size medium values:

Collection	length (l) [µm]	width (w) [µm]	l : w
holotype	15-16.3-18	4.5-5.6-6.5	2.9
LAMB 2299	12-16.1-21	5.0-6.7-8.0	2.4
LINDSAY 1090	10-12.4-15	4.5-6.2-7.0	2.0
LINDSAY 1299	12-15.3-19	5.0-6.3-8.0	2.4

Lecidea sphinctocidarum is hitherto only known from five localities on the South Orkney and South Shetland Islands.

(24) *Lecidea umbronata* (Hepp) Mudd

MUDD, Manual Brit. Lich.: 204 (1861); HERTEL, Beih. Nova Hedwigia 24: 37-40 (1967); HERTEL, Herzogia 2: 497-499 (1973) et 4: 379, 398 (1977) - *Bistortum umbronata* Hepp, Flechten Eur. no. 257 (1857).

Argentina: *Tierra del Fuego*: Dept. Ushuaia, Montes Martial bei Ushuaia oberhalb Acrosilla, Siltfelsen im Krummholzberich, S- und SE-Hänge, 700-1000 m, über kalkhaltigem Schiefer, 17. Januar 1989, J. POELT (OZU, several very small thalli beside *Farmoidea*).

This is the first Southern Hemisphere record of this calciphilous species which is closely related to *Lecidea lapicida*. Its Northern Hemisphere distribution is mapped in HERTEL (1977c: 398).

(25) *Lecidea verrucosa* Pisch.

POELT, Mitt. Bot. München 4: 187 (1961); HERTEL, Herzogia 1: 433-434 (1970) and 2: 300-301 (1973); HERTEL, Mitt. Bot. München 12: 142-143 (1975) and 17: 179 (1981).

Argentina: *Tierra del Fuego*: Montes Martial bei Ushuaia, oberhalb Acrosilla, Siltfelsen im Krummholzberich, S- und SE-Hänge, 700-1000 m, 17. Januar 1989, J. POELT (OZU).

Chile: *Tierra del Fuego*: Porvenir, Morro Piedra, on boulder in *Chiliotrichum* heath,

300 m, 31 December 1940, R. SANTESSON 6837 (S), 6858 (S, rich material), 6860 (S, very scanty material).

These are the first records of this species for the Southern Hemisphere. It is not clear whether the lichens are growing parasitically.

(26) *Lecidella elaeochroma* (Ach.) Choisy s. ampl.

CHOISY, Bull. Mens. Soc. Linn. Lyon 19: 19 (1950); HERTEL, Willdenowia 6: 248-249 (1971).

Prince Edward Islands: Marion Island: Transvaal Cove, Meteorological Station, 45° 53' S, 37° 51' E, on old wooden stain of the Geophysical Laboratory building, 6 May 1982, H. HERTEL 24452 (M).

Kerguelan Islands: Observatory Bay, 49° 25'12" S, 69° 53'22" E, on timber of old hut, British Transit of Venus Expedition, 20.-21 February 1930, members of B.A.N.Z.A.R.E., no. B195-1 (AD; det. C.W. DODGE as "Lecidea subpiana Nyl.").

Macquarie Island: Wireless Hill, N Head of Macquarie I., on wood of MAWSON's old Wireless Hut, 15 January 1964, R. FILSON 5715 (MEL 1041275).

The occurrence of this usually corticolous lichen on three, very remote, subantarctic islands with a species-poor, vascular plant flora which does not include trees or shrubs is remarkable. It is unclear, whether *Lecidella elaeochroma* was introduced by man or whether it is a very reproductive pioneer lichen, well-adapted to very long distance dispersal. On the Prince Edward Islands, I saw it only on the timber of huts, but not on natural, woody, plant remnants which are very rare in these perhumid, cool and constantly wind-swept islands, e.g. old roots of *Azorella selago*.

(27) *Lecidella schistiseda* (Zahlbr.) Hertel

HERTEL, Mitt. Bot. München 19: 443-444 (1983); Beih. Nova Hedwigia 79: 450 (1984) - *Lecidea schistiseda* Zahlbr., Denkschr. Akad. Wiss. Wien, math.-natur. Kl. 104: 309-311 (1941).

New Zealand: Otago: Old Man Range, Obelisk near Fruitlands, 45° 19' S, 169° 13' E, 1690 m, alpine Colomia-Raoulia-vegetation, siliceous rock, 6 November 1981, V. WIRTH (no. 11860) & D.D. MEURK (STU).

(28) *Lecidella stigmaria* (Ach.) Hertel & Leuckart

HERTEL, Mitt. Bot. München 21: 325 (1985); RAMBOLD, Biblioth. Lichensol. 34: 112 (1989).

Peru: Dept. Junín: Prov. Tarma, at km 35 on road from Tarma to Jauja (20 km SE of Tarma), 4100 m, calcareous outcrops and cliffs with low spiny shrubs of *Ribes* and *Mahonia*, 28 November 1962, H.H. & C.M. ILTIS LP-13 (WIS).

Argentina: Govt. Santa Cruz: Rio Gallegos, Estancia Olier-Alba, on basaltic rocks on the edge of the meseta just above the valley of Rio Gallegos, 18 January 1940, R. SANTESSON 7096 (S).

Tierra del Fuego: Dept. Rio Grande, N-Rand der Anden, Cerro Chenen E Hacienda Chenen, felsiger NE-Hang, 200-230 m, 8. Januar 1989, J. POELT (OZU).

New Zealand: Canterbury: Cave Stream and Flock Hill above Cave Stream, 650-

750 m, 43° 12' S, 171° 45' E, limestone outcrop in tussock grassland with scattered *Dicrania toumatou*, 9 September 1981, H. MAYRHOFER 2301 (GZU), 2310 (GZU). - Cass, Cass Hill near the University of Canterbury Field Station, 750 m, 43° 02' S, 171° 46' E, *Nothofagus solandri* var. *cliffordioides* forest, 7 September 1981, H. MAYRHOFER 2410 (GZU). - Rangitata River Co., Stew Point, 43° 43' S, 171° 05' E, boulders in the heavily grazed bottom of the valley, 17 January 1985, H. HERTEL 29633 (M).

Otago: Lake Hayes NE of Queenstown, boulders at the shore of the lake, 20 September 1981, H. MAYRHOFER 2246 (GZU).

Southland: Springhill, Sharks Tooth, April 1961, on limestone rocks, J. MURRAY 5847 (OTA), 5848 (OTA). - "Springfield", Otakaroma, 215 m, on rock (sandstone) bluff, with *Tragelia coarctata*, 27 February 1984, C.D. MEURK ML7 (hb. Meurk).

Tasmania: Table Cape, 3 miles N of Wynyard (N.W. coast), on boulders of andesite-olivine-dolerite at S.E. base of headland, against the sea, 9 January 1967, J.H. WILLIS (MEL 1031111 - TLC: stranorin, zeorins).

South Shetland Islands: Half Moon Island, 62° 35' S, 59° 55' W, hill S of the huts, on basaltic rock facing WNW, 30 m, 5 January 1966, D.C. LINDSAY 600 (AAS - a specimen with rather small apothecia, up to 0.6 mm in diam., hymenium 65-75 µm, spores 14-16 x 7-9 µm).

Antarctica: East Graham Land: Ross Island, Cape Oage, rock on talus slope, 6 m, 29 November 1945, I.M. LAMB, Operation Tabarin, no. 2845 (BM).

(29) *Megalospora gompholoma* (Mill.-Arg.) Sipman subsp. *gompholoma*

SIPMAN, Biblioth. Lichenol. 18: 105-106 (1983).

The following specimen is growing directly on rock and was found associated with undetermined herbarium material:

Auckland Islands: Rose Island: On maritime and fresh water rocks, directly on rock, 9 January 1963, P.W. JAMES (BM). - A second record for the Auckland Islands (cfr. SIPMAN 1983: 234).

(30) *Paraporiella leptocarpa* (Bab. & Mitt.) Rambold & Hertel

in RAMBOLD, Biblioth. Lichenol. 34: 259 (1989) - *Lecidea leptocarpa* Nyl. ex Bab. & Mitt. in HOOKER, Flora Tasmaniæ 2: 352, tab. CC, fig. C (1859); HERTEL, Beih. Nova Hedwigia 79: 464-465 (1984); GALLOWAY, Flora New Zealand Lich.: 234-235 (1985).

New Zealand: Auckland: Little Barrier Island, on clay, 30 m, April 1973, D.J. COURT H30.56 (AK). - Thames Co., Rabbit Island, on rocks of cliff tops, August 1973, B.W. & O.C. HAYWARD 49.147 (AK). - Coromandel Peninsula, Mt Te Moehau, Little Moehau, 36° 32' S, 175° 24' E, 850 m, on andesite, 24 February 1983, J.K. BARTLETT 22673 (M), 22678 (M). - Waitakere Coast, Whatipu, 37° 04' S, 174° 30' E, 25 m, on volcanic andesitic rock, 27 October 1983, J.K. BARTLETT 27014 (M). - Rangitoto Island, coastal rock, above spray zone, H.H. ALLAN (CHR 347196). - Rangitoto Island, 36° 48' 30" S, 174° 52' E, lava rock at shore, 10 January 1985, H. HERTEL 29024 (M); interior of the island, 10 January 1985, H. HERTEL 29027 (M). - Coromandel Co., Mt Maungatwhiri, near Whitianga, on rocks in grassland, August 1974, B.W. HAYWARD H44.239 (AK).

Gisborne: East Cape: Mt Hikurangi, c. 1200 m, 29 December 1982, J.K. BARTLETT 27006 (M).

Hawke's Bay: NW Ruahine, Otupai Range, western slopes, 39° 32' S, 176° 11' E,

1000–1100 m, 17 October 1983, J.K. BARTLETT 26999 (M), 27012 (M).

Wellington: York Bay (near Wellington), steep earth slope at the path between the forest and the road, 17 December 1926, G.E. & G. DU RIETZ 1277 (UPS). – North Tararu Range, Pahiatua Track, March 1970, D.J. GALLOWAY (?) (BM).

Nelson: Tasman Mts, Lonely Lake, c. 1270 m, 20 December 1983, J.K. BARTLETT 26976 (M). – Near Dunn Saddle, 41° 21' S, 173° 22' E, on ultrabasic rock, 1460 m, 12 January 1983, J.K. BARTLETT 24568 (M). – Ultramafic area near summit of Mt Dunn 1000 m, 15 July 1979, J.K. BARTLETT 27038 (M) – Upper Cobb Valley, headwaters of Burgoa Stream, c. 1180 m, 16 December 1983, J.K. BARTLETT 26972 (M). – Mt Cobb, 1585 m [undated], J.K. BARTLETT 24580 (M). – Lake Rototiti, 600 m, 14 January 1960, D. SCOTT, hb. Murray 435 (OTA).

Canterbury: Arthur's Pass National Park: Woolshed Hill, from Hawdon Shelter following the summit track, 42° 49' S, 171° 45' E, 700 m, rock outcrops in *Nothofagus solandri* var. *ciliifloroides* forest, 24 January 1983, H. HERTEL 29941 (M). – 9.3 km SSE of Arthur's Pass, Bealy Spur, along track to Grassmere Hut, 43° 02' S, 171° 33' E, on pebbles in open area, 17 December 1980, L. TIBELL 10103 (UPS). – Porters Pass, Torless Range, 970 m, on greywacke in alpine heath, 23 March 1980, J.A. ELIX 7647 (ANUC). – Bank's Peninsula, 1 km SE of Carew's Peak, 640 m, on basalt in tussock grassland, 10 June 1980, J.A. ELIX 8382 (ANUC). – Bank's Peninsula, on outcropping siliceous rocks facing SE, 1 January 1981, L. TIBELL 10173 (UPS). – Mt Peel, Lynn Stream valley, Long Speer above Acland's Hut, 43° 50' 50" S, 171° 11' E, 770 m, shaded rock face, 15 January 1983, H. HERTEL 29442 (M), 29448 (M).

Otago: W Red Hut, Herbert, 610 m, schist outcrops, 8 July 1970, P. CHILD 518 (Child). – Kidd's Bush, Lake Hawea, 300 m, 3 June 1973, P. CHILD 1803 (Child). – Flagstaff Hill near Dunedin, rock outcrop near summit, 45° 56' S, 170° 28' E, 650 m, 1 February 1983, H. HERTEL 30114 (M). – Hooper's Inlet, Otago Harbour, 18 February 1953, W. MARTIN (CHR 407364). – Summit of Mt Charles, Otago Peninsula, 400 m, 22 July 1970, P. CHILD 577 (Child). – Summit of Silver Peak (Pulpit Rock), 45° 45' S, 170° 26' 30" E, W of Waiau, horizontal schist layers near ground, 760 m, 29 January 1983, H. HERTEL 32345 (M, beside *Lecidea fuscocastanea*). – Mt Maungatua, 43° 34' S, 170° 05' E, swampy summit plateau with scattered rocks, 830–890 m, 31 January 1983, H. HERTEL 30089 (M). – Rock and Pillar Range, below University Hut, 900 m, on small quartz rock in open *Chionochloa rigidula* tussock grassland, 28 March 1980, C.D. MEURK (BM).

Southland: Dunrobin Stn., mid Aparima Valley, 247 m, 2 October 1984, C.D. MEURK ML32 (hb. Meurk). – West Dome, S slopes of East Peak, c. 600 m, 20 June 1982, C.D. MEURK ML35 (hb. Meurk).

Tasmania: King Island, Cape Wickham, 39° 34' S, 143° 37' E, 1969, J. WANGAREK (HO 63282). – King Island, 39° 33' S, 144° 00' E, 1968, J. WANGAREK, hb. BRATT no. 68/1497 (HO 63738). – Yarra Creek gorge, E. coast of King Island, 6 March 1966, J.H. WILLIS (MEL 1018329). – Mathinna, 41° 29' S, 147° 53' E, 26 October 1972, J. ADAM S, hb. BRATT no. 72/1155 (HO 65266). – Crater Lake Lookout, 41° 39' S, 145° 57' E, on quartzite, exposed and uncommon, 8 January 1972, G.C. BRATT 72/23 (HO 63730). – Coastal rocks between Shelly Point and Buxton Point, 42° 14' S, 148° 01' E, S of Swansea (N of Mayfield Bay), dolerite, 18 February 1985, H. MAYRHOFER (no. 7347) & H. HERTEL (GZU). – Orford Beach, 42° 34'S, 147° 52'E, on sandstone on exposed cliffs, very common, 17 November 1963, G.C. BRATT (no. 803) & J.A. CASHIN (HO 65323) – Orford Road between Bust-me-Gull Hill and Break-me-Neck Hill, 42° 37' S, 147° 38' E, 300 m, on dolerite in very open timber, common, 31 August 1968, G.C. BRATT (no. 68/842) & J.A. CASHIN (HO 65309). – Maria Island, 42° 38' S, 148° 05' E, on dolerite, exposed, Easter 1968, D.O. THOMAS (hb. BRATT 68/263) (HO 63759). – Mt Field National Park, track to Beatties Tarn, 42° 41' S, 146° 39' E, 920 m, 18 August 1963, G.C. BRATT & J.A. CASHIN, hb. BRATT no. 489a (HO 63713). – Black Hills Road summit, 42° 45' S, 147° 01' E, 390 m, on dolerite, 20 July 1963, G.C. BRATT (no. 386) & J.A. CASHIN (HO 63716). – Sorell-Nugent Road, Native Cherry Clump, 42° 45' S, 147° 41' E, 180 m, on dolerite, in the open, common, 15 July 1967, G.C. BRATT (no. 4114) & J.A. CASHIN (HO

CASHIN (HO 63673). - At Fossil Lake, 42° 47' S, 146° 29' E, 960 m, on rock at edge of lake, common, 7 June 1973, G.C. BRATT (HO 65291). - Grass Tree Hill, 42° 48' S, 147° 22' E, on exposed rock, 6 June 1964, G.C. & M.H. BRATT 1330 (HO 63712). - Port Arthur Road, 36.8 km from Sorell, 42° 50' S, 147° 38' E, on dolerite in the open, 7 March 1967, G.C. BRATT 3849a (HO 63698). - Lomavale, 24 km NW of Huonville, 42° 51' S, 146° 47' E, 70 m, on dolerite in the open, 28 June 1964, G.C. BRATT (no. 13544) & J.A. CASHIN (HO 63709). - Near Carbon Turn off on Port Arthur Road, 42° 52' S, 147° 41' E, 7 March 1967, G.C. BRATT & MATTHEWS, hb. BRATT 3845a (HO 63697). - Miller Track, Fern Tree, 42° 55' S, 147° 14' E, 800 m, on dolerite, 16 May 1964, G.C. BRATT (no. 1277) & J.A. CASHIN (HO 63719). - 14 km ESE of Huonville, Snug Mt, Snug Falls, 43° 05' S, 147° 12' E, on outcropping rock, 180 m, 8 March 1981, L. TIBELL 11277 (UPS, M). - Huon Road, 3 March 1891, (probably) W.A. WEYMOUTH (NSW 155807). - Mt Sprent, quartzite outcrops in *Gymnoschoenus* moorland, 800 m, 5 February 1987, O. KANTVILAS 5/87 (M). - Snug Falls Track, 43° 05' S, 147° 13' E, 360 m, 18 September 1966, G.C. BRATT (no. 3470) & M.H. BRATT (HO 63675). - Lune River Lagoon Mouth, 43° 26' S, 146° 54' E, 6 July 1968, G.C. BRATT 68/322 (HO 65308, beside *Diothallus pubescens*). - Scotts Peak Rd. near Celtic Hill, boulders in sedgeland heath, 400 m, quartzite, 11 August 1981, G. KANTVILAS & P.W. JAMES (BM). - Near Hobart(?) 1886(?) R.A. BASTOW(?) (H, ex hb. P.R.M. WILSON). - Summit of Mt Wellington, 1270 m, 5 January 1965, R. FILSON 6856 (MEL 1018537), 6860 (MEL 1031118); 1270 m, on exposed dolerite rocks, 16 January 1979, J.A. ELIX 5562 (ANUIC, beside *Poeltiaria coronandica*). - Mt Wellington, Myrtle Gully, 5 January 1965, R. FILSON 6837 (MEL 1031119), 6838 (MEL 1031119). - The Needles, outcrop in sedgeland-heath, 750 m, 5 January 1985, O. KANTVILAS 10/85 (hb. Kantvillas). - Crossing Plains, pebbles in sedgeland heath, 180 m, quartzite, 11 March 1984, G. KANTVILAS 127/84 (M). - Snug Tiers, mudstone and soil, disturbed ground in dry sclerophyll forest, 200 m, 12 October 1980, G. KANTVILAS 453/80 (hb. Kantvillas). - Prosser R., Orford, dolerite rocks in open *Eucalyptus* forest, 24 September 1981, O. KANTVILAS & A. HENSSEN 940/81 (hb. Kantvillas). - Sleepy Bay Road, 1 mile W of coastline, dry sclerophyll forest, on granite, 20 m, 2 February 1984, P. JAMES & G. KANTVILAS 160/84 (hb. Kantvillas). - Cox Bight, behind W beach, quartzite pebbles in sedgeland heath, 0 m, 5 March 1985, G. KANTVILAS 102/85 (hb. Kantvillas). - Kangaroo Point, no collector, no date (NSW 155746). - Mt Knocklofty, under dripping water, 1886, R.A. BASTOW (MEL 6977). - Mt Knocklofty, 23 August 1893, (probably) W.A. WEYMOUTH (NSW 153813). - Summit of the Acropolis, 7 January 1965, R. FILSON 6937 (MEL 1031116). - Lake St.Clair National Park, trail from Derwent River toward lakes at base of Mt Hugel, badly burnt forest, 24 February 1968, W.A. WEBER & D. McVEAN (COLO, L-49679).

All specimens contain confluentic acid together with 2'-O-methylperistolic and 2'-O-methylmicrophyllinic acid. *Paraporpidia leptocarpa* is a very common saxicolous lichen in most regions of New Zealand and Tasmania.

(31) *Poeltiaria coronandica* (Zahlbr.) Rambold & Hertel

In RAMBOLD, Biblioth. Lichenol. 34: 265 (1989) - *Lecidea coronandica* Zahlbr., Denkschr. Akad. Wiss. Wien, math.-natur. Kl. 104: 311 (1941) - HERTEL, Min. Bot. München 23: 326 (1987).

Tasmania: Lake St. Clair National Park, Cradle Mts, in the vicinity of the Labyrinth, 7 January 1965, R. FILSON 6978 (MEL 1031112). - Cradle Mountains National Park, 41° 39' S, 145° 57' E, on basalt boulders in snow beds, 1967, D. McVEAN 6753 (CBO). - Hounslow Heath, rock outcrops (schist) in alpine heath, 1080 m, 17 February 1984, P. JAMES & O. KANTVILAS 342/84 (hb. Kantvillas). - Ben Lomond National Park, 41° 32' S, 147° 30' E, on basalt pebbles in late snow area, 1967, D. McVEAN 6765 (CBO). - Mt

Rufus, ca. 1100 m, Bergwald-Region, lose Sandsteinplatten an einer Wegböschung, 20 February 1983, H. HERTEL 22903 (M). - Mt King William I, boulders in alpine heath, 1240 m, 28 January 1984, O. KANTVILAS 73/84 (hb. Kantvillas). - Summit of Mt Wellington, near Hobart, 1270 m, on exposed dolerite rocks, 16 January 1979, J.A. ELIX 5562 (ANUC, beside *Paraporpidia leptocarpa*).

(32) *Poeltiaria corrugans* (Ris.) Hertel

HERTEL, Beih. Nova Hedwigia 79: 431 (1984); HERTEL, Mitt. Bot. München 21: 310-312 (1985) et 23: 330 (1987); RAMBOLD, Biblioth. Lichenol. 34: 268 (1989).

Juan Fernandez (Chile): Masiatierra: Pelsen im obersten Teil des Villagratales, 10. Januar 1917, C. & I. SKOTTBERG s.n. (UPS, det. ZAHLEBRUCKNER 1924: 359-360 as "Lecidess leucopeltis Müll.Arg.").

New Zealand: Canterbury: Banks Peninsula, along the summit trail to Mt Sinclair, on basalt rocks in dense bush, 350 m, 14 February 1980, J.A. ELIX 7135 (ANUC).

Tasmania: Little Fisher River, on ?granite on rocks in rainforest floor, 880 m, 15 February 1984, P. JAMES & O. KANTVILAS 462/84 (hb. Kantvillas).

(33) *Poeltiaria fuscopurpurea* (Koerb.) Hertel

HERTEL, Beih. Nova Hedwigia 79: 431-432 (1984); HERTEL, Mitt. Bot. München 21: 311, 326 (1985); RAMBOLD, Biblioth. Lichenol. 34: 272 (1989).

New Zealand: Auckland: Three Kings Islands, Great Island, Castaway Valley, 12 November 1970, D. GALLOWAY (BM). - Whanauiti Beach, 35° 31' S, 173° 28' E, xanthicoleous on maritime rock, 15 March 1980, J.K. BARTLETT 23749 (hd.). - Bay of Plenty, Rurima Island, on rocks beneath Metrosideros areas, March 1984, B. & G. HAYWARD T21 (AK). - Bay of Plenty: Whale Island, camp V, on exposed rocks, no date, B. & G. HAYWARD T23 (AK).

Tasmania: Arve Loop Road, c. 27 km SW of Huonville, 43° 08' S, 146° 47' E, on quartzite, 21 December 1968, G.C. BRATT (no. 68/1461a), K.M. MACKAY et al. (HO 63743).

(34) *Poeltiaria perusta* (Nyl.) Hertel & Hafellner in Hertel

HERTEL, Beih. Nova Hedwigia 79: 463 (1984); HERTEL, Mitt. Bot. München 21: 330-331 (1987); RAMBOLD, Biblioth. Lichenol. 34: 275 (1989). - *Lecidess perusta* Nyl. in CROMBIE, Journ. Bot. (London) 12: 334 (1874).

Argentina: Tierra del Fuego: Dept. Ushuaia, steller Hang N Passo Garibaldi, gegenüber Lago Escondido, 400-500 m, Nothofagus-Wald mit großen Siliktfelsen, 10. Januar 1989, J. POELT (OZU).

New Zealand: Canterbury: Summit of Mt Peel, 54° 51' S, 171° 09' E, 1600-1650 m, on schistose rocks near ground, 16 January 1985, H. HERTEL 29514 (M).

Southland: Fiord Bot. Dist., alpine *Dianthus crassifolius* belt on the ridge S of Lake Harris Saddle, on rocks, 12 February 1927, O.E. & Greta DU RIETZ 1753:18 (UPS).

(35) *Porpidia albocaeruleans* (Wulfen) Hertel & Knoph

HERTEL & KNOPH, Beih. Nova Hedwigia 79: 433-434 (1984); HERTEL, Mitt.

itage Library, <http://www.biodiversitylibrary.org/>; www.b

Bot. München 21: 327-329 (1985) and 23: 331-332 (1987); HERTEL & KNOPH, Mitt. Bot. München 20: 467-488 (1984); RAMBOLD, Biblioth. Lichenol. 34: 278 (1989).

var. *albocrenulescens*

New Zealand: Auckland: Three Kings Islands, Great Island, Castaway Valley, 12 November 1970, D. GALLOWAY (BM).

Nelson: Kaihaka Coast, in coastal forest near Kaihaka Lakes, 40° 33' S, 172° 36' E, 12 December 1982, J.K. BARTLETT 22883 (M). - "Nova Zelandia" (no exact locality given), 17 July 1875, Ch. KNIGHT s.n. (UPS).

var. *polycarpiza* (Vain.) Rambold & Hertel

RAMBOLD & HERTEL in RAMBOLD, Biblioth. Lichenol. 34: 282 (1989) - *Le-
cidea polycarpiza* Vain., Ann. Acad. Sci. Fenn., Ser. A, 15 (6): 136-137 (1921) -
Hullia albocrenulescens var. *polycarpiza* (Vain.) Hertel, Herzogia 3: 373 (1975);
HERTEL, Khumbu Himal 6 (3): 207-210 (1977).

Papua New Guinea: Morobe District: Vicinity of Wau, along road from Wau to summit of Mt Keindu and Eddie Creek, roadside cutbanks, on rock outcrops and landslips, often with much seepage water, on a variety of exposures, 19 June 1968, W.A. WEBER & D. McVEAN (COLO L-50522).

(36) *Porpidea crustulata* (Ach.) Hertel & Knoph

HERTEL & KNOPH in HERTEL, Beih. Nova Hedwigia 79: 433-434 (1984); HER-
TEL, Mitt. Bot. München 21: 329-330 (1985) et 23: 332-333 (1987); SCHWAB,
Mitt. Bot. München 22: 380-388, 415-422 (1986); RAMBOLD, Biblioth. Liche-
nol. 34: 283 (1989).

Chile: Terr. Magallanes: Punta Arenas, Cerro Mina Rica, on stone in naked mineral-
soil (solifluction), 550 m, 4 March 1941, R. SANTESSON 6146 (S).

New Zealand: Wellington: Maungaroa vid Wellington, July 1874, S. BEROGREN
(S).

Otago: Old Man Range, summit plateau near Hyde Rock, 45° 22' S, 169° 12' E,
1670 m, stones on ground at foot of the tall tor, together with *Trapelia coarctata* and *Cer-
atodon delisei*, 3 February 1985, H. HERTEL 30329 (M).

Tasmania: Mt Knocklofty, 23 August 1893, (probably) W.A. WEYMOUTH (NSW
155801).

(37) *Porpidea stephanodes* (Stirt.) Hertel

HERTEL, Beih. Nova Hedwigia 79: 436-437 (1984); HERTEL, Mitt. Bot. Mün-
chen 23: 333 (1987) - *Lecidea stephanodes* Stirt. in CROMBIE, Journ. Linn. Soc.
Bot. London, 16: 221 (1877).

Macquarie Island: Summit of Mt Elder, on rock, 22 January 1972, R. HNATIUK
11843 (MEL 1027346), 11844 (MEL 1027342). - Mt Waite, about 45 m below the summit,
on rocks, SE aspect, 375 m, at Station 8a, 4 February 1964, Ph. ATKINSON no. RF-
5971 (MEL 1048263). - Mt Harwell, abundant in southerly aspect at Station 14, 335 m,
12 February 1964, R. FILSON 6036 (MEL 1048291). - NE corner of Lake Flynn, on rocks
with *Andreaea*, 3 February 1964, R. FILSON 5885 (MEL 1048256). - Approx. halve way

Along W side of Lake Strahan (Gratitide Lake), abundant in easterly aspect, 180 m, 4 February 1964, R. FILSON (no. 3942) & J. PHILLIPS (MEL 1041285). - Cliffs on the W side of the Island, 0.8 km S of Double Point, 3 February 1964, R. FILSON 5918 (MEL 1048282).

The above are the only localities known in addition to the type collection from Kerguelen Island. *Porpidia stephanodes* is easily recognized by its huge ascospores 40-60 x 20-26 µm (see HERTEL, 1984). No lichen substances were detected by TLC in any of the specimens. The development of an epilithic thallus varies considerably. Many specimens are cryptothalline, but some show a moderately thick dirty whitish to pale milky brownish thallus.

(38) *Protoparmelia petracoides* (Bab. & Mitt.) Hertel

HERTEL, Mitt. Bot. München 21: 330 (1987); HERTEL, Beih. Nova Hedwigia 79: 452 (1984); RAMBOLD, Biblioth. Lichenol. 34: 129 (1989).

South Africa: Cape Province: Summit plateau of Table Mt near Cape Town, 1000-1080 m, 7 June 1982, H. HERTEL 24131 (M), 25103 (M), 25130 (M). - Franschhoek Mts, Franschhoek Pass near Franschhoek (23 km E of Stellenbosch), 720-760 m, 28 May 1982, H. HERTEL 24798 (M).

New Zealand: Otago: East Cape, Mt Hikurangi, saxicolous on summit rocks, 1735 m, 27 December 1982, J.K. BARTLETT 24704 (M).

Nelson: Upper Cobb Valley, headwaters of Burgoo Stream, 1180 m, 16 December 1983, J.K. BARTLETT 26984 (M).

Canterbury: Lees Valley road, 7 km N of Oxford, 460 m, on greywacke rocks in open bush, 28 May 1980, J.A. ELIX 8304 (ANUC).

Otago: Alexandra, on schist rocks, 24 December 1963, W. MARTIN 6968 (CHR 407420). - Fruiland, 12 km S of Alexandra, 45° 20' S, 169° 19' E, 460 m, 6 November 1981, V. WIRTH (no. 14064) & C.D. MEURK (STU). - Rock and Pillar Range, 1190 m, 18 September 1981, C.D. MEURK ML17 (hb. Meurk). - Old Man Range, Obelisk near Fruiland, 45° 19' S, 169° 13' E, 1690 m, alpine *Calymine-Racoule* vegetation, 6 November 1981, V. WIRTH (no. 11866) & C.D. MEURK (STU). - Fiordland National Park, 5 km SE Homer Huts, Lyttles Flat, 44° 48' S, 168° 03' E, 660 m, 21 September 1981, H. MAYRHOFER 2262 (OZU).

Tasmania: Cradle Mts, Lake St. Clair National Park, on the western side of Mt Campbell, on rock outcrop overlooking Lake Dove, c. 45 m below summit, 1 March 1968, R. FILSON 10841 (MEL 1035723). - Lake St. Clair National Park, trail from Derwent Bridge toward lakes below Mt Hugel, badly burnt forest, 24 February 1968, W.A. WEBER & D. McVEAN (COLO L-49683, with *Lecanora cf. blanda* Nyl.). - Arm River, edge of river in wet forest, on dolerite, 440 m, 14 February 1984, P. JAMES & G. KANTVILAS 354/84 (hb. Kantvilas).

(39) *Rimularia prephotota* (Tuck.) Hertel & Rambold

HERTEL, Mitt. Bot. München 21: 334 (1987); RAMBOLD, Biblioth. Lichenol. 34: 307 (1989); HERTEL & RAMBOLD (in press) - *Lecidea prephotota* Tuck., Proceed. Acad. Arts Sci. 12: 181 (1877) - *Lambielia prephotota* (Tuck.) Hertel, Beih. Nova Hedwigia 79: 460 (1984); HERTEL, Mitt. Bot. München 21: 317-319 (1987).

Argentina: Tierra del Fuego: Dept. Ushuaia, Estancia Most, etwa 4 km W von Pampa de los Indios, *Nothofagus betuloides*-Wald, felsige Klüste, 0-30 m, 12. Januar 1989, J. POELT (OZU, 3 collections, one of it in parasitic contact with a species of *Catoplaea*).

- Montes Martial bei Ushuaia, oberhalb Acostilla, Sichtfelsen im Krummholzbereich, S- und SE-Hänge, 700-1000 m, 17. Januar 1989, J. POELT (OZU, 2 Proben).

Chile: Terr. Magallanes: Natales, Cerro Dorotea, on big boulder in a very thin forest, 9 May 1940, R. SANTESSON 2130 (S).

New Zealand: Westland: Mt Alack ridge, upper neve of Fox Glacier, 2550 m, site of Pioneer Hut, sheltered rock on S side below ridge crest, 11 January 1967, B.A. FINERAN 2403 (CANU). - Mt Haast, West Peak summit, summit rocks, 10 January 1967, B.A. FINERAN 2363 (CANU) and 2453 (CANU).

Canterbury: Phipps Peak, ridge to summit, 1830 m, near Arthur Pass, c. 42° 54' S, 171° E, 3 March 1966, B.A. FINERAN (CANU, beside *Lecidea lepicida*). - Cass, on boulders of greywacke in *Dicroidium-Festuca* tussock country above the Biological Station, 8 January 1927, G.E. & Greta DU RIETZ 1454:6 (UPS, beside *Lecidea atromorio* Knight). - Taumarunui, ca. 43° 07' S, 170° 13' E, on rock outcrop on ridge near Ball Hut, 8 April 1966, B.A. FINERAN 2329 (CANU), 2330 (CANU). - Kirkliston Range, 1670-1830 m, 25 March 1978, D.J. GALLOWAY s.n. (BM). - Lees Valley road, 7 km N of Oxford, 460 m, on greywacke rocks in open bush, 28 May 1980, J.A. ELIX 8304 (ANUC, beside *Protoparmelia petrovaeoides*). - Bank's Peninsula, Kennedy's Bush, hill with siliceous outcrops, 43° 39' S, 172° 36' E, 200 m, 1 November 1981, V. WIRTH (no. 11804) & L. GREENFIELD (STU). - Lake Pukaki Hotel, on non-calcareous boulders in *Festuca novae-zelandiae*-tussock at the road side, 12 March 1927, G.E. & G. DU RIETZ 2094b (UPS).

Otago: Rock and Pillar Range, 1190 m, 18 September 1981, C.D. MEURK ML17 (hb. Meurt). - Old Man Range, summit plateau between Hyde Rock and Obelisk, 1670 m, 3 February 1983, H. HERTEL 30344 (M, beside *Lecidella*).

South Georgia: About 10-15 m below "Summit 706 m" due E of Swinhoe Peak above Hamburg Glacier, on block scree on S-facing slope, probably never receiving direct sunshine, 6 November 1976, R.I. LEWIS SMITH 2326 (AAS, very scanty beside *Carbonea varicosa*, *Umbilicaria propagulifera* and *Ulmus sphaerocarpa*). - Royal Bay, above Köppen Point (GR 162 098), on dry, S-facing bird perching stone, 30 m, 25 January 1972, D.C. LINDSAY 4092 (AAS, beside *Tephromela strigosa*).

Antarctica: West Graham Land: Berthelot Group, Green Island, N facing inland cliff, 18 March 1935, British Graham Land ("Penots") Expedition no. 1081k (BM).

(40) *Schaereria fuscocinerea* (Nyl.) Clauzade & Roux

CLAUZADE & ROUX, Bull. Soc. Centre-Ouest, N.S., Num. Spéc. 7: 829 (1963); RAMBOLD, Biblioth. Lichenol. 34: 310 (1989). - *Lecidea fuscocinerea* Nyl., Nya Bot. Notiser: 177 (1852). - *Schaereria tenebrosa* (Plot.) Hertel & Poelt in HAWKS-WORTH, JAMES & COPPINS, Lichenologist 12: 107 (1980) - *Lecidea tenebrosa* Plot. in SCHUCHARDT, Bot. Zeitung (Berlin) 13: 146-147 (1855).

New Zealand: Canterbury: Banks Peninsula, Devil's Gap (between Lake Forsyth and Akaroa Harbour), 43° 49' 30" S, 172° 49' 30" E, on hard basaltic rock outcrop, E exposed, near summit, 670-710 m, 19 January 1983, H. HERTEL 29695 (M).

The species is widespread in the Northern Hemisphere. In addition to the above, all Southern Hemisphere records are from South Georgia (HERTEL 1977a) and from Australia (RAMBOLD 1989).

(41) *Sporastatia polypora* (Nyl.) Grunm.

GRUNMANN, Catal. Lich. Germaniae: 23 (1963); HERTEL, Mitt. Bot. München 23: 334-335 (1987).

Antarctica: East Graham Land: Trinity Peninsula, Hope Bay, Scar Hills, $63^{\circ} 25' S$, $57^{\circ} 01' W$, 90 m, 25 September 1945, I.M. LAMB, Operation Tabarin, no. D 2491 (BM, a tiny thallus within a mosaic of other crustose lichens).

(42) *Sporastatia acetuginosa* (Ach.) Massal.

MASSALONGO, Genes. Lich.: 9 (1854); MAGNUSSON, Rabenhorst: Kryptog. Fl. 9, Teil 1: 9-13 (1936); HERTEL, Mitt. Bot. München 21: 316-317 (1963) and 22: 334 (1987).

Argentina: Tierra del Fuego: Montes Martial bei Ushuaia, oberhalb Acrocailla, Silikatfelsen im Krummholzbereich, S- und SE-Hänge, 700-1000 m, 17. Januar 1989, J. POELT (GZU).

New Zealand: Westland: Mt Haast, West Peak summit, summit rocks, 10 January 1967, R.O. CUNNINGHAME (CANU, hb. PINERAN 2455).

Otago: The Remarkables, Double Cone above Lake Alta, $45^{\circ} 04' S$, $168^{\circ} 48' E$, 2280 m, exposed rocks, 5 February 1985, H. HERTEL 30443 (M, beside *Lecidea lapicida*).

South Georgia: Royal Bay, Mt Krokisius, 29 April 1902, C. SKOTTSBERG (S, det. C.W. DODGE as *Lecidea azurella* MELL. Arg.).

(43) *Stephanocyclos hessenianus* Hertel

HERTEL, Lecidaceae etc., fasc. V, no. 96 (1983); HERTEL, Beih. Nova Hedwigia 79: 438-439 (1984); HERTEL, Mitt. Bot. München 23: 335 (1987).

Kerguelen Island: (no locality given), 27 December 1897 - 20 January 1898, R. HALL (MEL 11101, loose apothecia only).

Stephanocyclos hessenianus is very common on the Prince Edward Islands (which are lying 4000 km to the West) and is also known from Western Patagonia and Tasmania (HERTEL 1987a).

(44) *Tephromela atra* (Huds.) Hafellner

in KALB, Lich. Neotrop., fasc. VII, no. 297 (1983); HERTEL & RAMBOLD, Bot. Jahrb. Syst. 107: 492-494 (1985); RAMBOLD, Biblioth. Lichenol. 34: 150 (1989) - *Lecanora atra* (Huds.) Ach., Lichenogr. Univ.: 344 (1810).

Only lecideoid/apicillloid forms with immersed apothecia (due to a heavily eroded thallus surface), found in the herbaria under *Lecidea*, are listed here.

South Georgia: Cumberland East Bay: c. 1 km W of Mt Duse (GR 133 127), 350 m, exposed dry greywacke outcrop facing W, 23 November 1971, D.C. LINDSAY 3141 (AAS, beside *Lecidea lapicida*).

Kerguelen Island: Cliffs above Lake du Val Studer, 11 February 1963, R. FILSON 4658 (MEL 1032387).

Auckland Islands: Auckland Island: Mt Eden, basalt rock of summit knob, 420 m, 11 April 1980, C.D. MEURK s.n. (BM).

Macquarie Island: W of Aurora Cave, coastal rocks, 21 January 1972, R. HNATIUK 11804 (MEL 1027290). - North Lusitania Bay, on scree, 270 m, 29 December 1971, R. HNATIUK 11592 (MEL 1027328).

itage Library, <http://www.biodiversitylibrary.org/>; www.

(45) *Trapelia coarctata* (Sm.) Choisy

CHOISY in WERNER, Bull. Soc. Sci. Nat. Maroc, J2: 160 (1932); HERTEL, Beih. Nova Hedwigia, 79: 457 (1984).

Papua New Guinea: Morobe Province: Road 4,6 km SW of Bulolo, 7° 14' S, 146° 36' E, 1000 m, large boulder in regrowth at edge of Awaracaria plantation, 16 January 1983, H. STREIMANN 33439 (BGC).

New Zealand: Auckland: Hokianga, 1874, S. BERGREN (S; already included in HELLBOM 1896: 104). - Omatangi, c. 600 m, 1874, S. BERGREN (S; already included in HELLBOM 1896: 104). - Coromandel, IX.1874, D. BERGREN (S; already included in HELLBOM 1896: 104).

Canterbury: Torlesse Range, Foggy Peak above Porter's Pass (between Cass and Springfield), 43° 18' S, 171° 45' E, ± 1250 m, sandstone pebbles partly imbedded into ground, 22 January 1983, H. HERTEL 29743 (M). - Mt Peel, Lynn Stream Valley, Long Speer above Acland's Hut, 43° 50' 30" S, 171° 11' E, 950 m, 15 January 1983, H. HERTEL 29463 (M, beside *Trapelia involuta*). - Dunedin, V.1874, S. BERGREN (S; already included in HELLBOM 1896: 104).

Otago: Old Man Range, summit plateau near Hyde Rock, 45° 22' S, 169° 12' E, 1670 m, stones on ground at foot of the tall tor, together with *Porpädia crustulata* and *Cetraria islandica*, 3 February 1983, H. HERTEL 30329 (M).

Southland: "Springfield", Otikerama, 215 m, on rock bluff, 27 February 1984, C.D. MEURK ML7 (hb. Meurk). - Street Manspout - Clifden, c. 3 km N of Blackmount Pass, 45° 44' S, 167° 37' E, pebbles loose on ground of a road cutting, 11 February 1983, H. HERTEL 30752 (M).

Tasmania: Moonah, 67 Sinclair Avenue, 42° 51' S, 147° 18' E, on dolerite, exposed and rare, 27 July 1968, O.C. BRATT 68/572 (HO 63807). - Murchison Highway, 1.6 km N of Que River, 41° 33' S, 145° 44' E, 7 April 1969, O.C. BRATT 69/261 (HO 63746). - "Tasmania" (without exact locality), FITZGERALD (NSW 153800).

(46) *Trapelia involuta* (Th. Teyl. in Mackay) Hertel

HERTEL, Herzogia 2: 508-513 (1973); HERTEL, Khumbu Himal 6 (3): 346-347 (1977). - *Lecanora involuta* Th. Teyl in MACKAY, Flora Hibernica : 134-135 (1836).

New Zealand: Auckland: Thames Co., Slipper Island, on clay bank, July 1973, B.W. HAYWARD H49.145 (AK).

Canterbury: Mt Peel, Lynn Stream Valley, Long Speer above Acland's Hut, 43° 50' 30" S, 171° 11' E, 950 m, 15 January 1983, H. HERTEL 29463 (M, beside *Trapelia coarctata*).

This specimen has a tiny, clearly lobulate thallus containing gyrophoric acid. *Trapelia involuta* is well known from Europe and North America (HERTEL 1973) and has been reported also from East Africa (HERTEL 1973) and Indonesia (HERTEL 1977a). It is a new record for Australasia.

(47) *Trapelia moortiana* (Carroll) P. James

P. JAMES in HERTEL, Herzogia 3: 405-406 (1975); HERTEL, Khumbu Himal 6 (3): 347-349 (1977); HERTEL, Mitt. Bot. München 27: 335-337 (1987) - *Lecidea moortiana* Carroll, Nat. Hist. Rev. 6: 529 (1859).

Tasmania: C. 2 km NW of Trial Harbour, 41° 55' S, 145° 11' E, on roadside gravel and pebbles in buttongrass moorland, 200 m, 6 April 1989, G. KANTVILAS 143/89 (M, dupl. ex HO).

This is a second record of this widely distributed species (see distribution map in HERTEL 1987b p. 337) for Australasia.

(48) *Tremolecia strata* (Ach.) Hertel

HERTEL, Khumba Himal 6 (3): 351-353 (1977); HERTEL, Beih. Nova Hedwigiae 79: 458 (1984); HERTEL, Mitt. Bot. München 21: 335-336 (1985) and 22: 337 (1987); SCHWAB, Mitt. Bot. München 22: 441-453 (1986); RAMBOLD, Biblioth. Lichenol. 34: 316 (1989) - *Lecidex strata* (Ach.) Wahlb., Fl. Lappon.: 475 (1812); HERTEL, Herzogia 2: 483 (1973).

Colombia: Depto. de Valle: Parque Nacional de Cali (Pacific Range of the Andes on the inland side), 3° 20' N, 76° 42' W, 3750 m, montane-subalpine rain forest, on exposed summit of mountain, 24 January 1976, D. HARTMAN 336 (COLO L-83469).

Argentina: Tierra del Fuego: Dept. Ushuaia, Montes Martial bei Ushuaia, oberhalb Acroilla, alpine Heiden, Silikatfelsen im Krummholzbereich, S- und SE-Hänge, 700-1000 m, 17. Januar 1989, J. POELT (GZU).

Uganda: Distr. Toro: Central summit group of Mt Ruwenzori, 4480 m, on rocks, April 1984, K. SCHMITT (Id.).

Tasmania: Mt Wellington, on dolerite, alpine rocks, 1260 m, 10 February 1984, P. JAMES & G. KANTVILAS 301/84 (hb. Kantvillas). - Cradle Mountains, Lake St. Clair National Park, Mt Rufus, 42° 07' S, 146° 06' E, W of Derwent Bridge, c. 1320 m, an Außenbastenfläche eines großen, harten, freistehenden Feisboches, 20. Februar 1985, H. HERTEL 32914 (M, sub *Rimularia prephota*).

South Georgia: About 10-15 m below "Summit 706 m" due E of Swinhoe Peak above Hamburg Glacier, on block scree on S-facing slope, probably never receiving direct sunshine, 6 November 1976, R.L. LEWIS SMITH 2523 (AAS). - Cumberland East Bay: Hope Point (GR 133 124), on moderately sheltered dry, non-nitrogenous greywacke outcrop facing S, 50 m, 24 November 1971, D.C. LINDSAY 3187 (AAS, beside *Lecidex lippicida*). - Royal Bay: near waterfall on SW side of Whale Valley (GR 158 098), on dry non-nitrogenous boulder, 30 m, 17 January 1972, D.C. LINDSAY 3945 (AAS, beside *Lecidex* sp.).

South Shetland Islands: King George Island: N side of Barton Peninsula, 62° 13' S, 58° 44' W, 3 m, on pebbles of raised beach, 22 January 1986, D.C. LINDSAY 736 (AAS).

Antarctica: East Graham Land: Trinity Peninsula, Hope Bay, N flank of Mt Flora, 300 m, 6 January 1949, I.M. LAMB, Operation Tabarin, no. 2834 (BM). - Summit ridge of Mt Flora, 520 m, 3 January 1945, D. JAMES & V. RUSSELL, Operation Tabarin, no. 2859 (BM). - Hope Bay, low hill on E. side of bay, 100 m, on loose stone, 16 April 1945, I.M. LAMB, Operation Tabarin, no. 2376 (BM), no. 2389 (BM). - Hope Bay, from a moraine between two ridges N of Mt Bransfield, 180 m, 10 October 1946, A. REECE, Falkland Island Dependencies Survey no. D-355-5 (BM). - Nobby Nunatak, south side of Hope Bay, 270 m, small boulder in shelter of larger ones, 15 January 1960, C.O. BRADING 216 (AAS).

West Graham Land: Argentine Islands: Galindez Island, 65° 15' S, 64° 15' W, from typical boulder area, just uncovered by snow melt, 27 December 1935, British Graham Land ("Pencos") Expedition 1934-37, no. 1328r (BM).

(49) *Trichothallia pahensis* (Zahlbr.) Hertel & Killas

HERTEL, Mitt. Bot. München 23: 337-338 (1987); RAMBOLD, Biblioth. Lichenol.

itage Library, <http://www.biodiversitylibrary.org/>; www.biodiversitylibrary.org/page/155.html
34: 155 (1989).

New Zealand: North Auckland: 17.5 km ENE of Warkworth, Tawharanui Peninsula, 3.5 km W of Tokatu Pt., $36^{\circ} 22' S$, $174^{\circ} 50' E$, on sea-shore rocks, 25 May 1981, L. TIBELL 13191 (UPS, beside *Lecanora subcoarctata*).

Southland: Southern shore of Bluff Peninsula, boulders in the lowest aerohyaline belt, 26 March 1977, Greta DU RIETZ 3007:6 (UPS), 3007:4 (UPS). - Howells Point near Riverton, $46^{\circ} 23' S$, $168^{\circ} 02' E$, coastal rocks, 11 February 1983, H. MAYRHOFER (no. 7269) & H. HERTEL (OZU).

Tasmania: Flinders Island, 4 km W of Emu, 17 April 1965, R. FILSON 7109 (MEL 564158). - Lune River Lagoon Mouth, $43^{\circ} 26' S$, $146^{\circ} 54' E$, 6 July 1968, G.C. BRATT 68/322 (HO 65308, beside *Paraporphidium leptoocarpum*). - Tasman Peninsula, coastal heath between Safety Cove and Crescent Bay near Standup Point, S of Port Arthur, $43^{\circ} 12' S$, $147^{\circ} 52' E$, c. 5 m, 24 October 1981, H. MAYRHOFER 3353 (OZU). - Blackmans Bay near Kingston, S of Hobart, $43^{\circ} 00' 40'' S$, $147^{\circ} 19' 30'' E$, Flowerpot Point SE of Blackmans Beach, coastal rocks, dolerite, 19 February 1983, H. MAYRHOFER (no. 7297) & H. HERTEL (OZU, Mayrhofer), 7310 (OZU), 7312 (OZU). - Coastal rocks between Shelly Point and Buxton Point, S of Swansea, $42^{\circ} 14' S$, $148^{\circ} 01' E$, 18 February 1983, H. MAYRHOFER (no. 7325) & H. HERTEL (Mayrhofer).

(50) *Xenolecia spadicommus* (Nyl. in Cramb.) Hertel

HERTEL, Beih. Nova Hedwigia 79: 440 (1984). - *Lecidea spadicommus* Nyl. in Crambie, Journ. Linn. Soc. London, Bot. 15: 233 (1876).

Argentina: Prov. Chubut: Lago Menendez, lado oeste, en piedras en la orilla del lago, por cierto sumergidas en épocas de mucha agua, 6 February 1950, I.M. LAMB 5948 (UPS).

This is a second record of this species, described from Isla Wellington in southern Chile.

LITERATURE

- CABRERA, A.J. 1971: Flora de la Repùblica Argentina. - Bol. Soc. Argent. Bot. 14: 1-42.
GALLOWAY, D.J. 1985: Flora of New Zealand Lichens. - Wellington.
HELLBOM, P.J. 1896: Lichenae Neo-Zelandicae seu Lichenae Novae Zealandiae a Mr. Berggren annis 1874-75 collecti. - Bihang K. Svenska Vet.-Akad. Handl. 21, Afld. III, no. 13.
HERTEL, H. 1973: Beiträge zur Kenntnis der Flechtenfamilie Lecideaceae V. - Herzogia 2: 479-515.
HERTEL, H. 1975: Beiträge zur Kenntnis der Flechtenfamilie Lecideaceae VI [*-Lecideaceae V* - Iapsu]. - Herzogia 3: 365-406.
HERTEL, H. 1977a: Osteinsbewohnende Arten der Sammelgattung *Lecidea* (Lichenes) aus Zentral-, Ost- und Südosten. Eine erste Übersicht. - Khunshu Himal, Ergebn. Forschungsunternehmens Nepal-Himalaya 6 (3): 143-378.
HERTEL, H. 1977b: *Lecidea* in der Arktis I. - Mitt. Bot. München 19: 337-352.
HERTEL, H. 1977c: Bemerkenswerte Flechtenfunde aus dem Gebiet des Kongofjordes und des Isfjordes (Spitzbergen). - Herzogia 4: 367-401.
HERTEL, H. 1984: Über siccicole, lecideoidae Flechten der Subantarktis. - Beih. Nova Hedwigia 79: 399-499 [Festschrift J. POELT].

- Heritage Library, <http://www.biodiversitylibrary.org/>; www.b
- HERTEL, H. 1983: New, or little-known New Zealand lecideoid lichens. - Mitt. Bot. München 21: 301-337.
- HERTEL, H. 1987a: Progress and problems in taxonomy of Antarctic saxicolous lecideoid Lichens. - Biblioth. Lichenol. 25: 219-242.
- HERTEL, H. 1987b: Bemerkenswerte Funde südhemisphärischer, saxicoler Arten der Sammelgattung *Lecidea*. - Mitt. Bot. München 23: 321-340.
- HERTEL, H. 1989: Problems in Monographing Antarctic crustose lichens. - Polarforschung 58: 65-76.
- HERTEL, H. & KNOPH, J.-G. 1984: *Porpidia albocerulescens*: eine weit verbreitete, doch in Europa seltene und vielfach verkannte Krustenflechte. - Mitt. Bot. München 20: 467-488.
- HERTEL, H. & RAMBOLD, G. 1990: Zur Kenntnis der Familie Rimulariaceae (Lecanorales). - Biblioth. Lichenol. (in press).
- LEUCKERT, Ch. & POELT, J. 1978: Über Nord-Süd-Gradienten von Chemotypen europäischer Flechten. - Plant. Syst. Evol. 130: 53-77.
- RAMBOLD, G. 1989: A monograph of the saxicolous lecideoid lichens of Australia (excl. Tasmania). - Biblioth. Lichenol. 34: 1-345.
- SCHWAB, A. 1986: Rostfarbene Arten der Sammelgattung *Lecides* (Lecanorales). Revision der Arten Mittel- und Nordeuropas. - Mitt. Bot. München 22: 221-476.
- SIPMAN, H.J.M. 1983: A monograph of the lichen family Megalosporaceae. - Biblioth. Lichenol. 18.
- WIRTH, V. 1987: Die Flechten Baden-Württembergs: Verbreitungsatlas. - Stuttgart (Hohenheim).

ZOBODAT -

www.zobodat.at

Zoologisch-Botanische
Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Mitteilungen der
Botanischen Staatssammlung München

Jahr/Year: 1989

Band/Volume: 28

Autor(en)/Author(s): Hertel Hannes

Artikel/Article: New records of lecideoid lichens from the Southern Hemisphere 211-238