

## Some new species of *Astragalus* L. sect. *Malacothrix* BUNGE in Iran

by

A.A. MAASSOUMI

### Summary:

5 new species of *Astragalus* L. sect. *Malacothrix* Bunge from Iran are described.

### Introduction:

During my revisional work on Iranian Astragali with basifixied hairs (BUNGE 1868/1869) since four years the section *Malacothrix* Bunge with a higher evolutionary drift was the focal object of my main work. The similarity of most taxa in vegetative characters and the great number of species have led to confusion also between the old and distinct taxa and between them and some recent described species. This confusion was augmented by the fact that most of the older species never were typified and the lack of adequate type material. In 1986 and 1990 I has succeeded to examin the type specimen of most of the older species borrowed by Dr. PODLECH from several European herbaria. Typification of some old species has been done by Dr. PODLECH and the author. This work is also based on the rich material of two big Iranian herbaria (TARI & EVIN) and several duplicate material of other provincial herbaria.

### *Astragalus expectatus* Maassoumi, spec. nov.

diffrerit ab *A. inexpectato* Maassoumi & Podl. foliolis 7-8 jugis (nec 10-18 jugis), bracteis anguste triangularibus dense nigropilososis (nec linearibus albo-nigropilosis), ab *A. plagiophacos* Maassoumi & Podl. calyce multo longiore et legumine lateraliter compresso (nec inflato).

Plantae perennes, acaules, pilis 0,3-0,8 mm longis obtectae. Caudex ramosus, caudiculos ad 20 cm longos emitens. Stipulae 4-6 mm longae, anguste triangulares, laxe pilosae vel inferiores subglabrate, basi breviter petiolo adnatae, inter se breviter connatae. Folia 2-3,5 cm longa, petiolo 0,5-0,7 cm longo sicut rachis pilis albis patentibus obtecto. Foliola 7-8 juga, contigua, 5-10 mm longa et 3-4 mm lata, anguste elliptica antice acutiuscula, utrinque dense semipatenter ad patenter pilosa, saepissime plicata. Inflorescentiae solitaires, pedunculo ad 6 cm longo basi pilis albis apicem versus albis nigrisque patentibus obtecto suffultae, dense capitatae multiflorae. Bractae anguste triangulares, acutae, 6-7 mm longae, dense nigropilosae. Flores subsessiles. Calyx ca. 12 mm longus, tubulosus, dense albopilosus, dentibus subulatis 4-5 mm longis albo-nigropilosis. Corolla glabra, in vivo flava, in sicco violacea. Vexillum ca. 15 mm longum, lamina obovata. Alae carinaque deficientes. Ovarium sessile, pilosum. Legumen ovatum, 10-11 mm longum et ca. 5 mm altum, ventro appланato-carinatum, dorso rotundatum et leviter sulcatum, apice in rostro curvato attenuatum, dense pilis brevissimis et insuper pilis mollibus patentibus 2,5-4 mm longis obtectum, complete biloculare.

Holotypus: Iran, Prov. Fars: 20 km from Nurabad to Kora Kan, Doshman-Ziary region (WP2), 1800 m., 31.5.1981, Mozaffarian 45716 (TARI).

*Astragalus ghoortapacensis* Maassoumi, spec. nov.

differt ab *A. iranico* Bunge indumento in rachide pendunculoque e pilis appressis (nec patentibus) consistente, ab *A. molle* M.B. foliolis minoribus et indumento in partibus vegetativis e pilis appressis consitente.

Plantae perennes, acaules, 12-14 cm altae. Stipulae 2-4 (-5) mm longae, triangulares, appresse albo-nigro-pilosae, basi petiolo breviter adnatae, inter se breviter connatae. Folia 6-8 cm longa, petiolo 1,5-2 cm longo, sicut rachis pilis brevissimis appressis vel subappressis albis vel apicem versus nigris etiam obtecto. Foliola 13-16 juga, ovata vel oblonga, apice rotundata vel late emarginata, utrinque dense subappresse pilosa. Inflorescentiae pedunculo 6-8 cm longo, dense pilis appressis, 0,2-0,3 (-0,5) mm longis albis vel apicem versus imprimis nigris obtecto suffultae, ovoidese vel cylindricae, multiflorae. Bractae lineares, albo-membranaceae, 3-4 (-5) mm longae, albo-nigropilosae. Calyx 10-12 mm longus, tubulosus, longitudinaliter purpureo-nervosus, laxe pilis ad 1 mm longis albis nigrisque patentibus obtectus. dentibus subulatis 3-4 mm longis. Corolla glabra, flava. Vexillum ignotum. Alae 19-20 mm longae, lamina anguste

obovata, apice versus dilatata, antice oblique incisa, ca. 7 mm longa et 2,5-3 mm lata, auricula ca. 1 mm longa provisa, ungue ca. 12 mm longo. Carina ca. 16 mm longa, lamina oblique elliptica, curvata, ca. 6 mm longa et 3 mm lata, ungue ca. 10 mm longo. Ovarium breviter stipitatum, pilosum. Legumen stipite ca. 1 mm longo suffultum, ovoides vel ellipticum, ca. 15 mm longum, 4-5 mm altum et ca. 3 mm latum, ventro rotundatum et leviter carinatum, dorso anguste sulcatum, apice in rostro breve curvato attenuatum, pilis brevissimis et longis lanatis obtectum, complete biloculare.

Holotypus: Iran, Prov. Esfahan, Semiroom, Ghoortapacy, 14.6.1986, M. Nowrooz 4307 (TARI).

*Astragalus khajiboulaghensis* Maassoumi, spec. nov.

differit ab *A. molle* MB. foliolis supra subglabris (nec patenter pilosis), stipulis 3-4 mm tantum longis basi petiolo breviter adnatis (nec per 4-5 mm petiolo adnatis).

Plantae perennes, acaules, 5-9 cm altae. Caudex valide ramosus. Stipulae viridiussculae, 3-4 mm longae, pilosae, basi per 2 mm petiolo adnatae. Folia 3-5 cm longa, petiolo 1,5-2 cm longo, sicut rachis dense pilis albis antrorsis 0,2-1 mm longis obtecto. Foliola 11-13 juga, 3-4 mm longa et 1-2 mm lata, anguste elliptica, antice rotundata, supra glabra vel marginem versus sparse vel laxe appressae pilosa, subtus dense pilis subappressis ad 0,5 mm longis obtecta. Inflorescentiae pedunculo 3-3,5 cm longo dense subappressae albopiloso spicem versus etiam pilis nigris immixtis suffultae, multiflorae. Bracteae 8-9 mm longae, anguste triangulares, albo-nigropilosae. Flores subsessiles. Calyx 15-16 mm longus, tubulosus, apice oblique truncatus, densiuscule pilis ad 3 mm longis albis vel spicem versus nigris immixtis obtectus, dentibus subulatis 6-7 mm longis. Corolla glabra, in vivo flava. Vexillum 19-20 mm longum, lamina ca. 7 mm lata, subrhomboidea, antice incisa. Alae ca. 16 mm longa, lamina anguste oblonga, antice rotundata, ca. 6 mm longa et 2 mm lata, auricula ca. 1,5 mm longa provisa, ungue ca. 10 mm longo. Carina ca. 15 mm longa, lamina oblique elliptica, ca. 6 mm longa et 3 mm lata. Ovarium dense pilosum. Legumen brevissime stipitatum, ovatum, ca. 10 mm longum et 4-5 mm altum, ventro fere rotundatum dorso profunde sulcatum, antice in rostro breve attenuatum, dense pilis ca. 4 mm longis obtectum, complete biloculare.

Holotypus: Iran, Prov. Azerbayejan: 60 km from Namin to Khaje-Boulagh, above Khaje-Boulagh, 1950 m, 21.6.1980, Mozaffarian & Nowrozi 34683 (TARI).

*Astragalus nahavandicus* Maassoumi, spec. nov.

differt ab *A. sarac* Eig foliolis anguste ellipticis vel linearibus (nec oblongis) supra glabris (nec pilosis), lamina alarum antice rotundata(nec oblique emarginata).

Plantae perennes, acaules, 11-13 cm altæ. Caudex ramosus, caudiculos elongatos emittens. Stipulae 4-5 mm longae acuminatae, dense pilis albis vel interdum nigris immixtis obtectae, basi ad 3 mm petiolo adnatae inter se breviter connatae. Folia 8-11 cm longa, petiolo 0,5-2 cm longo sicut rachis densiuscule pilis subappressis vel semipatentibus 0,3-0,5 mm longis et insuper laxe pilis ad 1 mm longis patentibus albis obtecto. Foliola 15-17 juga, 6-10 mm longa et 2-3 (-4) mm lata, anguste elliptica vel linearia, supra glabra vel marginem versus laxe pilosa, subtus laxe vel densiuscule pilis brevibus subappressis vel semipatentibus obtecta. Inflorescentiae pedunculo 5-6 cm longo pilis subappressis brevibus et ad 2 mm longis patentibus albis nigrisque obtecto suffultae, multiflorae elongatae. Bracteæ anguste triangulares, 5-6 mm longæ, albo-nigropilosas. Pedicelli brevissimi. Calx ca. 11 mm longus, tubulosus, basin versus leviter ventricosus pilis brevissimis appressis imprimis nigris et insuper ad 1 mm longis patentibus imprimis albis obtectus, dentibus subulatis ca. 1 mm longis. Corolla glabra, purpurea vel violacea. Vexillum 16-17 mm longum, lamina 8-9 mm lata, rhomboidea antice incisa. Alæ 15-16 mm longæ, lamina anguste oblonga, antice rotundata, ca. 7 mm longa et 2 mm lata, auricula ca. 1,5 mm longa provisa, ungue 7-8 mm longo. Carina 13-14 mm longa, lamina obliquè elliptica, ca. 6 mm longa. Ovarium sessile. pilosum. Legumen ignotum.

Holotypus: Iran, Prov. Hamadan, ca. 21 km from Nahavand on the road to Malayer, 2000 m, 11.5.1987, Maassoumi & Mirhosseini 59365 (TARI, Herb. PODLECH).

*Astragalus rusticus* Maassoumi, spec. nov.

differt ab *A. macrouro* Fischer & C.A.Meyer cui similis corolla violacea (nec flava) et vexillo rhomboidei (nec distincte ligulato).

Plantae perennes, acaules, 10-11 cm altæ. Stipulae 5-8 mm longæ, anguste triangulares, albopilosæ, basi per 2 mm petiolo adnatae, inter se breviter connatae. Folia 5-6 cm longa, petiolo 1-1,5 cm longo, sicut rachis dense pilis brevissimis et insuper pilis longis subappressis obtecto. Foliola 11-12 juga, 5-7 mm longa et 2-4 mm lata,

ritage Library, http://www.biodiversitylibrary.org/; www.b

elliptica. acuta vel obtusiuscula, utrinque appresse vel subappresse pilosa. Inflorescentiae solitaires, pedunculo 5-6 cm longo sicut rachis piloso suffultae, 7-9 florate. Bracteae anguste triangulares, 5-6 mm longae, albo-nigropilosae. Flores subsessiles. Calyx 11-12 mm longus, brevissime nigro- et longe albopilosae, dentibus subulatis, 3-3,5 mm longis. Corolla intense violacea. Vexillum ca. 10 mm longum, rhomboideum, antice emarginatum. Alae ca. 17 mm longae, lamina anguste oblonga, apicem versus leviter dilatata, rotundata, breviter auriculata, unque lamina sequante. Carina ca. 15 mm longa, lamina obliqua elliptica. Ovarium stipitatum, pilosum. Legumen ignotum.

Holotypus: Iran, Azerbayjan, Kandewan road to Esfahan, before Kahnou, 2090 m, 27.5.1987, Olfat & Sarkarat 233 (TARI).

# **ZOBODAT -**

## **www.zobodat.at**

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Mitteilungen der Botanischen Staatssammlung München

Jahr/Year: 1990

Band/Volume: [29](#)

Autor(en)/Author(s): Maassoumi Ali Asghar

Artikel/Article: [Some new species of  
Astragalus L. sect. Malacothrii BUNGE in Iran  
503-507](#)