

New Records of Tiger Beetle Species from the Malaysian Peninsular (II)¹⁾ (Coleoptera: Cicindelidae)

Jürgen WIESNER

Abstract: *Callytron doriai* (Horn, 1897) and *Enantiola hewittii* (Horn, 1908) are recorded for the first time from the Malaysian Peninsular (= Malacca). The number of tiger beetle species known from Malacca raised up to 71.

Key words: Cicindelidae, faunistic records, Malaysia, Malacca

The entomologists Uno ROOSILEHT and Aare LINDT from the Eesti Loodusmuuseum, Tallinn, Estonia, recently made a journey to the Malaysian Peninsular (Malacca). The tiger beetles, which were collected during that stay are listed in the following. All material is conserved in the Loodusmuuseum, some duplicates are deposited in the authors's collection.

Results

Tricondyla (Tricondyla) annulicornis Schmidt-Goebel, 1846

Specimens examined: 19.4.2006, Malaysia, Brinchang, 600 m, 4°32' 50''N, 101°17'54''E, leg. Uno Roosileht, 1 ♂; 24.4.2006, Malaysia, Brinchang, 600 m, 4°22'09''N, 101°20'00''E, leg. Aare Lindt, 1 ♂; 26.4.2006, Malaysia, Cameron Highlands, 600 m, 4°22'09''N, 101°20' 00''E, leg. Uno ROOSILEHT, 1 ♀.

Distribution: Myanmar, Malaysian Peninsular, Thailand, Laos, Vietnam.

¹⁾ 99th Contribution towards the knowledge of Cicindelidae.

***Neocollyris (Neocollyris) bonellii bonellii* (Guérin-Méneville, 1834)**

Specimens examined: 28.3.2006, Malaysia, Cameron Highlands, 600 m, 4°22'09''N, 101°20'00''E, leg. Uno ROOSILEHT, 2 ♂♂; 24.4.2006, Malaysia, Cameron Highlands, 600 m, 4°22'09''N, 101°20'00''E, leg. Uno ROOSILEHT, 1 ♂, 1 ♀.

Distribution: Jawa, Bali, Sumatra, Sumbawa, Sumba, Flores, Borneo (Kalimantan), Sulawesi, Malaysian Peninsular, Singapore, Nepal, India, Bangladesh, Myanmar, Thailand, Laos, Cambodia, Vietnam, China (Fujian, Guangdong, Guangxi, Hainan, Hong Kong, Hunan, Yunnan, Zhejiang).

***Neocollyris (Neocollyris) elongata* (Chaudoir, 1864)**

Specimens examined: 27.3.2006, Malaysia, Cameron Highlands, 600 m, 4°22'09''N, 101°20'00''E, leg. Uno ROOSILEHT, 1 ♀; 25.4.2006, Malaysia, Cameron Highlands, 600 m, 4°22'09''N, 101°20'00''E, leg. Aare LINDT, 1 ♂; 26.4.2006, Malaysia, Cameron Highlands, 600 m, 4°22'09''N, 101°20'00''E, leg. Uno ROOSILEHT, 1 ♀.

Distribution: Thailand, Malaysian Peninsular.

***Neocollyris (Neocollyris) chloroptera* (Chaudoir, 1860)**

Specimen examined: 24.8.2006, Malaysia, Pulau Tioman, 2°48'13''N, 104°09'31''E, leg. Uno ROOSILEHT, 1 ♂.

Distribution: Thailand, Malaysian Peninsular, ?Borneo, Sumatra (S. barat, Bengkulu).

***Neocollyris (Leptocollyris) subtilis subtilis* (Chaudoir, 1863)**

Specimens examined: 28.3.2006, Malaysia, Cameron Highlands, 600 m, 4°22'09''N, 101°20'00''E, leg. Uno ROOSILEHT, 1 ♂; 26.4.2006, Malaysia, Cameron Highlands, 600 m, 4°22'09''N, 101°20'00''E, leg. Uno ROOSILEHT, 1 ♂.

Distribution: Thailand, Malaysian Peninsular, Sumatra (S. utara, S. barat), Laos.

***Neocollyris (Stenocollyris) leucodactyla* (Chaudoir, 1860)**

Specimen examined: 24.4.2006, Malaysia, Cameron Highlands, 600 m, 4°22'09''N, 101°20'00''E, leg. Uno ROOSILEHT, 1 ♀.

Distribution: Malaysian Peninsular, Singapore, Sumatra (Aceh, S. utara, S. barat, Bengkulu, Lampung), Borneo (Brunei, Sarawak).

Neocollyris (Stenocollyris) oblita Naviaux, 1995

Specimen examined: 27.3.2006, Malaysia, Cameron Highlands, 600 m, 4°22'09''N, 101°20'00''E, leg. Uno ROOSILEHT, 1 ♀.

Distribution: Malaysian Peninsular, Sumatra, Borneo (Sarawak, Kalimantan barat).

Heptodonta analis analis (Fabricius, 1801)

Specimens examined: 25.3.2006, Malaysia, Cameron Highlands, 1500 m, 4°29'01''N, 101°22'95''E, leg. Uno ROOSILEHT, 1 ♂, 1 ♀; 3.4.2006, Malaysia, Jeli-Grik Road, 500 m, 5°38'18''N, 101°42'43''E, leg. Uno ROOSILEHT, 1 ♀.

Distribution: S Thailand, Malaysian Peninsular, Sumatra (S. utara, S. barat), Jawa, Borneo (Sarawak, Sabah), Sulawesi (S. selatan), Luzon.

Therates merkli Wiesner, 1996

Specimen examined: 9.4.2006, Malaysia, Pulau Tioman, 2°48'13''N, 104°09'31''E, leg. Uno ROOSILEHT, 1 ♂.

Distribution: Malaysian Peninsular.

Therates dimidiatus wallacei Thomson, 1857

Specimen examined: 24.4.2006, Malaysia, Cameron Highlands, 600 m, 4°22'09''N, 101°20'00''E, leg. Uno ROOSILEHT, 1 ♀.

Distribution: Thailand, Borneo, Singapore, Malaysian Peninsular, Sumatra.

Calomera angulata angulata (Fabricius, 1798)

Specimens examined: 11.4.2006, Malaysia, Kuantan, 3°16'44''N, 103°26'01''E, leg. Uno ROOSILEHT, 1 ♂; 24.8.2006, Malaysia, Pulau Tioman, 2°48'13''N, 104°09'31''E, leg. Uno ROOSILEHT, 1 ♂, 1 ♀.

Distribution: Afghanistan, Pakistan, Nepal, Sikkim, India, Myanmar, Sri Lanka, Thailand, Laos, Vietnam, Peninsula Malaysia, Sumatra, Sumbawa, Borneo, Philippines (Luzon), Taiwan, China (Anhui, Fujian, Guangdong, Guizhou, Hainan, Hebei, Henan, Hubei, Jiangsu, Jiangxi, Shaanxi, Shanxi, Sichuan, Yunnan, Zhejiang).

***Cosmodela aurulenta aurulenta* (Fabricius, 1801)**

Specimens examined: 3.4.2006, Malaysia, Jeli-Grik Road, 500 m, 5°38'18''N, 101°42'43''E, leg. Uno ROOSILEHT, 1 ♀; 4.4.2006, Malaysia, Jeli-Grik Road, 500 m, 5°38'18''N, 101°42'43''E, leg. Uno ROOSILEHT, 1 ♀. Distribution: Malaysian Peninsular, Borneo (Brunei, Sarawak, Sabah), Sumatra (Aceh, S. utara, S. barat, Bengkulu, S. selatan), Jawa, Bali, Sulawesi (S. utara). ?China (Fujian, Guangxi, Hainan, Jiangxi).

***Lophyra (Lophyra) fuliginosa* (Dejean, 1826)**

Specimens examined: 9.4.2006, Malaysia, Pulau Tioman, 2°48'13''N, 104°09'31''E, leg. Uno ROOSILEHT, 1 ♀; 19.4.2006, Malaysia, Brinchang, 600 m, 4°32'50''N, 101°17'54''E, leg. Uno ROOSILEHT, 1 ♀. Distribution: Sri Lanka, Myanmar, Peninsula Malaysia, Borneo, Bali, Jawa, Sumatra (S. utara, S. selatan), Thailand, Laos, Cambodia, Vietnam, China (Guangdong, Shanghai, Zhejiang).

***Cylindera (Verticina) versicolor* (MacLeay, 1825)**

Specimens examined: 29.3.2006, Malaysia, Cameron Highlands, 1800 m, 4°31'37''N, 101°23'26''E, leg. Uno ROOSILEHT, 1 ♂; 4.4.2006, Malaysia, Jeli-Grik Road, 500 m, 5°31'55''N, 101°15'07''E, leg. Uno ROOSILEHT, 1 ♀; 16.4.2006, Malaysia, Jeli-Grik Road, 500 m, 5°31'55''N, 101°15'07''E, leg. Uno ROOSILEHT, 1 ♀. Distribution: Thailand, Singapore, Malaysian Peninsular, Borneo (Sarawak, Sabah, Brunei), Sumatra (Aceh, S. utara, S. barat, Bengkulu, S. selatan, Lampung), Jawa.

***Cylindera (Leptinomera) catoptroides* (Horn, 1892)**

Specimens examined: 4.4.2006, Malaysia, Jeli-Grik Road, 500 m, 5°31'55''N, 101°15'07''E, leg. Uno ROOSILEHT, 3 ♀♀; 15.4.2006, Malaysia, Jeli-Grik Road, 5°38'18''N, 101°42'43''E, leg. Uno ROOSILEHT, 1 ♂, 2 ♀♀. Distribution: Malaysian Peninsular, Borneo (Sabah, Sarawak, Brunei), Sumatra (Aceh, S. utara, S. barat, Bengkulu, S. selatan, Lampung).

***Cylindera (Ifusina) viduata* (Fabricius, 1801)**

Specimens examined: 31.3.2006, Malaysia, Gua Musang, 4°44'38''N, 104°42'59''E, leg. Uno ROOSILEHT, 4 ♂♂, 4 ♀♀; 31.3.2006, Malaysia,

Kelantan Gua Musang, 228 m, 04°44'38''N, 101°42'59''E, leg. Aare LINDT, 1 ♂; 3.4.2006, Malaysia, Jeli-Grik Road, 500 m, 5°38'18''N, 101°42'43''E, leg. Uno ROOSILEHT, 2 ♂♂; 4.4.2006, Malaysia, Jeli-Grik Road, 500 m, 5°31'55''N, 101°15'07''E, leg. Uno ROOSILEHT, 1 ♀. 16.4.2006, Malaysia, Jeli-Grik Road, 500 m, 5°31'55''N, 101°15'07''E, leg. Uno ROOSILEHT, 1 ♀; 17.4.2006, Malaysia, Jeli-Grik Road, 500 m, 5°31'55''N, 101°15'07''E, leg. Uno ROOSILEHT, 2 ♀♀; 24.4.2006, Malaysia, Cameron Highlands, 600 m, 4°22'09''N, 101°20'00''E, leg. Uno ROOSILEHT, 1 ♂.

Distribution: India, Sikkim, Nepal, Bangladesh, Myanmar, Malaysian Peninsular, Borneo, Sumatra, Sumbawa, Sulawesi, Jawa, Bali, Thailand, Laos, Cambodia, Vietnam, Palawan, Leyte, Tawitawi, Mindanao, Papua New Guinea (Western Province), China (Guangdong, Hainan, Hong Kong, Shanghai, Yunnan).

***Cylindera (Ifasina) discreta discreta* (Schaum, 1863)**

Specimen examined: 26.9.2006, Malaysia, Gua Musang, 5°12'52''N, 101°55'54''E, leg. Uno ROOSILEHT, 1 ♀.

Distribution: Malaysian Peninsular, Borneo (Brunei, Sarawak, Sabah), Sumatra (Aceh, S. utara, S. barat, Bengkulu, S. selatan), Flores, Sebesi, Jawa, Sulawesi, Moluccas Toekang Besi Is., Buru, Australia (Northern Territory, N Queensland).

***Callytron doriai* (Horn, 1897)**

(Fig. 1)

Specimens examined: 7.4.2006, Malaysia, Mercung, 3°00'47''N, 103°25'36''E, leg. Uno ROOSILEHT, 4 ♀♀.

Distribution: Borneo (Sarawak), Sumatra (S. utara).

New record for the Malaysian Peninsular.

***Enantiola hewittii* (Horn, 1908)**

(Fig. 2)

Specimen examined: 7.4.2006, Malaysia, Mercung, 3°00'47''N, 103°25'36''E, leg. Uno ROOSILEHT, 1 ♀.

Distribution: Borneo (Sarawak), Vietnam, Cambodia.

New record for the Malaysian Peninsular.




Fig. 1: *Callytron doriai* (Horn, 1897)

Fig. 2: *Enantiola hewittii* (Horn, 1908)

Acknowledgement

The author is indebted to Uno ROOSILEHT for making the material available for the study.

Author:

Jürgen WIESNER, Dresdener Ring 11, D-38444 Wolfsburg, Germany.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Mitteilungen des Internationalen Entomologischen Vereins](#)

Jahr/Year: 2007

Band/Volume: [32 2007](#)

Autor(en)/Author(s): Wiesner Jürgen

Artikel/Article: [New Records of Tiger Beetle Species from the Malaysian Peninsular \(II\) \(Coleoptera: Cicindelidae\) 111-116](#)